

COMPUTATIONAL STATISTICS

PRIMER ON PROBABILITY AND STATISTICS

Luca Bortolussi

Department of Mathematics and Geosciences
University of Trieste

Office 303, third floor, E1 3
`luca@dmi.units.it`

Trieste, Winter Semester 2015/2016

OUTLINE

- 1 BASICS OF PROBABILITY THEORY
- 2 SOME PROBABILITY DISTRIBUTIONS
- 3 FITTING DISTRIBUTIONS
- 4 INFORMATION THEORY
- 5 DECISION THEORY

BASICS ON PROBABILITIES

SIGMA ALGEBRAS

Let Ω be a set, $\mathcal{S} \subseteq 2^\Omega$ is a σ -algebra iff

$$1) \emptyset \in \mathcal{S}$$

$$2) A \in \mathcal{S} \Rightarrow A^c \in \mathcal{S}$$

$$3) A_n \in \mathcal{S} \Rightarrow \bigcup_n A_n \in \mathcal{S}$$

BASICS ON PROBABILITIES

SIGMA ALGEBRAS

Let Ω be a set, $\mathcal{S} \subseteq 2^\Omega$ is a σ -algebra iff

- 1 $\emptyset, \Omega \in \mathcal{S}$;
- 2 $A \in \mathcal{S} \Rightarrow A^c \in \mathcal{S}$;
- 3 $A_n \in \mathcal{S}, n \in \mathbb{N} \Rightarrow \bigcup_n A_n \in \mathcal{S}$;

(Ω, \mathcal{S}) is called *measurable space*. Example: the Borel sigma algebra \mathcal{B} in \mathbb{R}^n , the smallest σ -algebra containing all open sets.

MEASURABLE FUNCTION

A function $f : (X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$ is measurable

BASICS ON PROBABILITIES

SIGMA ALGEBRAS

Let Ω be a set, $\mathcal{S} \subseteq 2^\Omega$ is a σ -algebra iff

- 1 $\emptyset, \Omega \in \mathcal{S}$;
- 2 $A \in \mathcal{S} \Rightarrow A^c \in \mathcal{S}$;
- 3 $A_n \in \mathcal{S}, n \in \mathbb{N} \Rightarrow \bigcup_n A_n \in \mathcal{S}$;

(Ω, \mathcal{S}) is called *measurable space*. Example: the Borel sigma algebra \mathcal{B} in \mathbb{R}^n , the smallest σ -algebra containing all open sets.

MEASURABLE FUNCTION

A function $f : (X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$ is measurable iff $f^{-1}(B) \in \mathcal{A}$ for each $B \in \mathcal{B}$

BASICS ON PROBABILITIES

PROBABILITY MEASURE

Let (Ω, \mathcal{S}) be a measurable space. A probability measure on (Ω, \mathcal{S}) is a function $\mu : \mathcal{S} \rightarrow [0, 1]$ such that

$$\mu(\Omega) = 1$$

$$A_n \text{ disjoint} \Rightarrow \mu\left(\bigcup_n A_n\right) = \sum \mu(A_n)$$

BASICS ON PROBABILITIES


PROBABILITY MEASURE

Let (Ω, \mathcal{S}) be a measurable space. A probability measure on (Ω, \mathcal{S}) is a function $\mu : \mathcal{S} \rightarrow [0, 1]$ such that

- 1 $\mu(\emptyset) = 0$
- 2 $\mu(A^c) = 1 - \mu(A)$
- 3 If $A_n \in \mathcal{S}$ disjoint, then $\mu(\bigcup_n A_n) = \sum_{n=0}^{\infty} \mu(A_n)$

PROBABILITY SPACE

$(\Omega, \mathcal{S}, \mu)$, with \mathcal{S} σ -algebra and μ probability measure on (Ω, \mathcal{S}) , is a probability space.



PROBABILITIES: AN INTUITION

- Random variables: results of non exactly reproducible experiments
- Either intrinsically random (e.g. quantum mechanics) or the system is incompletely known, cannot be controlled precisely

PROBABILITIES: AN INTUITION

- Random variables: results of non exactly reproducible experiments
- Either intrinsically random (e.g. quantum mechanics) or the system is incompletely known, cannot be controlled precisely
- The probability p_i of an experiment taking a certain value i is the frequency with which that value is taken in the limit of infinite experimental trials (**frequentist viewpoint**)

PROBABILITIES: AN INTUITION

- Random variables: results of non exactly reproducible experiments
- Either intrinsically random (e.g. quantum mechanics) or the system is incompletely known, cannot be controlled precisely
- The probability p_i of an experiment taking a certain value i is the frequency with which that value is taken in the limit of infinite experimental trials (**frequentist viewpoint**)
- Alternatively, we can take probability to be our belief that a certain value will be taken (**Bayesian viewpoint**)

RANDOM VARIABLES: FORMALLY

- Let $(\Omega, \mathcal{S}, \mu)$ be a probability space (the **sample space**) and (X, \mathcal{A}) be a measurable space.

$X: (\Omega, \mathcal{S}) \rightarrow (X, \mathcal{A})$ measurable.

$$P(X \in A) = \mu(X^{-1}(A))$$

$A \in \mathcal{A}$

RANDOM VARIABLES: FORMALLY

- Let $(\Omega, \mathcal{S}, \mu)$ be a probability space (the **sample space**) and $(\mathcal{X}, \mathcal{A})$ be a measurable space. A measurable function $x : (\Omega, \mathcal{S}) \rightarrow (\mathcal{X}, \mathcal{A})$ is called a **random variable**.
- The law of x

RANDOM VARIABLES: FORMALLY

- Let $(\Omega, \mathcal{S}, \mu)$ be a probability space (the **sample space**) and $(\mathcal{X}, \mathcal{A})$ be a measurable space. A measurable function $x : (\Omega, \mathcal{S}) \rightarrow (\mathcal{X}, \mathcal{A})$ is called a **random variable**.
- The law of x is $\mathbb{P}\{x \in A\} = \mu(x^{-1}(A))$, for each $A \in \mathcal{A}$, and it is a probability distribution in $(\mathcal{X}, \mathcal{A})$.
- Example: discrete random variables, with values in a countable state space S , with the σ -algebra 2^S .

RANDOM VARIABLES: FORMALLY

- Let $(\Omega, \mathcal{S}, \mu)$ be a probability space (the **sample space**) and $(\mathcal{X}, \mathcal{A})$ be a measurable space. A measurable function $x : (\Omega, \mathcal{S}) \rightarrow (\mathcal{X}, \mathcal{A})$ is called a **random variable**.
- The law of x is $\mathbb{P}\{x \in A\} = \mu(x^{-1}(A))$, for each $A \in \mathcal{A}$, and it is a probability distribution in $(\mathcal{X}, \mathcal{A})$.
- Example: discrete random variables, with values in a countable state space \mathcal{S} , with the σ -algebra $2^{\mathcal{S}}$.
- Example: real-valued random variables, with values in \mathbb{R} , with the Borel σ -algebra.

RANDOM VARIABLES, NOTATION AND MORE DEFINITIONS

- Let x and y be two random variables, $p(x = i, y = j)$ is the *joint probability* of x taking value i and y taking value j (with i and j in the respective spaces of possible values).
- Often just written $p(x, y)$ to indicate the function (as opposed to its evaluation over the outcomes i and j).
- $p(x|y)$ is the conditional probability, i.e. the probability of x if you know y has a certain value

RULES OF PROBABILITY (DISCRETE RV)

$x \in \mathcal{X}$ discrete

- **Normalisation**: the sum of the probabilities of all possible experimental outcomes must be 1,

$$\sum_{x \in \mathcal{X}} p(x) = 1$$

RULES OF PROBABILITY (DISCRETE RV)

- **Normalisation**: the sum of the probabilities of all possible experimental outcomes must be 1, $\sum_{x \in \mathcal{X}} p(x) = 1$
- **Sum rule**: the marginal probability $p(x)$ is given by summing the joint $p(x, y)$ over all possible values of y ,

$$p(x) = \sum_{y \in \mathcal{Y}} p(x, y)$$

RULES OF PROBABILITY (DISCRETE RV)

- **Normalisation**: the sum of the probabilities of all possible experimental outcomes must be 1, $\sum_{x \in \mathcal{X}} p(x) = 1$
- **Sum rule**: the marginal probability $p(x)$ is given by summing the joint $p(x, y)$ over all possible values of y ,

$$p(x) = \sum_{y \in \mathcal{Y}} p(x, y)$$

- **Product rule**: the joint is the product of the conditional and the marginal,

$$p(x, y) = p(x|y)p(y)$$

RULES OF PROBABILITY (DISCRETE RV)

- **Normalisation**: the sum of the probabilities of all possible experimental outcomes must be 1, $\sum_{x \in \mathcal{X}} p(x) = 1$
- **Sum rule**: the marginal probability $p(x)$ is given by summing the joint $p(x, y)$ over all possible values of y ,

$$p(x) = \sum_{y \in \mathcal{Y}} p(x, y)$$

- **Product rule**: the joint is the product of the conditional and the marginal, $p(x, y) = p(x|y)p(y)$
- **Bayes rule**: the posterior is the ratio of the joint and the marginal

$$\leadsto p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

RULES OF PROBABILITY (DISCRETE RV)

- **Normalisation**: the sum of the probabilities of all possible experimental outcomes must be 1, $\sum_{x \in \mathcal{X}} p(x) = 1$
- **Sum rule**: the marginal probability $p(x)$ is given by summing the joint $p(x, y)$ over all possible values of y ,

$$p(x) = \sum_{y \in \mathcal{Y}} p(x, y)$$

- **Product rule**: the joint is the product of the conditional and the marginal, $p(x, y) = p(x|y)p(y)$
- **Bayes rule**: the posterior is the ratio of the joint and the marginal

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

INDEPENDENCE

- Two random variables x and y are *independent* if their joint probability factorises in terms of marginals

$$P(x, y) = P(x)P(y)$$

INDEPENDENCE

- Two random variables x and y are *independent* if their joint probability factorises in terms of marginals

$$p(x, y) = p(x)p(y)$$

- Using the product rule, this is equivalent to the conditional being equal to the marginal

$$x, y \text{ ind} \Rightarrow P(x|y) = P(x)$$

INDEPENDENCE

- Two random variables x and y are *independent* if their joint probability factorises in terms of marginals

$$p(x, y) = p(x)p(y)$$

- Using the product rule, this is equivalent to the conditional being equal to the marginal

$$p(x, y) = p(x)p(y) \Leftrightarrow p(x|y) = p(x)$$

CONTINUOUS STATES

- If the state space \mathcal{X} is continuous some of the previous definitions must be modified
- The general case is mathematically difficult; we restrict ourselves to $\mathcal{X} = \mathbb{R}^n$ and to distributions which admit a *density*, i.e. a function

$$p: \mathcal{X} \rightarrow \mathbb{R}, p(x) \geq 0, \int_{\mathcal{X}} p(x) dx = 1$$

$$p(A) = \int_A p(x) dx$$

CONTINUOUS STATES

- If the state space \mathcal{X} is continuous some of the previous definitions must be modified
- The general case is mathematically difficult; we restrict ourselves to $\mathcal{X} = \mathbb{R}^n$ and to distributions which admit a *density*, i.e. a function

$$p : \mathcal{X} \rightarrow \mathbb{R} \quad \text{s.t.} \quad p(x) \geq 0, \forall x \quad \text{and} \quad \int_{\mathcal{X}} p(x) dx = 1$$

- Formally, these are absolute continuous measures with respect to the Lebesgue measure, with density function playing the role of a Radon-Nikodym derivative.

CONTINUOUS STATES

- If the state space \mathcal{X} is continuous some of the previous definitions must be modified
- The general case is mathematically difficult; we restrict ourselves to $\mathcal{X} = \mathbb{R}^n$ and to distributions which admit a *density*, i.e. a function

$$p : \mathcal{X} \rightarrow \mathbb{R} \quad \text{s.t.} \quad p(x) \geq 0, \forall x \quad \text{and} \quad \int_{\mathcal{X}} p(x) dx = 1$$

- Formally, these are absolute continuous measures with respect to the Lebesgue measure, with density function playing the role of a Radon-Nikodym derivative.
- It can be shown that the rules of probability distributions hold also for probability densities
- Notice that $p(x)$ is NOT the probability of the random variable being in state x (that is always zero for bounded densities); probabilities are only defined as integrals over subsets of \mathcal{X}

PROBABILISTIC INFERENCE

- In logics, an inference system is given by a set of inference rules, allowing to infer logical consequences from a set of facts/ axioms.

PROBABILISTIC INFERENCE

- In logics, an inference system is given by a set of inference rules, allowing to infer logical consequences from a set of facts/ axioms.
- The rules of probability define an inference system generalising logical ones to reason under uncertainty.

PROBABILISTIC INFERENCE

- In logics, an inference system is given by a set of inference rules, allowing to infer logical consequences from a set of facts/ axioms.
- The rules of probability define an inference system generalising logical ones to reason under uncertainty.
- Typically, we have a probabilistic model, and possibly evidence (e.g. experimental observations), and we want to deduce consequences – here compute probabilities.
- We do this by consistent applications of the rules of probability.

PROBABILISTIC INFERENCE: EXAMPLE

- Scientists¹ found that people that enjoy working 14 hours per day (HW) almost inevitably eat Frico (F):
 $p(F|HW) = 0.8$. The probability of being a HW is rather low, about 10^{-4} .

¹cf. Monon Behaviour

PROBABILISTIC INFERENCE: EXAMPLE

- Scientists¹ found that people that enjoy working 14 hours per day (HW) almost inevitably eat Frico (F):
 $p(F|HW) = 0.8$. The probability of being a HW is rather low, about 10^{-4} .
- Assuming eating Frico is quite common, $p(F) = 0.4$, what is the probability that a Frico eater is a HW?

$$p(HW|F) = \frac{p(F|HW) p(HW)}{p(F)} = \frac{0.8 \cdot 10^{-4}}{0.4} = 2 \cdot 10^{-4}$$

¹cf. Monon Behaviour

PROBABILISTIC INFERENCE: EXAMPLE

- Scientists¹ found that people that enjoy working 14 hours per day (HW) almost inevitably eat Frico (F):
 $p(F|HW) = 0.8$. The probability of being a HW is rather low, about 10^{-4} .
- Assuming eating Frico is quite common, $p(F) = 0.4$, what is the probability that a Frico eater is a HW? By Bayes rule:

$$p(HW|F) = \frac{p(F|HW)p(HW)}{p(F)} = \frac{0.8 \cdot 10^{-4}}{0.4} = 2 \cdot 10^{-4}$$

- As Frico eating is rare worldwide, say $p(F) = \underline{2 \cdot 10^{-4}}$

¹cf. Monon Behaviour

PROBABILISTIC INFERENCE: EXAMPLE

- Scientists¹ found that people that enjoy working 14 hours per day (HW) almost inevitably eat Frico (F):
 $p(F|HW) = 0.8$. The probability of being a HW is rather low, about 10^{-4} .
- Assuming eating Frico is quite common, $p(F) = 0.4$, what is the probability that a Frico eater is a HW? By Bayes rule:

$$p(HW|F) = \frac{p(F|HW)p(HW)}{p(F)} = \frac{0.8 \cdot 10^{-4}}{0.4} = 2 \cdot 10^{-4}$$

- As Frico eating is rare worldwide, say $p(F) = 2 \cdot 10^{-4}$

$$p(HW|F) = \frac{p(F|HW)p(HW)}{p(F)} = \frac{0.8 \cdot 10^{-4}}{2 \cdot 10^{-4}} = 0.4$$

¹cf. Monon Behaviour

PROBABILISTIC INFERENCE: EXAMPLE

Example 1.4 (Who's in the bathroom?). Consider a household of three people, Alice, Bob and Cecil. Cecil wants to go to the bathroom but finds it occupied. He then goes to Alice's room and sees she is there. Since Cecil knows that only either Alice or Bob can be in the bathroom, from this he infers that Bob must be in the bathroom.

PROBABILISTIC INFERENCE: EXAMPLE

Example 1.4 (Who's in the bathroom?). Consider a household of three people, Alice, Bob and Cecil. Cecil wants to go to the bathroom but finds it occupied. He then goes to Alice's room and sees she is there. Since Cecil knows that only either Alice or Bob can be in the bathroom, from this he infers that Bob must be in the bathroom.

To arrive at the same conclusion in a mathematical framework, we define the following events

$$\rightarrow A = \text{Alice is in her bedroom}, \quad B = \text{Bob is in his bedroom}, \quad O = \text{Bathroom occupied} \quad (1.2.11)$$

We can encode the information that if either Alice or Bob are not in their bedrooms, then they must be in the bathroom (they might both be in the bathroom) as

$$\rightarrow p(O = \text{tr} | A = \text{fa}, B) = 1, \quad p(O = \text{tr} | A, B = \text{fa}) = 1 \quad (1.2.12)$$

The first term expresses that the bathroom is occupied if Alice is not in her bedroom, wherever Bob is. Similarly, the second term expresses bathroom occupancy as long as Bob is not in his bedroom. Then

$$p(B = \text{fa} | O = \text{tr}, A = \text{tr}) = \frac{p(B = \text{fa}, O = \text{tr}, A = \text{tr})}{p(O = \text{tr}, A = \text{tr})} = \frac{p(O = \text{tr} | A = \text{tr}, B = \text{fa})p(A = \text{tr}, B = \text{fa})}{p(O = \text{tr}, A = \text{tr})} \quad (1.2.13)$$

where

$$p(O = \text{tr}, A = \text{tr}) = p(O = \text{tr} | A = \text{tr}, B = \text{fa})p(A = \text{tr}, B = \text{fa}) + p(O = \text{tr} | A = \text{tr}, B = \text{tr})p(A = \text{tr}, B = \text{tr}) \quad (1.2.14)$$

Using the fact $p(O = \text{tr} | A = \text{tr}, B = \text{fa}) = 1$ and $p(O = \text{tr} | A = \text{tr}, B = \text{tr}) = 0$, which encodes that if Alice is in her room and Bob is not, the bathroom must be occupied, and similarly, if both Alice and Bob are in their rooms, the bathroom cannot be occupied,

$$p(B = \text{fa} | O = \text{tr}, A = \text{tr}) = \frac{p(A = \text{tr}, B = \text{fa})}{p(A = \text{tr}, B = \text{fa})} = 1 \quad (1.2.15)$$

This example is interesting since we are not required to make a full probabilistic model in this case thanks to the limiting nature of the probabilities (we don't need to specify $p(A, B)$). The situation is common in limiting situations of probabilities being either 0 or 1, corresponding to traditional logic systems.

PROBABILISTIC INFERENCE: EXAMPLE

Exercise 1.3 (Adapted from [181]). *There are two boxes. Box 1 contains three red and five white balls and box 2 contains two red and five white balls. A box is chosen at random $p(\text{box} = 1) = p(\text{box} = 2) = 0.5$ and a ball chosen at random from this box turns out to be red. What is the posterior probability that the red ball came from box 1?*

$$\begin{aligned}
 P(\text{Box}=1 | B=\text{red}) &= \frac{P(B=\text{red} | \text{Box}=1) \cdot P(\text{Box}=1)}{P(B=\text{red})} \\
 &= \frac{\frac{3}{8} \cdot \frac{1}{2}}{\frac{3}{8} \cdot \frac{1}{2} + \frac{2}{7} \cdot \frac{1}{2}} \\
 &= \frac{3}{31}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\sum_i P(B=\text{red}, \text{Box}=i)}{P(B=\text{red} | \text{Box}=1) \cdot P(\text{Box}=1) + P(B=\text{red} | \text{Box}=2) \cdot P(\text{Box}=2)}
 \end{aligned}$$

DISTRIBUTIONS AND EXPECTATIONS

- A probability distribution for finite state space can be given by a table, in general is given by a functional form
- Probability distributions (over numerical objects) are useful to compute expectations of functions

DISTRIBUTIONS AND EXPECTATIONS

- A probability distribution for finite state space can be given by a table, in general is given by a functional form
- Probability distributions (over numerical objects) are useful to compute expectations of functions

$$\langle f \rangle = \sum_{x \in \mathcal{X}} f(x)p(x)$$

$$\langle x \rangle = \sum_{x \in \mathcal{X}} x \cdot p(x)$$

- Important expectations are the *mean* $\langle x \rangle$ and *variance* $\sim \text{var}(x) = \langle (x - \langle x \rangle)^2 \rangle$.

DISTRIBUTIONS AND EXPECTATIONS

- A probability distribution for finite state space can be given by a table, in general is given by a functional form
- Probability distributions (over numerical objects) are useful to compute expectations of functions

$$\langle f \rangle = \sum_{x \in \mathcal{X}} f(x)p(x)$$

- Important expectations are the *mean* $\langle x \rangle$ and *variance* $\text{var}(x) = \langle (x - \langle x \rangle)^2 \rangle$.
- For more variables, also the *covariance* $\text{cov}(x, y) = \langle (x - \langle x \rangle)(y - \langle y \rangle) \rangle$ or its scaled relative the *correlation* $\text{corr}(x, y) = \text{cov}(x, y) / \sqrt{\text{var}(x)\text{var}(y)}$

DISTRIBUTIONS AND EXPECTATIONS

EXERCISE

if two variables are independent, then their correlation is zero.
NOT TRUE viceversa (no correlation does not imply independence)

COMPUTING EXPECTATIONS

- If you know analytically the probability distribution and can compute the sums (integrals), no problem

sc X is continuous

$$\Rightarrow \langle f \rangle = \int f(x) p(x) dx$$

COMPUTING EXPECTATIONS

- If you know analytically the probability distribution and can compute the sums (integrals), no problem
- If you know the distribution but cannot compute the sums (integrals), enter the magical realm of approximate inference (fun but out of scope)

COMPUTING EXPECTATIONS

- If you know analytically the probability distribution and can compute the sums (integrals), no problem
- If you know the distribution but cannot compute the sums (integrals), enter the magical realm of approximate inference (fun but out of scope)
- If you know nothing but have N_S samples, then use a sample approximation
- Approximate the probability of an outcome with the *frequency* in the sample

COMPUTING EXPECTATIONS

- If you know analytically the probability distribution and can compute the sums (integrals), no problem
- If you know the distribution but cannot compute the sums (integrals), enter the magical realm of approximate inference (fun but out of scope)
- If you know nothing but have N_S samples, then use a sample approximation
- Approximate the probability of an outcome with the *frequency* in the sample

$$\langle f(x) \rangle \simeq \sum_x \frac{n_x}{N_S} f(x) = \frac{1}{N_S} \sum_{i=1}^{N_S} f(x_i)$$

$x_1, \dots, x_{N_S} \sim p(x)$

OUTLINE

- 1 BASICS OF PROBABILITY THEORY
- 2 SOME PROBABILITY DISTRIBUTIONS
- 3 FITTING DISTRIBUTIONS
- 4 INFORMATION THEORY
- 5 DECISION THEORY

BASIC DISCRETE DISTRIBUTIONS

DISCRETE/ CATEGORICAL DISTRIBUTION

a random variable can take N distinct values with probability p_i , $i = 1, \dots, N$. Formally

$$p(x = i) = p_i \quad \sum_{i=1}^N p_i = 1$$

BASIC DISCRETE DISTRIBUTIONS

BERNOULLI DISTRIBUTION

a discrete random variable x with two outcomes: 1, with probability $p(x = 1|\theta) = \theta$ and 0, with probability $1 - \theta$. Compute mean and variance.

BASIC DISCRETE DISTRIBUTIONS

BERNOULLI DISTRIBUTION

a discrete random variable x with two outcomes: 1, with probability $p(x = 1|\theta) = \theta$ and 0, with probability $1 - \theta$. Compute mean and variance.

- $\langle x \rangle = \theta$
- $\text{var}(x) = \theta(1 - \theta)$

BASIC DISCRETE DISTRIBUTIONS

BINOMIAL DISTRIBUTION

describes the outcome of n Bernoulli trials. The probability of k successes is

BASIC DISCRETE DISTRIBUTIONS

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

BINOMIAL DISTRIBUTION

describes the outcome of n Bernoulli trials. The probability of k successes is

$$p(y = k|\theta) = \binom{n}{k} \theta^k (1-\theta)^{n-k}$$

$\leftarrow k! \right.$
 $\leftarrow 010-11 \dots 0$
 \leftarrow

Compute mean and variance.

$\langle \rangle$ is a linear operator

$$\langle aX + bY \rangle = a\langle X \rangle + b\langle Y \rangle$$

$$Y \sim \text{Binomial} \quad Y = X_1 + \dots + X_n$$

$\leftarrow X_i \sim \text{Bernoulli}$
 $P(X_i = 1) = \theta$

$$\langle Y \rangle = \langle X_1 + \dots + X_n \rangle = \sum_i \langle X_i \rangle = n \cdot \theta$$

BASIC DISCRETE DISTRIBUTIONS


BINOMIAL DISTRIBUTION

describes the outcome of n Bernoulli trials. The probability of k successes is

$$p(y = k|\theta) = \binom{n}{k} \theta^k (1 - \theta)^{n-k}$$

Compute mean and variance.

- $\langle x \rangle = n\theta$
- $\text{var}(x) = n\theta(1 - \theta)$



BASIC DISCRETE DISTRIBUTIONS

MULTINOMIAL DISTRIBUTION

describes the outcome of n trials of a categorical distribution on $\{1, \dots, K\}$ with probabilities $\theta = (\theta_1, \dots, \theta_K)$. The probability of observing y_i outcomes of type i is

$$p(y_1, \dots, y_K | \theta) = \frac{n!}{y_1! \dots y_K!} \prod_{i=1}^K \theta_i^{y_i}$$

Compute mean and variance.

$$\langle y_i \rangle = n \theta_i$$

BASIC DISCRETE DISTRIBUTIONS

MULTINOMIAL DISTRIBUTION

describes the outcome of n trials of a categorical distribution on $\{1, \dots, K\}$ with probabilities $\theta = (\theta_1, \dots, \theta_K)$. The probability of observing y_i outcomes of type i is

$$p(y_1, \dots, y_K | \theta) = \frac{n!}{y_1! \cdots y_K!} \prod_{i=1}^K \theta_i^{y_i}$$

Compute mean and variance.

- $\langle y_i \rangle = n\theta_i$
- $\text{var}(x) = n\theta_i(1 - \theta_i)$
- $\text{cov}(x) = -n\theta_i\theta_j$

POISSON DISTRIBUTION

a distribution over non-negative integers

POISSON DISTRIBUTION

a distribution over non-negative integers

$$p(n|\lambda) = \frac{\lambda^n}{n!} \exp[-\lambda]$$

The parameter λ is often called the *rate* of the distribution. The Poisson distribution is often used for *rare events*, e.g. decaying of particles or binding of DNA fragments to a probe.
Compute mean and variance.

POISSON DISTRIBUTION


a distribution over non-negative integers

$$p(n|\lambda) = \frac{\lambda^n}{n!} \exp[-\lambda]$$

The parameter λ is often called the *rate* of the distribution. The Poisson distribution is often used for *rare events*, e.g. decaying of particles or binding of DNA fragments to a probe.

Compute mean and variance.

- $\langle x \rangle = \lambda$
- $\text{var}(x) = \lambda$
- A binomial with parameter $\theta = \lambda/n$ converges to a Poisson for $n \rightarrow \infty$



BASIC CONTINUOUS DISTRIBUTIONS

UNIFORM DISTRIBUTION

A variable x with constant density, over its domain of definition, which is an interval $[a, b] \subset \mathbb{R}$.

BASIC CONTINUOUS DISTRIBUTIONS

UNIFORM DISTRIBUTION


A variable x with constant density, over its domain of definition, which is an interval $[a, b] \subset \mathbb{R}$. Hence $p(x) = 1/(b - a)$, if $x \in [a, b]$.

BASIC CONTINUOUS DISTRIBUTIONS

UNIFORM DISTRIBUTION

A variable x with constant density, over its domain of definition, which is an interval $[a, b] \subset \mathbb{R}$. Hence $p(x) = 1/(b - a)$, if $x \in [a, b]$.

- $\langle x \rangle = (a + b)/2$ *↙*
- $\text{var}(x) = (a^2 + b^2 + ab)/3$ *↘*



EXPONENTIAL DISTRIBUTION

Typically used for the time of occurrence of a random event, like the decay of a particle or the arrival of a customer in a shop. For $x \geq 0$,

EXPONENTIAL DISTRIBUTION

Typically used for the time of occurrence of a random event, like the decay of a particle or the arrival of a customer in a shop. For $x \geq 0$,

$$p(x|\lambda) = \lambda e^{-\lambda x}$$

with λ known as the rate of the distribution.


EXPONENTIAL DISTRIBUTION

Typically used for the time of occurrence of a random event, like the decay of a particle or the arrival of a customer in a shop. For $x \geq 0$,

$$p(x|\lambda) = \lambda e^{-\lambda x}$$

with λ known as the rate of the distribution.

- $\langle x \rangle = 1/\lambda$
- $\text{var}(x) = 1/\lambda^2$



GAMMA DISTRIBUTION

Defined for $x \geq 0$, by the density

$$p(x|\alpha, \beta) = \frac{1}{\beta\Gamma(\alpha)} (x/\beta)^{\alpha-1} e^{-x/\beta}$$

where α is the shape parameter, β is the scale parameter, and $\Gamma(\alpha)$ is the gamma function

$$\Gamma(a) = \int_0^{\infty} t^{a-1} e^{-t} dt$$

GAMMA DISTRIBUTION


Defined for $x \geq 0$, by the density

$$p(x|\alpha, \beta) = \frac{1}{\beta\Gamma(\alpha)} (x/\beta)^{\alpha-1} e^{-x/\beta}$$

where α is the shape parameter, β is the scale parameter, and $\Gamma(\alpha)$ is the gamma function

$$\Gamma(a) = \int_0^{\infty} t^{a-1} e^{-t} dt$$

- $\langle x \rangle = \alpha/\beta$
- $\text{var}(x) = \alpha/\beta^2$
- $\text{Gamma}(1, \lambda) \equiv \text{Exp}(\lambda)$



INVERSE GAMMA DISTRIBUTION

Defined for $x \geq 0$, by the density

$$p(x|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} (1/x)^{\alpha+1} e^{-\beta/x}$$

where α is the shape parameter, β is the scale parameter, and $\Gamma(\alpha)$ is the gamma function


INVERSE GAMMA DISTRIBUTION

Defined for $x \geq 0$, by the density

$$p(x|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} (1/x)^{\alpha+1} e^{-\beta/x}$$

where α is the shape parameter, β is the scale parameter, and $\Gamma(\alpha)$ is the gamma function

- $\langle x \rangle = \beta/(\alpha - 1), \alpha > 1$
- $\text{var}(x) = \beta^2/((\alpha - 1)^2(\alpha - 2))$



BETA DISTRIBUTION

Defined for $x \in [0, 1]$ by

$$p(x|\alpha, \beta) = \frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1}$$

where the Beta function $B(\alpha, \beta)$ is defined by $[B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$


BETA DISTRIBUTION

Defined for $x \in [0, 1]$ by

$$p(x|\alpha, \beta) = \frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1}$$

where the Beta function $B(\alpha, \beta)$ is defined by $[B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$

- $\langle x \rangle = \alpha / (\alpha + \beta)$
- $\text{var}(x) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$



DIRICHLET DISTRIBUTION

It is a distribution over discrete probability distributions on k elements. It depends on k parameters $\mathbf{u} = u_1, \dots, u_k$. Its pdf is

$$p_{dir}(\mathbf{x}|\mathbf{u}) = p(\mathbf{x}|\mathbf{u}) = \frac{1}{B(\mathbf{u})} \prod_{i=1}^k x_i^{u_i-1}$$

$$\text{with } B(\mathbf{u}) = \frac{\prod_{i=1}^k \Gamma(u_i)}{\Gamma(\sum_{i=1}^k u_i)}$$


DIRICHLET DISTRIBUTION

It is a distribution over discrete probability distributions on k elements. It depends on k parameters $\mathbf{u} = u_1, \dots, u_k$. Its pdf is

$$p_{dir}(\mathbf{x}|\mathbf{u}) = p(\mathbf{x}|\mathbf{u}) = \frac{1}{B(\mathbf{u})} \prod_{i=1}^k x_i^{u_i-1}$$

with $B(\mathbf{u}) = \frac{\prod_{i=1}^k \Gamma(u_i)}{\Gamma(\sum_{i=1}^k u_i)}$

- $\langle x \rangle = \alpha / (\alpha + \beta)$
- $\text{var}(x) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
- $p_{dir}(\mathbf{x}|\mathbf{u}_1)p_{dir}(\mathbf{x}|\mathbf{u}_2) = p_{dir}(\mathbf{x}|\mathbf{u}_1 + \mathbf{u}_2)$
- $\int_{x_j} p_{dir}(\mathbf{x}|\mathbf{u}) dx_j = p_{dir}(\mathbf{x}_{/j}|\mathbf{u}_{/j})$
- $\int_{\mathbf{x}_{/j}} p_{dir}(\mathbf{x}|\mathbf{u}) d\mathbf{x}_{/j} = p_{beta}(x_j|u_j, \sum_{i \neq j} u_i)$



BASIC CONTINUOUS DISTRIBUTIONS

One of the most used distributions in Machine Learning


$$p(x|\mu, \sigma^2) = \mathcal{N}(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2}(x-\mu)^2}$$

BASIC CONTINUOUS DISTRIBUTIONS

One of the most used distributions in Machine Learning

$$p(x|\mu, \sigma^2) = \mathcal{N}(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2}(x-\mu)^2}$$

- $\langle x \rangle = \mu$
- $\text{var}(x) = \sigma^2$
- For $\mu = 1, \sigma^2 = 1$ we talk about a standard normal distribution.
- $1/\sigma^2$ is known as *precision*.



STUDENT'S T-DISTRIBUTION

$$p(x|\mu, \lambda, \nu) = Student(x|\mu, \lambda, \nu) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})} \left(\frac{\lambda}{\nu\pi}\right)^{\frac{1}{2}} \left[1 + \frac{\lambda(x-\mu)^2}{\nu}\right]^{-\frac{\nu+1}{2}} \quad (8.3.24)$$

where μ is the mean, ν the degrees of freedom, and λ scales the distribution. The variance is given by

$$\text{var}(x) = \frac{\nu}{\lambda(\nu-2)}, \text{ for } \nu > 2 \quad (8.3.25)$$

For $\nu \rightarrow \infty$ the distribution tends to a Gaussian with mean μ and variance $1/\lambda$. As ν decreases the tails of the distribution become fatter.

STUDENT'S T-DISTRIBUTION

$$p(x|\mu, \lambda, \nu) = Student(x|\mu, \lambda, \nu) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})} \left(\frac{\lambda}{\nu\pi}\right)^{\frac{1}{2}} \left[1 + \frac{\lambda(x-\mu)^2}{\nu}\right]^{-\frac{\nu+1}{2}} \quad (8.3.24)$$

where μ is the mean, ν the degrees of freedom, and λ scales the distribution. The variance is given by


$$\text{var}(x) = \frac{\nu}{\lambda(\nu-2)}, \text{ for } \nu > 2 \quad (8.3.25)$$

For $\nu \rightarrow \infty$ the distribution tends to a Gaussian with mean μ and variance $1/\lambda$. As ν decreases the tails of the distribution become fatter.

The t -distribution can be derived from a *scaled mixture*

$$p(x|\mu, a, b) = \int_{\tau=0}^{\infty} \mathcal{N}(x|\mu, \tau^{-1}) \text{Gam}^{is}(\tau|a, b) d\tau$$

With $\nu = 2a$ degrees of freedom and scale $\lambda = a/b$.




MULTIVARIATE NORMAL DISTRIBUTION


This is the most important distribution we will use, and generalises the 1d normal. In d dimensions

$$p(\mathbf{x}|\mu, \Sigma) = \mathcal{N}(\mathbf{x}|\mu, \Sigma) = \frac{1}{\sqrt{(2\pi)^d \det(\Sigma)}} \exp\left(-\frac{1}{2}(\mathbf{x} - \mu)\Sigma^{-1}(\mathbf{x} - \mu)^T\right)$$

It holds $\mu = \langle \mathbf{x} \rangle$, and $\Sigma = \text{cov}(\mathbf{x}, \mathbf{x}) = \langle (\mathbf{x} - \mu)(\mathbf{x} - \mu)^T \rangle$



(a)



(b)

PROPERTIES OF MULTIVARIATE NORMAL

Completing the square

A useful technique in manipulating Gaussians is completing the square. For example, the expression

$$\exp\left(-\frac{1}{2}\mathbf{x}^T\mathbf{A}\mathbf{x} + \mathbf{b}^T\mathbf{x}\right) \quad (8.4.10)$$

can be transformed as follows. First we complete the square:

$$\frac{1}{2}\mathbf{x}^T\mathbf{A}\mathbf{x} - \mathbf{b}^T\mathbf{x} = \frac{1}{2}(\mathbf{x} - \mathbf{A}^{-1}\mathbf{b})^T\mathbf{A}(\mathbf{x} - \mathbf{A}^{-1}\mathbf{b}) - \frac{1}{2}\mathbf{b}^T\mathbf{A}^{-1}\mathbf{b} \quad (8.4.11)$$

Hence

$$\exp\left(-\frac{1}{2}\mathbf{x}^T\mathbf{A}\mathbf{x} + \mathbf{b}^T\mathbf{x}\right) = \mathcal{N}(\mathbf{x}|\mathbf{A}^{-1}\mathbf{b}, \mathbf{A}^{-1})\sqrt{\det(2\pi\mathbf{A}^{-1})}\exp\left(\frac{1}{2}\mathbf{b}^T\mathbf{A}^{-1}\mathbf{b}\right) \quad (8.4.12)$$

$\rho(\mathbf{x}|\mathbf{A}, \mathbf{b}) \propto \exp\left(-\frac{1}{2}\mathbf{x}\mathbf{A}\mathbf{x}^T + \mathbf{b}^T\mathbf{x}\right)$ is known as the canonical representation, and it is normal with mean $\mathbf{A}^{-1}\mathbf{b}$ and covariance \mathbf{A}^{-1} .

Linear transformation


Result 8.3 (Linear Transform of a Gaussian). Let \mathbf{y} be linearly related to \mathbf{x} through

$$\mathbf{y} = \mathbf{M}\mathbf{x} + \boldsymbol{\eta} \quad (8.4.14)$$


where $\mathbf{x} \perp \boldsymbol{\eta}$, $\boldsymbol{\eta} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, and $\mathbf{x} \sim \mathcal{N}(\boldsymbol{\mu}_x, \boldsymbol{\Sigma}_x)$. Then the marginal $p(\mathbf{y}) = \int_{\mathbf{x}} p(\mathbf{y}|\mathbf{x})p(\mathbf{x})$ is a Gaussian

$$p(\mathbf{y}) = \mathcal{N}(\mathbf{y}|\mathbf{M}\boldsymbol{\mu}_x + \boldsymbol{\mu}, \mathbf{M}\boldsymbol{\Sigma}_x\mathbf{M}^T + \boldsymbol{\Sigma}) \quad (8.4.15)$$

PROPERTIES OF MULTIVARIATE NORMAL



(a)



(b)

Eigendecomposition

$$\Sigma = \mathbf{E}\mathbf{\Lambda}\mathbf{E}^T \quad (8.4.5)$$

where $\mathbf{E}^T\mathbf{E} = \mathbf{I}$ and $\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_D)$. In the case of a covariance matrix, all the eigenvalues λ_i are positive. This means that one can use the transformation

$$\mathbf{y} = \mathbf{\Lambda}^{-\frac{1}{2}}\mathbf{E}^T(\mathbf{x} - \boldsymbol{\mu}) \quad (8.4.6)$$

so that

$$(\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) = (\mathbf{x} - \boldsymbol{\mu})^T \mathbf{E}\mathbf{\Lambda}^{-1}\mathbf{E}^T (\mathbf{x} - \boldsymbol{\mu}) = \mathbf{y}^T \mathbf{y} \quad (8.4.7)$$

So by rescaling, we can obtain a product of d -univariate standard normal distributions, one per dimension.

PROPERTIES OF MULTIVARIATE NORMAL

Marginal and conditional of multivariate Gaussians

Result 8.4 (Partitioned Gaussian). Consider a distribution $\mathcal{N}(\mathbf{z}|\boldsymbol{\mu}, \boldsymbol{\Sigma})$ defined jointly over two vectors \mathbf{x} and \mathbf{y} of potentially differing dimensions,

$$\mathbf{z} = \begin{pmatrix} \mathbf{x} \\ \mathbf{y} \end{pmatrix} \tag{8.4.16}$$

with corresponding mean and partitioned covariance

$$\boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_x \\ \boldsymbol{\mu}_y \end{pmatrix} \quad \boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{xx} & \boldsymbol{\Sigma}_{xy} \\ \boldsymbol{\Sigma}_{yx} & \boldsymbol{\Sigma}_{yy} \end{pmatrix} \tag{8.4.17}$$

where $\boldsymbol{\Sigma}_{yx} \equiv \boldsymbol{\Sigma}_{xy}^\top$. The marginal distribution is given by

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_x, \boldsymbol{\Sigma}_{xx}) \tag{8.4.18}$$

and conditional

$$p(\mathbf{x}|\mathbf{y}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_x + \boldsymbol{\Sigma}_{xy}\boldsymbol{\Sigma}_{yy}^{-1}(\mathbf{y} - \boldsymbol{\mu}_y), \boldsymbol{\Sigma}_{xx} - \boldsymbol{\Sigma}_{xy}\boldsymbol{\Sigma}_{yy}^{-1}\boldsymbol{\Sigma}_{yx}) \tag{8.4.19}$$

PROPERTIES OF MULTIVARIATE NORMAL

Product of multivariate Gaussians

Result 8.2 (Product of two Gaussians). The product of two Gaussians is another Gaussian, with a multiplicative factor, exercise(8.35):

$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_1, \boldsymbol{\Sigma}_1)\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_2, \boldsymbol{\Sigma}_2) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) \frac{\exp\left(-\frac{1}{2}(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)^\top \mathbf{S}^{-1}(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)\right)}{\sqrt{\det(2\pi\mathbf{S})}} \quad (8.4.8)$$

where $\mathbf{S} \equiv \boldsymbol{\Sigma}_1 + \boldsymbol{\Sigma}_2$ and the mean and covariance are given by

$$\boldsymbol{\mu} = \boldsymbol{\Sigma}_1 \mathbf{S}^{-1} \boldsymbol{\mu}_2 + \boldsymbol{\Sigma}_2 \mathbf{S}^{-1} \boldsymbol{\mu}_1 \quad \boldsymbol{\Sigma} = \boldsymbol{\Sigma}_1 \mathbf{S}^{-1} \boldsymbol{\Sigma}_2 \quad (8.4.9)$$

Gaussian average of a quadratic function

Result 8.5 (Gaussian average of a quadratic function).

$$\left\langle \mathbf{x}^\top \mathbf{A} \mathbf{x} \right\rangle_{\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma})} = \boldsymbol{\mu}^\top \mathbf{A} \boldsymbol{\mu} + \text{trace}(\mathbf{A} \boldsymbol{\Sigma})$$

THE CURSE OF DIMENSIONALITY

Exercise 1: Suppose you want to explore uniformly a region by gridding it. How many grid points do you need?

THE CURSE OF DIMENSIONALITY

Exercise II: Suppose you sample from a uniform distribution in d dimensions. What is the probability of finding a point inside the region $[\epsilon, 1 - \epsilon]^d$?

THE CURSE OF DIMENSIONALITY

Exercise III: Suppose you sample from a spherical Gaussian distribution. Where do the points lie as the dimensions increase?

MIXTURES: HOW TO BUILD MORE DISTRIBUTIONS

- More general distributions can be built via mixtures: e.g.

$$p(x|\mu_{1,\dots,n}, \sigma_{1,\dots,n}^2) = \sum_i \pi_i \mathcal{N}(\mu_i, \sigma_i^2)$$

where the *mixing coefficients* π_i are discretely distributed

MIXTURES: HOW TO BUILD MORE DISTRIBUTIONS

- More general distributions can be built via mixtures: e.g.

$$p(x|\mu_{1,\dots,n}, \sigma_{1,\dots,n}^2) = \sum_i \pi_i \mathcal{N}(\mu_i, \sigma_i^2)$$

where the *mixing coefficients* π_i are discretely distributed

- You can interpret this as a two stage hierarchical process: choose one component out of a discrete distribution, then choose the distribution for that component

MIXTURES: HOW TO BUILD MORE DISTRIBUTIONS

- **IMPORTANT CONCEPT:** the mixture

$$p(x|\mu_{1,\dots,n}, \sigma_{1,\dots,n}^2) = \sum_i \pi_i \mathcal{N}(\mu_i, \sigma_i^2)$$

is an example of *latent variable model*, with a latent class variable and an observed continuous value. The mixture is the marginal distribution for the observations (w.r.t. the latent variable)

MIXTURES: HOW TO BUILD MORE DISTRIBUTIONS

- **IMPORTANT CONCEPT:** the mixture

$$p(x|\mu_{1,\dots,n}, \sigma_{1,\dots,n}^2) = \sum_i \pi_i \mathcal{N}(\mu_i, \sigma_i^2)$$

is an example of *latent variable model*, with a latent class variable and an observed continuous value. The mixture is the marginal distribution for the observations (w.r.t. the latent variable)

- The probability of the latent variables given the observations can be obtained using Bayes' theorem.

CONTINUOUS MIXTURES: SOME COOL DISTRIBUTIONS

- No need for the mixing distribution (latent variable) to be discrete

CONTINUOUS MIXTURES: SOME COOL DISTRIBUTIONS

- No need for the mixing distribution (latent variable) to be discrete
- Suppose you are interested in the means of normally distributed samples (possibly with different variances/precisions): Marginalising the precision in a Gaussian using a Gamma mixing distribution yields a *Student t-distribution*

CONTINUOUS MIXTURES: SOME COOL DISTRIBUTIONS

- No need for the mixing distribution (latent variable) to be discrete
- Suppose you are interested in the means of normally distributed samples (possibly with different variances/precisions): Marginalising the precision in a Gaussian using a Gamma mixing distribution yields a *Student t-distribution*
- Suppose you have multiple rare event processes happening with slightly different rates: Marginalising the rate in a Poisson distribution using a Gamma mixing distribution yields a *negative binomial* distribution

OUTLINE

- 1 BASICS OF PROBABILITY THEORY
- 2 SOME PROBABILITY DISTRIBUTIONS
- 3 FITTING DISTRIBUTIONS
- 4 INFORMATION THEORY
- 5 DECISION THEORY

PARAMETERS?

- Many distributions are written as conditional probabilities *given* the parameters: $p(x|\theta)$
- Often the values of the parameters are not known
- If we have observations, we can try to estimate the parameters from such data.
- We assume to have independent and identically distributed (i.i.d.) observations of $p(x|\theta_{true})$: $\mathbf{x} = x_1, \dots, x_N$.

MAXIMUM LIKELIHOOD

- Likelihood for i.i.d. observations $\mathbf{x} = x_1, \dots, x_N$:

$$p(\mathbf{x}|\theta) = \prod_{i=1}^N p(x_i|\theta)$$

MAXIMUM LIKELIHOOD

- Likelihood for i.i.d. observations $\mathbf{x} = x_1, \dots, x_N$:

$$p(\mathbf{x}|\theta) = \prod_{i=1}^N p(x_i|\theta)$$

- Choose the parameters that best explain the observations: we pick θ by maximum likelihood:

$$\hat{\theta} = \operatorname{argmax}_{\theta} \left[\prod_i p(x_i|\theta) \right]$$

MAXIMUM A POSTERIORI

- Suppose we can encode prior knowledge (or absence of it) in a **prior** distribution over parameters, $p(\theta)$.
- We can then compute the posterior distribution, given i.i.d. observations $\mathbf{x} = x_1, \dots, x_N$, by Bayes theorem:

$$p(\theta|\mathbf{x}) = \frac{p(\mathbf{x}|\theta)p(\theta)}{p(\mathbf{x})}$$

where

$$p(\mathbf{x}) = \int_{\theta} p(\mathbf{x}|\theta)p(\theta)d\theta$$

MAXIMUM A POSTERIORI

- Suppose we can encode prior knowledge (or absence of it) in a **prior** distribution over parameters, $p(\theta)$.
- We can then compute the posterior distribution, given i.i.d. observations $\mathbf{x} = x_1, \dots, x_N$, by Bayes theorem:

$$p(\theta|\mathbf{x}) = \frac{p(\mathbf{x}|\theta)p(\theta)}{p(\mathbf{x})}$$

where

$$p(\mathbf{x}) = \int_{\theta} p(\mathbf{x}|\theta)p(\theta)d\theta$$

- Estimate θ_{true} by the *maximum a posteriori (MAP)* estimate

$$\hat{\theta}_{MAP} = \operatorname{argmax}_{\theta} \left[p(\theta) \prod_i p(x_i|\theta) \right]$$

EXERCISE: FITTING A DISCRETE DISTRIBUTION

- We have a discrete distribution with values in $K = \{1, \dots, k\}$, with parameters $\mu = \mu_1, \dots, \mu_k$, $\sum_i \mu_i = 1$.
- We have independent observations $\mathbf{x} = x_1, \dots, x_N$, each taking values in K .
- The likelihood is

$$\mathcal{L}(\mu) = p(\mathbf{x}|\mu) = \prod_{i=1}^N p(x_i|\mu)$$

- Compute the Maximum Likelihood estimate of μ . What is the intuitive meaning of the result? What happens if one of the D values is not represented in your sample?

EXERCISE: FITTING A DISCRETE DISTRIBUTION

EXERCISE II: FITTING A GAUSSIAN DISTRIBUTION

We have independent, real valued observations $\mathbf{x} = x_1, \dots, x_N$.
Fit a Gaussian by maximum likelihood.

BAYESIAN ESTIMATION

- The Bayesian approach fully quantifies uncertainty
- The parameters are treated as additional random variables with their own *prior* distribution $p(\theta)$
- The observation likelihood is combined with the prior to obtain a *posterior* distribution via Bayes' theorem

$$p(\theta|x) = \frac{p(x|\theta)p(\theta)}{p(x)}$$

- The distribution of the observable x (*predictive* distribution) is obtained as

$$p(x|x) = \int p(x|\theta)p(\theta|x)d\theta$$

EXERCISE: BAYESIAN FITTING OF GAUSSIANS

- Let data $x_i \quad i = 1, \dots, N$ be distributed according to a Gaussian with mean μ and variance σ^2
- Let the prior distribution over the mean μ be a Gaussian with mean m and variance v^2
- Compute the posterior and predictive distribution

ESTIMATORS

- A procedure to calculate an expectation is called an *estimator*
- e.g., fitting a Gaussian to data by maximum likelihood provides the M.L. estimator for mean and variance, or Bayesian posterior mean
- An estimator will be a noisy estimate of the true value, due to finite sample effects
- An estimator \hat{f} is *unbiased* if its expectation (under the joint distribution of the data set) coincides with the true value
- The ML estimator of variance for a gaussian is biased:

$$\langle \hat{\sigma}^2 \rangle = \frac{N-1}{N} \sigma^2.$$

CONJUGATE PRIORS

- The Bayesian way has advantages in that it quantifies uncertainty and regularizes naturally
- BUT computing the normalisation in Bayes theorem is very hard
- The case when it is possible is when the prior and the posterior are of the same form (*conjugate*)
- Example + Exercise: Binomial and Beta.
- Example: discrete and Dirichlet
- Exercise: conjugate priors for the univariate normal (mean)

CONJUGATE PRIORS: BINOMIAL AND BETA

Show that the Beta is the conjugate prior for the Binomial distribution.

BASICS PROBABILITY SUMMARY

- Probabilities make random variables quantitative
- Basic rules (product and sum) link various probabilities and allow to define the concept of independence
- Basic distributions are worth being familiar with: Gaussian, Gamma, Poisson
- Expectations and parameters can be computed from samples

OUTLINE

- 1 BASICS OF PROBABILITY THEORY
- 2 SOME PROBABILITY DISTRIBUTIONS
- 3 FITTING DISTRIBUTIONS
- 4 INFORMATION THEORY
- 5 DECISION THEORY

ENTROPY AND DIVERGENCE

- Probability theory is the basis of information theory (interesting, but not the topic of this course).
- An important quantity is the *entropy* of a distribution

ENTROPY AND DIVERGENCE

- Probability theory is the basis of information theory (interesting, but not the topic of this course).
- An important quantity is the *entropy* of a distribution

$$H[p] = - \sum_i p_i \log_2 p_i$$

- Entropy measures the level of disorder of a distribution; for discrete distributions, it is always ≥ 0 and 0 only for deterministic distributions

ENTROPY AND DIVERGENCE

- Probability theory is the basis of information theory (interesting, but not the topic of this course).
- An important quantity is the *entropy* of a distribution

$$H[p] = - \sum_i p_i \log_2 p_i$$

- Entropy measures the level of disorder of a distribution; for discrete distributions, it is always ≥ 0 and 0 only for deterministic distributions
- The *relative entropy* or *Kullback-Leibler (KL) divergence* between two distributions is

ENTROPY AND DIVERGENCE

- Probability theory is the basis of information theory (interesting, but not the topic of this course).
- An important quantity is the *entropy* of a distribution

$$H[p] = - \sum_i p_i \log_2 p_i$$

- Entropy measures the level of disorder of a distribution; for discrete distributions, it is always ≥ 0 and 0 only for deterministic distributions
- The *relative entropy* or *Kullback-Leibler (KL) divergence* between two distributions is

$$KL[q||p] = \sum_i q_i \log \frac{q_i}{p_i}$$

- Fact: KL is convex and ≥ 0

JUSTIFICATION FOR MAXIMUM LIKELIHOOD

- Given a data set $\{x_i\}$, $i = 1, \dots, N$, let the empirical distribution be

$$p_{emp}(x) = \frac{1}{N} \sum_{i=1}^N \mathbb{I}(x_i)$$

with \mathbb{I} the indicator function of a set

- To find a suitable distribution q to model the data, one may wish to minimize the Kullback-Leibler divergence

$$KL[p_{emp}||q] = H[p_{emp}] - \langle \log q(x) \rangle_{p_{emp}} = -\frac{1}{N} \sum \log q(x_i)$$

- Maximum likelihood is equivalent to minimizing a KL divergence with the empirical distribution*

OUTLINE

- 1 BASICS OF PROBABILITY THEORY
- 2 SOME PROBABILITY DISTRIBUTIONS
- 3 FITTING DISTRIBUTIONS
- 4 INFORMATION THEORY
- 5 DECISION THEORY

XXXX

???