

Journal of Medicinal Chemistry

© Copyright 2007 by the American Chemical Society

Volume 50, Number 18

September 6, 2007

Letters

2-Amino-3,4-dihydroquinazolines as Inhibitors of BACE-1 (β -Site APP Cleaving Enzyme): Use of Structure Based Design to Convert a Micromolar Hit into a Nanomolar Lead

Ellen W. Baxter,^{*,§} Kelly A. Conway,[§] Ludo Kennis,[†] François Bischoff,[†] Marc H. Mercken,[†] Hans L. De Winter,[†] Charles H. Reynolds,[§] Brett A. Tounge,[§] Chi Luo,[§] Malcolm K. Scott,[§] Yifang Huang,[§] Mirielle Braeken,[†] Serge M. A. Pieters,[†] Didier J. C. Berthelot,[†] Stefan Masure,[†] Wouter D. Bruinzeel,[†] Alfonso D. Jordan,[§] Michael H. Parker,[§] Robert E. Boyd,[§] Junya Qu,[§] Richard S. Alexander,[§] Douglas E. Brenneman,[§] and Allen B. Reitz^{*,§}

Johnson & Johnson Pharmaceutical Research and Development, LLC, Spring House, Pennsylvania 19477, and Turnhoutseweg 30, 2340 Beerse, Belgium

Received May 8, 2007

Abstract: A new aspartic protease inhibitory chemotype bearing a 2-amino-3,4-dihydroquinazoline ring was identified by high-throughput screening for the inhibition of BACE-1. X-ray crystallography revealed that the exocyclic amino group participated in a hydrogen bonding array with the two catalytic aspartic acids of BACE-1 (Asp₃₂, Asp₂₂₈). BACE-1 inhibitory potency was increased (0.9 μ M to 11 nM K_i) by substitution into the unoccupied S₁' pocket.

Alzheimer's disease (AD) is a chronic degenerative disorder of the central nervous system, resulting in severe cognitive deficits along with psychiatric complications.¹ More than 20 million individuals suffer from this condition worldwide, and this number is expected to grow dramatically. The current standards of care are the cholinesterase inhibitors such as galantamine, which are indicated for mild to moderate AD, and the NMDA receptor antagonist memantine for the moderate to severe form of the disease. A compelling molecular target among the various strategies being pursued² is the aspartic

Table 1. Inhibition of BACE by 1–3a

cmpd	K_i (nM)
1	900
2	158
3	30
3a	11





Figure 1. Left: superposition of OM99-2 (peptidic inhibitor) and 1 in the active site of BACE-1. Right: ribbon diagram representation of 1 with the two catalytic aspartic acids highlighted bound into BACE-1 (maroon), showing the flap over the active site moved when compared with BACE-1 when OM99-2 is bound in (turquoise).

protease BACE-1.³ This membrane bound type 1 protein catalyzes the initial proteolytic cleavage of the amyloid precursor protein (APP) en route to the β -amyloid_{1–40(42)} peptides. The mouse homozygous knock-out shows significant reduction of β -amyloid levels,⁴ and this gene deletion in the Tg2576 animal model of β -amyloid overexpression results in the attenuation of the cognitive deficit.⁵ BACE-1 has also recently been shown to play a role in myelination in the peripheral and central nervous system during development and may have cognitive and synaptic functions independent of APP processing.⁶ Nevertheless, BACE-1 inhibitors have been shown to lower β -amyloid levels in the brains of mice⁷ and thus may treat the etiology of the disease and not just the symptoms.



A screen of our corporate compound collection resulted in the identification of a variety of hits in the 1–10 μ M range. We focused SAR development on 2-aminoquinazoline 1, which

* To whom correspondence should be addressed. Phone: 215-628-7119 (E.B.); 215-628-5615 (A.R.). Fax: 215-628-4985 (E.B.); 215-540-4776 (A.R.). E-mail: ebaxter@prdus.jnj.com (E.B.); areitz@prdus.jnj.com (A.R.).

[§] Johnson & Johnson Pharmaceutical Research and Development, Spring House, PA.

[†] Johnson & Johnson Pharmaceutical Research and Development, 2340 Beerse, Belgium.





Figure 2. Left: the 2-amino-3,4-dihydroquinazoline fragment of **1** and the Asp₃₂ and Asp₂₂₈ sidechains of BACE-1, with H-bond distances in Å between the Asp carboxylate oxygens and the hydrogens. Right: the X-ray structure of **3a** bound in to BACE-1, showing the α -cyclohexyl substituent filling the S₁' pocket.

inhibited BACE-1 with a 0.9 μ M K_i (Table 1). An X-ray structure of **1** bound into the BACE-1 protein was obtained as part of a collaboration with Astex Therapeutics. The BACE-1 protein used in the crystallography studies was prepared by recombinant insect cell expression.⁸ The structure of **1** in the active site of BACE-1 was compact, with the side chain bending back onto itself in a hairpin turn orientation (Figure 1). This allowed the *N*-cyclohexyl substituent to occupy the critical S₁ binding pocket. In addition, the flap region of the protein adopted an "open" structure in which the Tyr⁷¹C α carbon is displaced by 4 Å relative to the peptidic OM99-2⁹ bound into BACE-1. Similar flexibility has been observed for the flap region of other aspartyl proteases such as renin and HIV-1 protease.

An especially intriguing feature of this structure is the exquisite hydrogen bonding array that **1** forms with the two catalytic aspartates (Asp₃₂, Asp₂₂₈; Figure 2). Related 2-amino-pyridines, 2,3-diaminopyridines, 1-aminoisoquinolines, and 2-aminoquinolines have been identified by the use of a fragment-screening approach as BACE-1 inhibitors.¹⁰


To reinforce the hairpin conformation in the side chain, we prepared **2**, which bears a 1,3-disubstituted phenyl ring in the side chain. This modification also incorporated 6-phenoxy substitution instead of the 6-benzoyl group present in **1**. A 6-fold increase in potency for BACE-1 inhibition was observed for **2** (158 nM K_i) when compared with **1**. Further examination of



the X-ray structure of **1** bound into BACE-1 revealed that the vacant hydrophobic S₁' pocket could be filled by substitution with *S* stereochemistry on the α -carbon of the side chain. Compound **3** was prepared as a racemate initially and was found to be a potent BACE-1 inhibitor (30 nM K_i). The enantiomers of **3** were obtained independently, and the activity resided predominantly in the *S* enantiomer as predicted (**3a**, 11 nM K_i , Figure 2).


Chemistry. 2-Amino-3,4-dihydroquinazoline **1** was prepared starting with the conversion of ethyl cyanoacetate **4** into amide **5**, which afforded amine **6** after reduction (Scheme 1). Reductive amination of 3-benzoyl-6-nitrobenzaldehyde **7**¹¹ and **6** provided benzyl amine **8**. Reduction of the nitro group of **8** followed by

Scheme 1^a




^a Reagents and conditions: (a) MeNH-*c*-C₆H₁₁, EtOH, reflux; (b) H₂, Raney Ni, NH₃/MeOH; (c) NaBH(OAc)₃, CH₂Cl₂; (d) H₂, Pd/C, thiophene, MeOH; (e) BrCN, EtOH, reflux.

Scheme 2^a



^a Reagents and conditions: (a) MeNH-*c*-C₆H₁₁, HBTU, NEM, DMF; (b) TFA, CH₂Cl₂; (c) K₂CO₃, DMF, 120 °C, 1 h; (d) NaBH(OAc)₃, CH₂Cl₂; (e) SnCl₂ 2H₂O, EtOH, reflux; (f) BrCN, EtOH.

Scheme 3^a




^a Reagents and conditions: (a) MeNH-*c*-C₆H₁₁, *i*-BuOCOCl, TEA, CH₂Cl₂; (b) NH₄OAc, NaBH₃CN, MeOH; (c) NaBH₃CN, HOAc, **13**, MeOH; (d) H₂, 10% Pd/C, MeOH; (e) BrCN, EtOH, reflux.

treatment with cyanogen bromide led to 2-amino-3,4-dihydroquinazoline **1**.

The preparation of **2** was initiated by conversion of **9** to amide **10** and deprotection to provide **11** (Scheme 2). Nitrobenzaldehyde **12** was converted to phenoxy derivative **13**.¹² Reductive amination of **13** and **11** afforded benzylamine **14**, which was converted to aminoquinazoline **2**. The chemistry to prepare **3** (racemic) started with amidation of ketoacid **15**¹³ to give **16**, followed by reductive amination to afford primary amine **17** (Scheme 3). Conversion to target **3** was conducted by the use of the same sequence as for **1** and **2**. The single enantiomers of **3** were obtained via a synthetic strategy to prepare γ -substituted- γ -aminoacids devised by Smrcina and Majer,¹⁴ which resulted in no detectable epimerization on a reaction scale of up to 200 g.

^a Abbreviations: APP, amyloid precursor protein; BACE-1, β -site APP cleaving enzyme or Asp-2 or memapsin-2; NEM, *N*-ethylmorpholine.

Scheme 4^a

^a Reagents and conditions: (a) Meldrum's acid, DMAP, EDCI, CH₂Cl₂; (b) NaBH₄, HOAc; (c) toluene, reflux; (d) NaOH, acetone; (e) MeNH-*c*-C₆H₁₁, HOBT, EDCI, TEA, DMF; (f) TFA, CH₂Cl₂; (g) **13**, NaBH(OAc)₃, CH₂Cl₂; (h) H₂, Pd/C; (i) BrCN, EtOH.

For the preparation of **3a**, (*R*)-cyclohexylglycine **18** (>97% e.e.) was treated with Meldrum's acid under basic conditions to afford **19**, and the ketone functionality of **19** was subsequently reduced with NaBH₄ to provide **20**. Thermolysis of **20** provided lactam **21**, which was saponified to **22**, suitable for further homologation to **3a** (Scheme 4).

Pharmacology. Compound **3a** was further examined in a variety of assays to probe selectivity and suitability for further consideration as a lead. It displayed modest selectivity for BACE-1 over the aspartic proteases renin (2.7 μM IC₅₀) and cathepsin D (0.11 μM IC₅₀) as well as the hERG channel (0.14 μM IC₅₀). The Caco-2 apparent permeability coefficients *P*_{app}A to B and *P*_{app}B to A values for compound **3a** were determined to be 0.28 × 10⁻⁶ cm/s and 3.38 × 10⁻⁶ cm/s, respectively. The efflux ratio of 12.2 suggests that this compound is a PgP substrate, although testing against PgP itself would need to be done to confirm this hypothesis. Compound **3a** exhibited excellent potency in a cellular assay, which measures the inhibition of Aβ₁₋₄₀ secretion in CHO cells transfected with the Swedish familial AD mutant APP (K670N/M671L). Additionally, **3a** lowered β-amyloid₁₋₄₀ in plasma by 40–70% in rats after p.o. administration (30 mg/kg), 3 h post-dosing.

Discussion. A variety of compounds have been reported that inhibit BACE-1.^{3,15} Many of these are statine-like pseudo-transition state inhibitors, and others have more dramatically altered chemotypes such as the 2-aminoheterocycles recently reported¹⁰ that are similar to the 2-amino-3,4-dihydroquinazolines disclosed here. Considerable computational work has also been conducted examining the BACE-1 active site in an effort to design more potent inhibitors or add suitable substitution to address ADME or other liabilities.¹⁶ There were several very distinctive features of our 2-amino-3,4-dihydroquinazoline chemotype. The 2-amino-3,4-dihydroquinazolines adopted a compact structure bearing a hairpin turn of the side chain and anchored at the active site by virtue of an extensive hydrogen bonding array of the two catalytic aspartic acids with the exocyclic amino group. As a consequence, the protein flap over the active site was pushed away by a distance of about 4 Å when compared to the structure of OM99-2 (peptidic inhibitor) bound in to BACE-1. Because only the S₁ pocket of the protein was filled with initial 2-amino-3,4-dihydroquinazoline ligand **1** we were able to explicitly design several modifications that served to improve potency. The hairpin turn was accommodated by the incorporation of a structural constraint (e.g., **2**), and the S₁' pocket was filled by (S)-α-(*c*-C₆H₁₂) substitution.

Conclusion. The 2-amino-3,4-dihydroquinazoline substructure is a new fragment for aspartic protease inhibition such as

against BACE-1, which is an important target in the search for new Alzheimer's disease treatments. The exocyclic amino group was shown to be involved in a hydrogen bonding array with the two aspartic acids at the BACE-1 active site. X-ray crystallography analysis of **1** (0.9 μM K_i) bound in to BACE-1 was critical in allowing us to design additional analogs with improved potency. For example, because the side chain of **1** adopted an energetically unfavored hairpin conformation, 1,3-phenyl disubstituted derivative **2** was prepared, which displayed improved potency (158 nM K_i) relative to **1**. Careful analysis of the structure of **1** bound into BACE-1 revealed that the S₁' pocket could be accessed by (S)-α-substitution off of the side chain, so we designed and prepared by (S)-α-(*c*-C₆H₁₁) derivative **3a**, which had excellent potency for BACE-1 inhibition (11 nM K_i). These structural modifications (**2** and **3a**) would not have been given high priority for synthesis without the insight provided by the crystal structure of **1** bound in to BACE-1. We were able to transition from a micromolar hit to a nanomolar lead suitable for extensive SAR development by the use of ligand–BACE-1 crystallographic analysis.

Acknowledgment. We thank Carlos Plata-Salaman, Kevin Felsenstein, Chris Teleha, Bill Jones, Laura Reany, Scott Ballentine, Sam Nortey, Jennifer Piesvaux, Tina Ross, Andrew Darrow, Sergey Ilyin, Christopher Creighton, Umar Maharroof, and Eric Strobel for helpful assistance and support. We also acknowledge Astex Therapeutics for obtaining the X-ray crystal structures.

Supporting Information Available: Chemistry experimentals for the preparation of **1**, **2**, **3** and **3a**. X-ray structure coordinates for **1** and **3a** bound in to BACE-1 have been deposited into the Protein Databank (www.pdb.org, file codes 2Q15 and 2Q11, respectively). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Goedert, M.; Spillantini, M. G. A Century of Alzheimer's Disease. *Science* **2006**, *314*, 777–781. (b) Roberson, E. D.; Mucke, L. 100 Years and Counting: Prospects for Defeating Alzheimer's Disease. *Science* **2006**, *314*, 781–784.
- (2) (a) Robichaud, A. J. Approaches to Palliative Therapies for Alzheimer's Disease. *Curr. Top. Med. Chem.* **2006**, *6*, 553–568. (b) Churcher, I. Tau Therapeutic Strategies for the Treatment of Alzheimer's Disease. *Curr. Top. Med. Chem.* **2006**, *6*, 579–595. (c) Catalano, S.; Dodson, E. C.; Henze, D. A.; Joyce, J. G.; Krafft, G. A.; Kinney, G. G. The Role of Amyloid-Beta Derived Diffusible Ligands (ADDLs) in Alzheimer's Disease. *Curr. Top. Med. Chem.* **2006**, *6*, 597–608.
- (3) (a) John, V. Human β-Secretase (BACE) and BACE Inhibitors: Progress Report. *Curr. Top. Med. Chem.* **2006**, *6*, 569–578. (b) Baxter, E. W.; Reitz, A. B. BACE Inhibitors for the Treatment of Alzheimer's Disease. *Annu. Rep. Med. Chem.* **2005**, *40*, 35–48. (c) John, V.; Beck, J. P.; Bienkowski, M. J.; Sinha, S.; Heinrichson, R. L. Human β-Secretase (BACE) and BACE Inhibitors. *J. Med. Chem.* **2003**, *46*, 4625–4630.
- (4) (a) Luo, Y.; Bolon, B.; Damore, M. A.; Fitzpatrick, D.; Liu, H.; Zhang, J.; Yan, Q.; Vassar, R.; Citron, M. BACE1 (β-Secretase) Knockout Mice do not Acquire Compensatory Gene Expression Changes or Develop Neural Lesions over Time. *Neurobiol. Dis.* **2003**, *14*, 81–88 and references cited therein. (b) Kouhei, N.; Gaku, S.; Yuko, H.; Gray, A. J.; Maeda, M.; Hirata-Fukae, C.; Becker, A. G.; Hosono, M.; Sakaguchi, I.; Minami, S. S.; Nakajima, Y.; Li, H.-F.; Takeyama, C.; Kihara, T.; Ota, A.; Wong, P. C.; Aisen, P. S.; Kato, A.; Kinoshita, N.; Matsuoka, Y. BACE1 Inhibition Reduces Endogenous Abeta and Alters APP Processing in Wild-Type Mice. *J. Neurochem.* **2006**, *99*, 1555–1563.
- (5) Ohno, M.; Sametsky, E. A.; Younkin, L. H.; Oakley, H.; Younkin, S. G.; Citron, M.; Vassar, R.; Disterhoft, J. F. BACE1 Deficiency Rescues Memory Deficits and Cholinergic Dysfunction in a Mouse Model of Alzheimer's Disease. *Neuron* **2004**, *41*, 27–33.
- (6) (a) Hu, X.; Hicks, C. W.; He, W.; Wong, P.; Macklin, W. B.; Trapp, B. D.; Yan, R. BACE1 Modulates Myelination in the Central and Peripheral Nervous System. *Nat. Neurosci.* **2006**, *9*, 1520–1525. (b) Laird, F. M.; Cai, H.; Savonenko, A. V.; Farah, M. H.; He, K.

- Melnikova, T.; Wen, H.; Chiang, H.-Ch.; Xu, G.; Koliatsos, V. E.; Borchelt, D. R.; Price, D. L.; Lee, H.-K.; Wong, P. C. BACE1, a Major Determinant of Selective Vulnerability of the Brain to Amyloid- β Amyloidogenesis, is Essential for Cognitive, Emotional, and Synaptic Functions. *J. Neurosci.* **2005**, *25*, 11693–11709.
- (7) (a) Asai, M.; Hattori, C.; Iwata, N.; Saido, T. C.; Sasagawa, N.; Szabó, B.; Hashimoto, Y.; Maruyama, K.; Tanuma, S.; Kiso, Y.; Ishiura, S. The Novel β -Secretase Inhibitor KMI-429 Reduces Amyloid β Peptide Reduction in Amyloid Precursor Protein Transgenic and Wild Type Mice. *J. Neurochem.* **2006**, *96*, 533–540. (b) Stachel, S. J.; Coburn, C. A.; Sankaranarayanan, S.; Price, E. A.; Pietrak, B. L.; Huang, Q.; Lineberger, J.; Espeseth, A. S.; Jin, L.; Ellis, J.; Holloway, M. K.; Munshi, S.; Allison, T.; Hazuda, D.; Simon, A. J.; Graham, S. L.; Vacca, J. P. Macrocyclic Inhibitors of β -Secretase: Functional Activity in an Animal Model. *J. Med. Chem.* **2006**, *49*, 6147–6150. (c) Hussain, I.; Hawkins, J.; Harrison, D.; Hille, C.; Wayne, G.; Cutler, L.; Buck, T.; Walter, D.; Demont, E.; Howes, C.; Naylor, A.; Jeffrey, P.; Gonzalez, M. I.; Dingwall, C.; Michel, A.; Redshaw, S.; Davis, J. B. Oral Administration of a Potent and Selective Non-Peptidic BACE-1 Inhibitor Decreases β -Cleavage of Amyloid Precursor Protein and Amyloid- β Production In Vivo. *J. Neurochem.* **2007**, *100*, 802–809. (d) Chang, W.-P.; Koelsch, G.; Wong, S.; Downs, D.; Da, H.; Weerasena, V.; Gordon, B.; Devasamudram, T.; Bilcer, G.; Ghosh, A. K.; Tang, J. In Vivo Inhibition of A β Production by Memapsin-2 (β -Secretase) Inhibitors. *J. Neurochem.* **2004**, *89*, 1409–1416.
- (8) (a) Bruinzeel, W.; Yon, J.; Giovannelli, S.; Masure, S. Recombinant Insect Cell Expression and Purification of Human β -Secretase (BACE-1) for X-ray Crystallography. *Protein Expression Purif.* **2002**, *26*, 139–148. (b) Patel, S.; Vuillard, L. P.; Cleasby, A.; Murray, C. W.; Yon, J. Apo and Inhibitor Complex Structures of BACE (β -Secretase). *J. Mol. Biol.* **2004**, *343*, 407–416.
- (9) Ghosh, A. K.; Shin, D.; Downs, D.; Koelsch, G.; Lin, X.; Ermolieff, J.; Tang, J. *J. Am. Chem. Soc.* **2000**, *122*, 3522–3523.
- (10) (a) Congreve, M.; Aharony, D.; Albert, J.; Callaghan, O.; Campbell, J.; Carr, R. A. E.; Chessari, G.; Cowan, S.; Edwards, P. D.; Frederickson, M.; McMenamin, R.; Murray, C. W.; Patel, S.; Wallis, N. Application of Fragment Screening by X-ray Crystallography to the Discovery of Aminopyridines as Inhibitors of β -Secretase. *J. Med. Chem.* **2007**, *50*, 1124–1132. (b) Murray, C. W.; Callaghan, O.; Chessari, G.; Cleasby, A.; Congreve, M.; Frederickson, M.; Hartshorn, M. J.; McManamin, R.; Patel, S.; Wallis, N. Application of Fragment Screening by X-ray Crystallography to β -Secretase. *J. Med. Chem.* **2007**, *50*, 1116–1123.
- (11) Freyne, E. J. E.; Venet, M. G.; Raeymaekers, A. H. M.; Sanz, G. C. Preparation of (1*H*-Azol-1-ylmethyl)quinolines, -quinazolines, and -quinoxalines as Drugs. European Patent 371,564, June 6, 1990.
- (12) For an alternative synthesis of **13**: Katritzky, A. R.; Wang, Z.; Hall, C. D.; Akhmedov, N. G. The Preparation of 5-Substituted and 4,5-Disubstituted 2,1-Benzisoxazoles. *ARKIVOC* **2003**, *2*, 49–58.
- (13) Betancourt de Perez, R. M.; Fuentes, L. M.; Larson, G. L.; Barnes, C. L.; Heeg, M. J. The Chemistry of α -Silyl Carbonyl Compounds. 12. A Synthesis of 4-Oxo Carboxylic Acids, 4-Oxo Aldehydes, and 1,4-Diketones from γ -Lactones. *J. Org. Chem.* **1986**, *51*, 2039–2043.
- (14) Smrcina, M.; Majer, P.; Majrova, E.; Guerassina, T. A.; Eissenstat, M. A. Facile Stereoselective Synthesis of γ -Substituted γ -Amino Acids from the Corresponding α -Amino Acids. *Tetrahedron* **1997**, *53*, 12867–12874.
- (15) (a) Maillard, M. C.; Hom, R. K.; Benzon, T. E.; Moon, J. B.; Mamo, S.; Bienkowski, M.; Tomasselli, A. G.; Woods, D. D.; Prince, D. B.; Paddock, D. J.; Emmons, T. L.; Tucker, J. A.; Dappen, M. S.; Brogley, L.; Thorsett, E. D.; Jewett, N.; Sinha, S.; John, V. Design, Synthesis, and Crystal Structure of Hydroxyethyl Secondary Amine-Based Peptidomimetic Inhibitors of Human β -Secretase. *J. Med. Chem.* **2007**, *50*, 776–771. (b) Rajapakse, H. A.; Nantermet, P. G.; Selnick, H. G.; Munshi, S.; McGaughey, G. B.; Lindsley, S. R.; Young, M. B.; Lai, M.-T.; Espeseth, A. S.; Shi, X.-P.; Colussi, D.; Pietrak, B.; Crouthamel, M.-Ch.; Tugusheva, K.; Huang, Q.; Xu, M.; Simon, A. J.; Kuo, L.; Hazuda, D. J.; Graham, S.; Vacca, J. P. Discovery of Oxadiazoyl Tertiary Carbinamine Inhibitors of β -Secretase (BACE-1). *J. Med. Chem.* **2006**, *49*, 7270–7273. (c) Patey, S. J.; Edwards, E. A.; Yates, E. A.; Turnbull, J. E. Heparin Derivatives as Inhibitors of BACE-1, the Alzheimer's β -Secretase, with Reduced Activity against Factor Xa and Other Proteases. *J. Med. Chem.* **2006**, *49*, 6129–6132. (d) Hanessian, S.; Yang, G.; Rondeau, J.-M.; Newmann, U.; Betschart, C.; Tintelnnot-Blomley, M.; Garino, C.; Tomita, T.; Pietrancosta, N.; Laras, Y.; Rosas, R.; Herbet, G.; Maigret, B.; Quelever, G.; Iwatsubo, T.; Kraus, J.-L. *J. Med. Chem.* **2006**, *49*, 4275–4285.
- (16) For example, see: (a) Tounge, B. A.; Rajamani, R.; Baxter, E. W.; Reitz, A. B.; Reynolds, C. H. Linear Interaction Energy Models for β -Secretase (BACE) Inhibitors: Role of van der Waals, Electrostatic and Continuum-Solvation Terms. *J. Mol. Graph. Model.* **2006**, *24*, 475–484. (b) Moitessier, N.; Therrien, E.; Hanessian, S. A Method for Induced-Fit Docking, Scoring, and Ranking of Flexible Ligands. Application to Peptidic and Pseudo-peptidic β -Secretase (BACE 1) Inhibitors. *J. Med. Chem.* **2006**, *49*, 5885–5894.

JM0705408