

Determination of crystal structures by X-ray diffraction

Outline

- 1 Bragg and Von Laue formulation of X-ray diffraction by a crystal
- 2 Experimental geometries suggested by the Laue condition
- 3 The geometrical structure factor
- 4 The atomic form factor


- 1 Bragg and Von Laue formulation of X-ray diffraction by a crystal
- 2 Experimental geometries suggested by the Laue condition
- 3 The geometrical structure factor
- 4 The atomic form factor

Diffraction by a crystal

The electromagnetic probe

X-ray diffraction

- Interatomic distances are of the order of Å
 - 10^{-8}cm
 - $E = \hbar\omega = \frac{hc}{\lambda} \sim 12.3 \times 10^3 \text{ eV}$
- Wavelength and energies characteristic of X-rays
- Sharp peaks of scattered radiation
 - due to long range order
 - not found for amorphous solids or liquids




X-ray diffraction pattern from a crystal

Diffraction by a crystal

The electromagnetic probe

X-ray diffraction

- We consider a **rigid** lattice of ions
- Effect of vibrations:
 - decrease the intensity of the scattered peaks
 - contribute to the diffuse background




X-ray diffraction pattern from a crystal

Diffraction by a crystal

X-ray diffraction

Equivalent Formulations

- **Bragg** formulation
 - used by crystallographers
- **Von Laue** formulation
 - exploits the reciprocal lattice
 - closer to the solid-state approach



X-ray diffraction pattern from a crystal

X-ray diffraction by a crystal

Bragg formulation

Bragg's interpretation of X-ray diffraction


- Crystal composed of parallel planes (**lattice planes**)
 - separated by a distance d
- **Conditions** for the appearance of sharp **diffraction peaks**
 - X-rays are **specularly reflected** by the crystal planes
 - **constructive interference** of reflected X-rays
- **Bragg's condition:** $n\lambda = 2d \sin \theta$
 - n : **order of reflection**
 - θ : **angle of incidence** on the crystal's plane

X-ray diffraction by a crystal


Bragg interpretation of X-ray diffraction

Simple derivation of Bragg condition

- Condition for **constructive interference**:
 - path difference ($2d \sin \theta$) equals an integral number of wavelengths
 - total angle of deflection of the incident rays: 2θ



reflection from a family of lattice planes





Bragg angle θ

X-ray diffraction by a crystal

Bragg's interpretation of X-ray diffraction

Further observations

- A large number of reflections arise as a result of
 - **different wavelengths** of incident X-rays
 - **different reflection orders** n for a given set of planes
 - **different set** of lattice planes (infinitely many)




Two possible resolutions of the same crystal lattice into planes

X-ray diffraction by a crystal

Von Laue formulation

Assumptions

- Crystal composed of **scatterers** at the sites \mathbf{R} of a Bravais lattice
 - atoms, ions
- Peaks are observed for directions of constructive interference between all scattered rays
- **no** resolution of the lattice into crystal planes
- **no** need to assume specular reflection




two scattering centers separated by a displacement vector \mathbf{d}

X-ray diffraction by a crystal

Von Laue formulation

Derivation of the condition of constructive interference

- Wave vector of incident radiation: $\mathbf{k} = \frac{2\pi}{\lambda} \hat{\mathbf{n}}$
- Wave vector of scattered radiation: $\mathbf{k}' = \frac{2\pi}{\lambda} \hat{\mathbf{n}}'$
 - elastic scattering
- Path difference: $\mathbf{d} \cdot (\hat{\mathbf{n}} - \hat{\mathbf{n}}')$
 - $\mathbf{d} \cdot (\mathbf{k} - \mathbf{k}') = 2\pi m$ (m integer)




two scattering centers separated by a displacement vector \mathbf{d}

X-ray diffraction by a crystal

Von Laue formulation

Derivation of the condition of constructive interference

- For **all** scatterers in the lattice: $\mathbf{R} \cdot (\mathbf{k} - \mathbf{k}') = 2\pi m, \forall \mathbf{R}$
 - **all scattered rays** interfere constructively
- Alternatively: $e^{i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{R}} = 1$
 - $\mathbf{k} - \mathbf{k}'$ is a **reciprocal lattice vector** \mathbf{K}




two scattering centers separated by a displacement vector \mathbf{d}

X-ray diffraction by a crystal

Von Laue formulation

Another geometrical interpretation

- $\mathbf{k} - \mathbf{k}'$ is a reciprocal lattice vector \mathbf{K}
- $k = |\mathbf{k} - \mathbf{K}|$ and squaring
- $\mathbf{k} \cdot \hat{\mathbf{K}} = \frac{1}{2}K$
 - **component** of \mathbf{k} along \mathbf{K}




k-space plane (Bragg plane)

X-ray diffraction by a crystal

Equivalence of Bragg and Von Laue formulations

Proof

- Von Laue condition: $\mathbf{k}' - \mathbf{k} = \mathbf{K}$ ($k' = k$)
- \mathbf{K} is \perp to a family of direct lattice planes




\mathbf{K} bisects the angle between \mathbf{k} and \mathbf{k}'

X-ray diffraction by a crystal

Equivalence of Bragg and Von Laue formulations

Proof

- if d distance between planes, $|\mathbf{K}| = 2k \sin \theta = n|\mathbf{K}_0| = n \frac{2\pi}{d}$
- $k \sin \theta = \frac{n\pi}{d}$ (Bragg condition)
- Reflection from the lattice planes $\perp \mathbf{K}$
- The order of reflection is $n = \frac{|\mathbf{K}|}{|\mathbf{K}_0|}$



\mathbf{K} bisects the angle between \mathbf{k} and \mathbf{k}'

- 1 Bragg and Von Laue formulation of X-ray diffraction by a crystal
- 2 Experimental geometries suggested by the Laue condition**
- 3 The geometrical structure factor
- 4 The atomic form factor

Experimental geometries suggested by the Laue condition

The Laue condition

Devising experimental setups


- **Laue condition:** the **tip** of \mathbf{k} must lie on a Bragg plane
 - **k -space plane**
- Difficult to realize for fixed **orientation** and λ
- **How do we achieve enough sampling of the reciprocal space?**
 - **vary** the wavelength of X-rays
 - **vary** the direction of incidence (i.e. **relative** orientation of the crystal)
- **Ewald** construction

Experimental geometries suggested by the Laue condition

A geometrical construction

The Ewald sphere

- Draw a sphere of radius k centered on the tip of \mathbf{k} ($k = \frac{2\pi}{\lambda}$)
 - passes through the origin
- Diffraction peaks for lattice points on the surface of the sphere
 - \mathbf{k}' satisfies the Laue condition




the Ewald construction

Experimental geometries suggested by the Laue condition

The Laue method


- Use polychromatic X-rays (from λ_1 to λ_0)
 - fixed orientation of the crystal and incident direction \hat{n}
 - $\mathbf{k}_1 = \frac{2\pi}{\lambda_1} \hat{n}$, $\mathbf{k}_0 = \frac{2\pi}{\lambda_0} \hat{n}$
- Diffracted rays in correspondence to multiple reciprocal lattice points
 - region between the two spheres



Experimental geometries suggested by the Laue condition

The rotating-crystal method

- Use **monochromatic** X-rays of fixed incident direction
- Vary the **orientation** of the crystal
 - rotation around a fixed axis
 - the reciprocal lattice **rotates around the same axis by the same amount**





the Ewald construction for the rotating-crystal method

Experimental geometries suggested by the Laue condition

The Debye-Scherrer Method

Powder Method

- Rotating-crystal method with rotation axis **over all possible directions**
 - **finely dispersed powder** (randomly oriented crystals)
 - Each **K** generates a **sphere** of radius K
- All **K** such that $K < 2k$ generates a **cone** of diffracted radiation
 - $K = 2k \sin \frac{1}{2}\phi$



the Ewald construction for the Debye-Scherrer Method


- 1 Bragg and Von Laue formulation of X-ray diffraction by a crystal
- 2 Experimental geometries suggested by the Laue condition
- 3 The geometrical structure factor**
- 4 The atomic form factor

Diffraction by a monoatomic lattice with a basis

The geometrical structure factor

Several identical scatterers in the primitive cell

- n scatterers at positions $\{\mathbf{d}_i\}_{i=1,\dots,n}$
 - n -atom **basis** (e.g. diamond structure: $n=2$)
- For a **Bragg peak** with $\mathbf{K} = \mathbf{k}' - \mathbf{k}$
 - **constructive/destructive** interference btw scattered rays
 - **Phase difference**: $\mathbf{K} \cdot (\mathbf{d}_i - \mathbf{d}_j)$



path difference btw rays scattered by centers at a distance d

Diffraction by a monoatomic lattice with a basis

The geometrical structure factor

Several identical scatterers in the primitive cell

- The **amplitude** of the rays will differ by a **factor** $e^{i\mathbf{K}\cdot(\mathbf{d}_i-\mathbf{d}_j)}$
- For the n scatterers the amplitudes are in the **ratio**:

$$e^{i\mathbf{K}\cdot\mathbf{d}_1} : e^{i\mathbf{K}\cdot\mathbf{d}_2} : \dots : e^{i\mathbf{K}\cdot\mathbf{d}_n}$$

- The **total amplitude** of X-ray scattered by the cell contains the factor

$$S_{\mathbf{K}} = \sum_{j=1}^n e^{i\mathbf{K}\cdot\mathbf{d}_j}$$

- $S_{\mathbf{K}}$: **geometrical structure factor**
- $I_{\mathbf{K}} \propto |S_{\mathbf{K}}|^2$

Diffraction by a monoatomic lattice with a basis

The geometrical structure factor

Absolute intensity in a Bragg peak

- The **intensity** depends on \mathbf{K} through $S_{\mathbf{K}}$
- **Not** the only source of \mathbf{K} dependence
 - characteristic **angular dependence** of the scattering process
 - **internal structure** of the scatterer
- $S_{\mathbf{K}}$ alone **cannot** be used to predict the absolute intensity
- When $S_{\mathbf{K}} = 0 \implies I_{\mathbf{K}} = 0$
 - complete **destructive** interference

Vanishing structure factor

Examples

bcc viewed as a sc lattice with a basis


- The reciprocal lattice is **fcc**
- **bcc** can be regarded as a **sc lattice with a basis**
 - **primitive vectors:** $a\hat{x}$, $a\hat{y}$, $a\hat{z}$
 - **basis:** $\mathbf{d}_1 = 0$, $\mathbf{d}_2 = (\frac{a}{2})(\hat{x} + \hat{y} + \hat{z})$
- \mathbf{K} must be a vector of the reciprocal lattice
 - $\mathbf{K} = \frac{2\pi}{a}(n_1\hat{x} + n_2\hat{y} + n_3\hat{z})$
- $S_{\mathbf{K}} = 1 + e^{i\pi(n_1+n_2+n_3)} = 1 + (-1)^{n_1+n_2+n_3}$
 - $S_{\mathbf{K}} = 2$ when $n_1 + n_2 + n_3$ is **even**
 - $S_{\mathbf{K}} = 0$ when $n_1 + n_2 + n_3$ is **odd**

Vanishing structure factor

Examples

bcc viewed as a sc lattice with a basis

- \mathbf{K} vectors for which $S_{\mathbf{K}} = 0$ will have **no Bragg reflection**
 - **odd** number of nearest-neighbour bonds
 - from the origin
- \mathbf{K} vectors for which $S_{\mathbf{K}} \neq 0$ define a **reciprocal fcc lattice**
 - side of $\frac{4\pi}{a}$



\mathbf{K} points for which $S_{\mathbf{K}} = 2$ (black circles) and $S_{\mathbf{K}} = 0$ (white circles)

Vanishing structure factor

Examples

Monoatomic diamond lattice (C, Si, Ge, grey tin)


- Not a Bravais lattice
- Viewed as a fcc lattice with a two-atom basis
 - $\mathbf{a}_1 = \frac{a}{2}(\hat{y} + \hat{z})$ etc
 - basis: $\mathbf{d}_1 = 0$, $\mathbf{d}_2 = \frac{a}{4}(\hat{x} + \hat{y} + \hat{z})$
- \mathbf{K} must be a vector of the bcc reciprocal lattice $\mathbf{K} = \sum_i n_i \mathbf{b}_i$
 - cubic cell of side of $\frac{4\pi}{a}$
 - $\mathbf{b}_1 = \frac{2\pi}{a}(\hat{y} + \hat{z} - \hat{x})$ etc
- $S_{\mathbf{K}} = 1 + e^{i\frac{\pi}{2}(n_1+n_2+n_3)}$
 - $S_{\mathbf{K}} = 2$ when $n_1 + n_2 + n_3$ is twice an even number
 - $S_{\mathbf{K}} = 0$ when $n_1 + n_2 + n_3$ is twice an odd number
 - $S_{\mathbf{K}} = 1 \pm i$ when $n_1 + n_2 + n_3$ is odd

Vanishing structure factor

Examples

Monoatomic diamond lattice (C, Si, Ge, grey tin)

- $\mathbf{K} = \sum_i n_i \mathbf{b}_i = \frac{4\pi}{a} (\nu_1 \hat{\mathbf{x}} + \nu_2 \hat{\mathbf{y}} + \nu_3 \hat{\mathbf{z}})$
 - $\nu_j = \frac{1}{2}(n_1 + n_2 + n_3) - n_j$
 - $\sum_j \nu_j = \frac{1}{2}(n_1 + n_2 + n_3)$
- The **bcc** is viewed as composed of two **sc lattices**
- The **first** contains the origin ($\mathbf{K} = 0$)
 - ν_i are integers ($n_1 + n_2 + n_3$ twice an even/odd)
 - $S_{\mathbf{K}} = 0, 2$ ($S_{\mathbf{K}} = 0$ when $\sum_j \nu_j$ is odd, as before)




\mathbf{K} points for which $S_{\mathbf{K}} = 2$, $S_{\mathbf{K}} = 1 \pm i$, and $S_{\mathbf{K}} = 0$ (white circles)

Vanishing structure factor

Examples

Monoatomic diamond lattice (C, Si, Ge, grey tin)

- $\mathbf{K} = \sum_i n_i \mathbf{b}_i = \frac{4\pi}{a} (\nu_1 \hat{\mathbf{x}} + \nu_2 \hat{\mathbf{y}} + \nu_3 \hat{\mathbf{z}})$
 - $\nu_j = \frac{1}{2}(n_1 + n_2 + n_3) - n_j$
 - $\sum_j \nu_j = \frac{1}{2}(n_1 + n_2 + n_3)$
- The **bcc** is viewed as composed of two **sc lattices**
- The **second** contains $\mathbf{K} = \frac{4\pi}{a} \frac{1}{2} (\hat{\mathbf{x}} + \hat{\mathbf{y}} + \hat{\mathbf{z}})$
 - all ν_i must be **integer** + $\frac{1}{2}$ ($n_1 + n_2 + n_3$ **odd**)
 - $S_{\mathbf{K}} = 1 \pm i$



\mathbf{K} points for which $S_{\mathbf{K}} = 2$, $S_{\mathbf{K}} = 1 \pm i$, and $S_{\mathbf{K}} = 0$ (white circles)

- 1 Bragg and Von Laue formulation of X-ray diffraction by a crystal
- 2 Experimental geometries suggested by the Laue condition
- 3 The geometrical structure factor
- 4 The atomic form factor**

Diffraction by a polyatomic crystal

The atomic form factor

Scattering by different centers in the basis

- If the scatterers are not identical

$$S_{\mathbf{K}} = \sum_{j=1}^n f_j(\mathbf{K}) e^{i\mathbf{K} \cdot \mathbf{d}_j}$$

- $f_j(\mathbf{K})$: atomic form factor
- depends on its internal structure
- identical centers have identical $f_j(\mathbf{K})$
- consistent with previous treatment

Diffraction by a polyatomic crystal

The atomic form factor

Scattering by different centers in the basis

- In simple treatments

$$f_j(\mathbf{K}) = -\frac{1}{e} \int d\mathbf{r} e^{i\mathbf{K}\cdot\mathbf{r}} \rho_j(\mathbf{r})$$

- **Fourier transform of $\rho_j(\mathbf{r})$**
- **$\rho_j(\mathbf{r})$** : electronic charge density of ion of **type j** at **$\mathbf{r} = 0$**