

**Università di Trieste**  
**Corso di Laurea in Geologia**

**Anno accademico 2016 - 2017**

**Geologia Marina**

Parte I

**Modulo 1.2** Oceani. Morfologia, struttura ed evoluzione

Docente  
**Martina Buseti**

# The oceans

1. The present oceans of the Earth
2. Morphology of the ocean sea floor
3. Structure of the ocean
4. Evolution of the ocean

# 1. Etymology

The term Ocean derives from Ὠκεανὸς (ὠκεανός), greek river-god that was believed to surround the world, the external sea (not the Mediterranean).

But the rooth of word are from sanscrit ACAYANA, in the sense of “containing the waters.



Okeanos Fontana di Trevi

Okeanos is one of the Titans, son of Uran (sky) and Gea (earth), husband of **Teti**, and father of all the fluvial divinities.



# The Oceans are important because they:

- Cover 71-72% of the Earth surface
- Contain the 97% of the water of the Earth


but less than 5-10% has been explored





# The present Oceans of the world






In the 1953, the International Hydrographic Bureau defined three oceans:

- the Pacific Ocean
- the Atlantic Ocean
- the Indian Ocean

# Arctic Ocean






# Southern Ocean

In 2000, the International Hydrographic Organization (IHO), published a draft definition of a new ocean:


## the Southern Ocean

surrounding Antarctica and extending to 60° S.  
Up to now, the Southern Ocean is not formally defined by the IHO.






# IHS's delineation of the Southern Ocean



# The Antarctic Convergence: a possible limit of the Southern Ocean



## **Antarctic Convergence**

a curve encircling Antarctica, varying in latitude seasonally, where cold, northward-flowing Antarctic waters meets the relatively warmer water of the subantarctic

Ocean	Area	Average Depth (m)	Deepest depth (m)
Pacific Ocean	165,250,000 km <sup>2</sup>	4,028 m	Mariana Trench 11,033 m
Atlantic Ocean	106,400,000 km <sup>2</sup>	3,926 m	Puerto Rico Trench 8,604 m
Indian Ocean	73,560,000 km <sup>2</sup>	3,963 m	Java Trench, 7,725 m
Southern Ocean	20,330,000 km <sup>2</sup>	4,000 to 5,000 m	the southern end of the South Sandwich 7,236 m
Arctic Ocean	13,990,000 km <sup>2</sup>	1,205 m	Eurasia Basin, 5,540 m



# SEA FLOOR MORPHOLOGY





Harry Hess (1906 – 1969)

Professor of geology at Princeton University

During the Second World War, Hess was the captain of a ship equipped with a SONAR (SOund NAvigation and Ranging, invented in 1917 by Paul Langevin).

Hess discovered the Mid-oceanic ridges and the guyots, and in the '62 he published the Sea floor Spreading theory, fundamental for the Plate Tectonic theory.



The SONAR principle


# SEA FLOOR MORPHOLOGY 1977



Marie Tharp and Bruce Heezen, oceanographers of the Columbia University's Lamont Geological Observatory. They discovered the 60.000 km of underwater ridges. The map was painted by Heinrich C. Berann.




# Mid-Ocean Ridges



- The largest feature of the ocean floor.
- Linear belt of submarine mountains and active volcanoes about 60.000 km long.
- Plate boundaries: new magma forces its way up between two plates and pushes them apart.




# Global sea floor topography from gravity data derived from satellite altimetry and shipboard depth soundings



SMITH AND SANDWELL, 1996




## Global gravity map of the oceans - from GEOSAT and ERS-1




SMITH AND SANDWELL, 1996



## Global gravity map of the oceans from GEOSAT and ERS-1



## Global sea floor topography from gravity data and shipboard depth soundings




SMITH AND SANDWELL, 1996

Dixon *et al.*, *JGR*, (1983) have summarized the basic theory for estimating sea floor topography from gravity anomalies.

Models of the isostatic compensation of sea floor topography furnish a spectral transfer function that predicts the gravity anomaly expected from sea floor topography.


This transfer function depends on: mean depth, crustal density and thickness, and elastic lithosphere thickness.







## Marine gravity model of the North Atlantic




Red dots show locations of earthquakes with magnitude  $> 5.5$  and they highlight the present-day location of the seafloor spreading ridges and transform faults.


This gravity information shows the details of the plate tectonic history of the rifting of these continents including the subtle signatures of fracture zones that are currently buried by sediment.



# SEA FLOOR MORPHOLOGY




# SEA FLOOR MORPHOLOGY






## Geomorphic features map of the world's oceans.



Harris, Macmillan-Lawler, Rupp, Baker, 2014. **Geomorphology of the oceans.**  
Marine Geology, 352, 2014, 4–24. <http://dx.doi.org/10.1016/j.margeo.2014.01.011>


# The Ocean formation



continental rift




# SEA FLOOR MORPHOLOGY





Mid-oceanic ridge system is 60,000 km long, 2000 km wide, 3000 m<sup>2</sup> high













Paired lines on either side of the axis show the amount of crust generated in the last 10 million years at the current opening rates.

Dotted lines show the volcanic arcs, which are lines of volcanoes and volcanic islands formed from magma rising from the subducted plate. The distortion caused by plotting these lines on a Mercator projection is indicated by the hourglass-shaped graph on the right, which gives the amount of crust generated at a 100 mm/yr opening rate (1 mm = 0.04 in.).





half spreading rate in mm/year



half spreading rate in mm/year

## ***Pacific-Antarctic Ridge: fast-spreading, broad and smooth***




## ***Mid-Atlantic Ridge: slow-spreading, narrow and rough***



In places where spreading is fastest (more than 80 millimeters per year), the ridge has relatively gentle topography and is roughly dome-shaped in cross-section as a result of the many layers of lava that build up over time.

At slow- and ultra-slow spreading centers, the ridge is much more rugged, and spreading is dominated more by tectonic processes rather than volcanism.


*The more prominent ridges and valleys on the flanks are fracture zones (transform fault zones) that were crossed at an oblique angle.*  
(After B. C. Heezen, *The deep-sea floor*, in S. K. Runcorn, ed., *Continental Drift*, Academic Press, 1962)



(a)

Oblique view of fast- and slow-spreading mid-ocean ridges, showing differences in morphology along the ridge.

a) *Fast-spreading East Pacific Rise at 19°S, viewed toward the north.*



(b)




b) *Slow-spreading Mid-Atlantic Ridge at 30°N and the Atlantis transform view toward the northeast.*

*Images made with GeoMapApp software with multibeam sonar data (each with 2× vertical exaggeration).*

*(W. Haxby 2006, GeoMapApp; Marine Geosciences Data Management System, <http://www.GeoMapApp.org/>)*



## TRANSFORM FAULTS




A transform fault or transform boundary, also known as conservative plate boundary since these faults neither create nor destroy lithosphere, is a type of fault whose relative motion is predominantly horizontal in either sinistral or dextral direction.

## TRANSFORM FAULTS

First discovered in the Pacific Ocean by Vacquier (1965): left-lateral offset along the Mendocino and Pioneer faults amount of 1450 km, while the right-lateral offset across the Murray fault is 600 km in the west and only 150 km in the east.


Wilson (1965) termed the faults “Transform” as:  
the lateral displacement across the fault is taken up by transforming it into either the formation of new lithosphere at a terminated ocean ridge segment or lithosphere subduction at a trench.

**The transform faults can form a tectonic plate boundary**






# Global sea floor topography from gravity data derived from satellite altimetry and shipboard depth soundings




SMITH AND SANDWELL, 1996







The Romanche Trench bisects the Mid-Atlantic Ridge just north of the equator at the narrowest part of the Atlantic between Brazil and West Africa.











La faglie principali nella Terra Vittoria Settentrionale e nel Mare di Ross.





# Triple junction



## Dorsale medio-oceanica







- Oceanic plates thin crust (MOHO at about 6 km)
- Oceanic plates thicken as they cool (boundary between convecting and non-convecting mantle deepens)
- Colder (older) plates sink (subduction)
- Examples: Pacific, Nazca, Cocos, Phillipine
- On-land examples of ocean crust: ophiolites
- Oceanic crust emplaced upon continents
- Thickness of 8-10 km
- Ophiolite stratigraphy same worldwide => same processes operate worldwide

## Pillow lava along Juan de Fuca Ridge



The compositions of materials erupted at the mid-ocean ridges are tholeiitic basalts called **mid-ocean ridge basalts (MORB)**.



## Magnetic Susceptibility

$$K = \frac{M}{H}$$

K = magnetic susceptibility


M = induced magnetization in the material

H = applied magnetic field

$$K < 0$$

### DIAMAGNETIC

weak negative susceptibility  
(water, organic matter, quartz,  
feldspars, calcium carbonate)




Quarzo, calcite e dolomite

$$K > 0$$

### PARAMAGNETIC


weak positive susceptibility  
(many Fe-containing minerals and salts,  
eg. biotite, muscovite, pyroxene, olivine,  
ferrous sulfate)



$$K \gg 0$$





### FERROMAGNETIC

strong positive susceptibility  
(pure iron like magnetite and hematite,  
nickel, chromium)



Ossidi di ferro (magnetite  $\text{Fe}_3\text{O}_4$ , ematite  $\alpha\text{-Fe}_2\text{O}_3$ , ilmenite  $\text{FeTiO}_3$ , magnemite  $\gamma\text{-Fe}_2\text{O}_3$  e ulvospinello  $\text{Fe}_2\text{TiO}_4$ ), idrossidi (es. la goethite ( $\alpha\text{-FeO}\cdot\text{OH}$ ), solfuri (es. pirite  $\text{FeS}_2$ ).

## Present magnetic field








Mare di Ross (Antartide) 2002 – M/N Italica

Instruments to measure the earth magnetic field in the ocean:


- magnetometer

- gradiometer composed by two magnetometers to filter time variation in the magnetic field



The magnetic anomalies are numbered as  $C_n$  (Chrono  $n$ ) or  $A_n$  (Anomaly  $n$ ) ( $C_1$  or  $A_1$  is the youngest and  $C_{23}$  or  $A_{23}$  is older).







- 1 - sedimenti non magnetici
- 2 - colate e strato di basalto a cuscini (altamente magnetico: in nero, polarità normale; in bianco, polarità inversa)
- 3 - crosta oceanica a gabbri (debolmente magnetica)


# Geomagnetic Polarity Time Scale (GPTS)

Cande & Kent 1995






**between reversals**




**during a reversal**

NASA computer simulation using the model of Glatzmaier and Roberts. The tubes represent magnetic field lines blue when the field points towards the center and yellow when away. The rotation axis of the Earth is centered and vertical. The dense clusters of lines are within the Earth's core.







## Stato termico dei ridge e piane abissali




**Figure 4-24** The solid lines are isotherms,  $T - T_s$  (°K), in the oceanic lithosphere from Equation (4-125). The data points are the thicknesses of the oceanic lithosphere in the Pacific determined from studies of Rayleigh wave dispersion data. (From A. R. Leeds, L. Knopoff, and E. G. Kausel, Variations of upper mantle structure under the Pacific Ocean, *Science*, **186**, 141–143, 1974.)

# Cooling of oceanic crust






## Basins in the oceans



The numbers indicate contour depths of major ocean basins based on the most shallow, closed, bathymetric contour that defines the basin outline, illustrating that the deepest basins are located in the northwest Pacific.

# Abyssal Plain




Result from the blanketing of the oceanic crust by fine-grained sediments, mainly clay and silt from turbidity currents and from pelagic sediments.

Metallic nodules are common in some areas of the plains, with varying concentrations of metals, including manganese, iron, nickel, cobalt, and copper.



## Total Sediment Thickness of the World's Oceans & Marginal Seas



A digital total sediment thickness database for the world's oceans and marginal seas is being compiled by the National Geophysical Data Center (NGDC), Marine Geology & Geophysics Division. The data are gridded with a spacing of 5 arc-minutes by 5 arc-minutes. Sediment thickness data were compiled from three principle sources: previously published isopach maps; ocean drilling results, both ODP and DSDP; and seismic reflection profiles archived at NGDC as well as seismic data and isopach maps available as part of the IOC's Geological/Geophysical Atlas of the Pacific (GAPA) project.

The distribution of sediments in the oceans is controlled by five primary factors:

- 1) Age of the underlying crust
- 2) Tectonic history of the ocean crust
- 3) Structural trends in basement
- 4) Nature and location of sediment sources, and
- 5) The nature of the sedimentary processes delivering sediments to depocenters






The data values are in meters and represent the depth to acoustic basement. It should be noted that acoustic basement may not actually represent the base of the sediments. These data are intended to provide a minimum value for the thickness of the sediment in a particular geographic region.


<http://www.ngdc.noaa.gov/mgg/sedthick/sedthick.html>



## HOT SPOT

## EXPLANATION


-  **Divergent plate boundaries—**  
Where new crust is generated  
as the plates pull away from  
each other.
-  **Convergent plate boundaries—**  
Where crust is consumed in the  
Earth's interior as one plate  
dives under another.
-  **Transform plate boundaries—**  
Where crust is neither produced  
nor destroyed as plates slide  
horizontally past each other.
-  **Plate boundary zones—Broad  
belts in which deformation is  
diffuse and boundaries are not  
well defined.**
-  **Selected prominent hotspots**



Hotspots are places where molten rock from the earth's mantle is erupting at the surface. They are in the middle of the plate.

Two hypothesis:

- a) the hotspots move relative to the earth;
- b) the hotspots are fixed to the earth.




## ICELAND HOT SPOT

- mid oceanic ridge rises above sea level
- 20 million years old
- some of the most active volcanoes in the world are located in Iceland
- is situated on a hotspot

A detailed map of Iceland lying in the middle of the Atlantic Ocean. It shows the mid Atlantic ridge cutting through Iceland along with the location of its hotspot.

Image taken from


[Iceland Keck](#)




## ICELAND HOT SPOT (or mantle plume)

The starting plume head, **hundreds of degrees hotter than normal mantle**, spreads sideways, incorporates surrounding mantle, and buoyantly uplifts a region roughly 1000 km in diameter to produce a **topographic bulge about 1 km high**.


(Image by D. Müller, University of Sydney).








## HAWAII HOT SPOT



The Hawaiian Ridge-Emperor Seamounts chain:

- extends some 6,000 km,
- composed by 80 volcanoes
- started 70 Ma ago, and sharp bend indicates change of motion at 43 Ma, possibly due to India-Asia collision
- is stationary

## HAWAII HOT SPOT




As the Pacific Plate continues to move west-northwest, the Island of Hawaii will be carried beyond the hotspot by plate motion, setting the stage for the formation of a new volcanic island in its place. In fact, this process may be under way. **Loihi Seamount**, an active submarine volcano, is **forming about 35 km off the southern coast of Hawaii**. Loihi already has risen about 3 km above the ocean floor to within 1 km of the ocean surface. According to the hotspot theory, assuming Loihi continues to grow, it will become the next island in the Hawaiian chain. In the geologic future, Loihi may eventually become fused with the Island of Hawaii, which itself is composed of five volcanoes knitted together-Kohala, Mauna Kea, Hualalai, Mauna Loa, and Kilauea.




# Morphological Features

## Seamounts, volcanic island and guyots


Copyright © McGraw-Hill Companies, Inc. Permission required for reproduction or display.



A





B




C

Subsidence of sea floor





# SEAMOUNT




Seamounts: undersea mountains rising from the bottom of the sea with a minimum elevation of 1,000 meters, that do not break the water's surface.

Seamounts are usually isolated and cone-shaped, are mostly volcanic and therefore found on oceanic crust, formed near mid-oceanic ridges, hotspot and island-arc convergent settings.

A seamount tall enough to break the sea surface is called an oceanic island, e.g., the islands of Hawaii, the Azores and Bermuda.

# Seamount Locations


Kitchingman and Lai 2004



about 60.000 seamounts



# Volcanic Island




Isole canarie




# Gifford Guyot


Faust-Capel basins reconnaissance survey (Tan0713)




# GUYOT




1. Volcanic Island



2. Wave erosion




3. Guyot




# The classification





## Classification of the Marine Zones



### Temperature

$T = 12^{\circ}\text{C}$   
at about 700-1000 m depth  
in the tropic

$T = 4^{\circ}\text{C}$   
at about 4000 m depth  
in the tropic

### Environmental classification:


- Littoral
- sublittoral
- bathyal
- abyssal
- hadal

### Light classification:

- photic
- aphotic


### Pelagic:

- neritic
- oceanic:
  - epipelagic (photic zone)
  - mesopelagic (down to  $T=12^{\circ}\text{C}$ )
  - bathypelagic ( $12^{\circ}\text{C} < T > 4^{\circ}\text{C}$ )
  - abyssalpelagic
  - hadalpelagic





Such a miniscule amount of light penetrates beyond a depth of 200 meters that photosynthesis is no longer possible.





Colore	Frequenza	Lunghezza d'onda
Violetto	668-789 THz	380–450 nm
Blu	631-668 THz	450–475 nm
Ciano	606-631 THz	476-495 nm
Verde	526-606 THz	495–570 nm
Giallo	508-526 THz	570–590 nm
Arancione	484-508 THz	590–620 nm
Rosso	400-484 THz	620–750 nm





# Ancient Oceans



## The origin of the water of the ocean

The water of the paleo-oceans formed on Earth 3.8 billion years ago (the Earth is 4.5 b years old) by two sources:

- outgassing whereby gases are released from molten rock in the mantle of the planet by volcanic activity;
- bombardment by comets and meteorites bringing with them gases which contributed to the Earth's atmosphere (some meteorites are formed by 20% of water).

Some of the gases in the new atmosphere were methane ( $\text{CH}_4$ ), ammonia ( $\text{NH}_3$ ), water vapor ( $\text{H}_2\text{O}$ ), and carbon dioxide ( $\text{CO}_2$ ).

The water on Earth stayed in gaseous form until the planet's surface cooled below 100°C.

At this time, 3.8 billion years ago, water condensed into rain and poured onto the land. Water collected in low lying areas which gradually became the primitive oceans.

At 3.5 billions years ago, the first photosynthetic organisms appeared, and they produced oxigen that enreached the primitive atmosphere.

The geochemical cycles had their beginnings here, with minerals entering the oceans from the land and sky and minerals leaving the oceans through tectonic activity and by evaporation/deposition processes.

These cycles were well established about 1 billion years ago.  
Since then the chemically composition of the oceans has remained constant.

## The **Giapeto Ocean**, between the Laurentia and Baltica continents

- formed in the Cambrian, about 510 milion years ago,
- disappear in the Devonian, about 400 milion years ago.



## Gli oceani nel passato




Map courtesy of CR Scotese, PALEOMAP Project

L'Oceano Tetide (o semplicemente Tetide) era un braccio oceanico disposto in senso Est-Ovest che, nei tempi geologici compresi tra il Permiano ed il Miocene separava l'Africa settentrionale dall'Europa e dall'Asia.

L'apertura dell'Oceano Tetide avvenne circa 250 milioni di anni fa, tra il Permiano ed il Triassico inferiore e portò alla separazione tra un blocco continentale settentrionale (Laurasia) ed uno meridionale (Gondwana). L'allontanamento delle due parti del Pangea proseguì fino al Giurassico, quando i movimenti delle placche tettoniche si invertirono ed iniziò una contrazione dell'Oceano Tetide stesso.

Il movimento dell'Africa era solidale con quello della placca adriatica, che forse ne rappresentava una parte settentrionale. La collisione della placca adriatica con il continente europeo chiuse la Tetide nella regione centrale del Mediterraneo, dando origine alla catena montuosa delle Alpi. Altre microplacche intrappolate tra le due maggiori (africana ed europea) contribuirono a formare altre catene montuose europee orientate generalmente in direzione Est-Ovest, mentre nella zona mediorientale la placca arabica collideva con l'Asia. A completare la chiusura della Tetide, l'India, staccatasi dal continente meridionale di Gondwana durante il Giurassico, si scontrò con l'Asia dando origine alla catena himalayana.


## ATLANTIC OCEAN: initial stage



### Fragmentation of the Pangea (Late Giurassic):

- Opening of the Central Atlantic (A) and the Ligurian-Piedmont Basin (LP – western Tethys)
- Continental rifting of the future Southern Atlantic (R)
- Subduction of the Paleo Tethys (PT) in the Permo-Triassic


# The break up of Pangaea




PERMIAN  
225 million years ago




TRIASSIC  
200 million years ago



JURASSIC  
135 million years ago



CRETACEOUS  
65 million years ago




PRESENT DAY



# General Bathymetric Chart of the Oceans - GEBCO

<http://www.gebco.net>





## International Hydrographic Organization

It is the [inter-governmental organisation](#) representing the [hydrographic](#) community. It enjoys observer status at the [UN](#) where it is the recognised competent authority on hydrographic surveying and [nautical charting](#).

### Functions

The principal work undertaken by the IHO is:

- To bring about a close and permanent association between national hydrographic offices.
- To study matters relating to [hydrography](#) and allied sciences and techniques.
- To further the exchange of nautical charts and documents between hydrographic officers of member governments.
- To tender guidance and advice upon request, in particular to countries engaged in setting up or expanding their hydrographic service.
- To encourage coordination of hydrographic surveys with relevant oceanographic activities.
- To extend and facilitate the application of oceanographic knowledge for the benefit of navigators.
- To cooperate with international organizations and scientific institutions which have related objectives.
- The IHO develops hydrographic and nautical charting standards to be agreed upon by its Member States. All Member States then follow those standards in their surveys, nautical charts, and publications. The almost universal use of the standards means that the products and services from the world's hydrographic and oceanographic offices are increasingly consistent and recognisable for all seafarers and for other users. Much has been done in the field of standardisation since the Bureau (now the IHO) was founded.

## Web sites

<http://www.oceanleadership.org/>

## Paleomagnetism

<http://www.minerva.unito.it/SIS/Paleomagnetismo/paleo4.htm>

## Plate tectonic

<http://www.ucl.ac.uk/EarthSci/people/lidunka/GEOL2014/Geophysics1-%20Plate%20tectonics/PLATE%20TECTONICS.htm>

## Woods Hole Oceanographic Institution

<http://www.whoi.edu/main/ocean-topics>

## NOAA National Geophysical Data Center

<http://www.ngdc.noaa.gov>

## Ocean gravity

[http://topex.ucsd.edu/grav\\_outreach/index.html#natlanticano](http://topex.ucsd.edu/grav_outreach/index.html#natlanticano)



## Bibliography

Global Tectonics, 3rd Edition

Philip Kearey, Keith A. Klepeis, Frederick J. Vine

2009, Wiley-Blackwell