


LECTURE 4

HISTONE METHYLATION AND DNA METHYLATION

LECTURE 4


HISTONE METHYLATION MECHANISMS

HISTONE LYSINE AND ARGININE METHYL TRANSFERASES (HKMETs and HRMETs))



HISTONE LYSINE METHYL TRANSFERASES (HKMETs)

all HKMETs contain a conserved SET domain that catalyzes the methylation of Lysines (K)
(exception Dot1 – no SET domain)




50 SET domain proteins are
Categorized according to sequence
homology
into 6 HKMET subfamilies

1. SET1 family
2. SET2 family
3. SUV39 family
4. EZH family
5. SMYD family
6. PRDM family
7. other SET domain HKMETs

50 SET domain proteins contain many
other protein domains
→ Interaction with other proteins or
DNA

HKMET HRMET SUBSTRATES ON HUMAN HISTONES



HKMETS epigenetic writers are substrate specific and can result in gene repression but also gene activation →→→
The epigenetic reader that binds to the modified histone K residue at the individual histone tail makes the difference

Effect on gene activity Best studied examples of histone methylation:

	Substrate	Histone lysine methyltransferases
activation	H3K4	SET9, SET1, MLL, ASH1L, SMYD3, PRDM9, SETMAR
repression	H3K9	SUV39H1, SUV39H2, EHMT1, EHMT2, SETDB1, PRDM2, ASH1L
repression	H3K27	EZH2, EHMT2
activation	H3K36	NSD1, SETD2/HYPB, SETMAR
activation	H3K79	DOT1L
repression	H4K20	SET8, SUV420H1, SUV420H2, NSD1, ASH1L

Fig. 1. Histone modifications. (A) The modifications on human histones include methylation (Me) on arginine and lysine residues, acetylation (Ac) on lysine residues, phosphorylation (P) on serine and threonine residues and ubiquitination (Ub) on lysine residues. (B) The enzymes responsible for methylation of human histone lysine residues are listed according to their target sites. Histone lysine methyltransferases (HKMTs) are very specific but redundant in several cases.

HISTONE MODIFICATIONS AND EPIGENETIC READERS

Protein domains that bind to histone modifications




Figure 1 Readers of histone PTMs. Recognition of the methylated (me) lysine, methylated (me) arginine, acetylated (ac) lysine and phosphorylated (ph) serine and threonine residues of the N-terminal histone H3 tail by indicated readers.

A large number of proteins contain these protein domains:

→ High complexity in gene regulation that


→ Creation of large numbers of EPIGENOMES

Table 1 Histone readers and their target PTMs


Recognition of	Reader	Histone PTM
Methyllysine	ADD	H3K9me3
	Ankyrin	H3K9me2, H3K9me1
	BAH	H4K20me2
	Chromo-barrel	H3K36me3, H3K36me2, H4K20me1, H3K4me1
	Chromodomain	H3K9me3, H3K9me2, H3K27me3, H3K27me2
	DCD	H3K4me3, H3K4me2, H3K4me1
	MBT	H3Kme1, H3Kme2, H4Kme1, H4Kme2
	PHD	H3K4me3, H3K4me2, H3K9me3
	PWWP	H3K36me3, H4K20me1, H4K20me3, H3K79me3
	TTD	H3K4me3, H3K9me3, H4K20me2
Methylarginine	Tudor	H3K36me3
	WD40	H3K27me3, H3K9me3
	zf-CW	H3K4me3
Acetyllysine	ADD	H4R3me2s
	Tudor	H3Rme2, H4Rme2
	WD40	H3R2me2
	Bromodomain	H3Kac, H4Kac, H2AKac, H2BKac
Phosphoserine or phosphothreonine	DBD	H3KacKac, H4KacKac
	DPF	H3Kac
	Double PH	H3K56ac
Unmodified histone	14-3-3	H3S10ph, H3S28ph
	BIR	H3T3ph
Unmodified histone	Tandem BRCT	H2AXS139ph
	ADD	H3un
	PHD	H3un
	WD40	H3un

THE SET DOMAIN

THE SET DOMAIN



THE ACTIVE SITE IN THE SET DOMAIN



THE BIOCHEMISTRY OF HISTONE LYSINE METHYLATION

The source of the methyl group is **S-adenosyl-l-methionine (AdoMet) or (SAM)**, which is converted to **S-adenosyl-l-homocysteine (AdoHcy)** in the reaction.



S-Adenosyl methionine is a common cosubstrate involved in methyl group transfers, transsulfuration, and aminopropylation.

SAM = enzymatic cofactor


SAM is after ATP the most commonly used cofactor used by the cell

Although these anabolic reactions occur throughout the body, most **SAM is produced and consumed in the liver**. More than 40 methyl transfers from SAM are known, to various substrates such as nucleic acids, proteins, lipids and secondary metabolites. It is made from adenosine triphosphate (ATP) and methionine by methionine adenosyltransferase. SAM was first discovered in Italy by Giulio Cantoni in 1952.

S-adenosyl-l-methionine (AdoMet) or (SAM),



THE BIOCHEMISTRY OF HISTONE LYSINE METHYLATION



Catalytic mechanism

In order for the reaction to proceed, S-Adenosyl methionine (SAM) and the lysine residue of the substrate histone tail must first be bound and properly oriented in the catalytic pocket of the SET domain. Next, a nearby tyrosine residue deprotonates the ϵ -amino group of the lysine residue. The lysine chain then makes a nucleophilic attack on the methyl group on the sulfur atom of the SAM molecule, transferring the methyl group to the lysine side chain.


ENZYMATIC ASSAY TO DETECT KMTase ACTIVITY

Experiment:

Overexpression of **myc-tagged-SUV39H1 KMT** in HeLa cells

Use an antibody to immunoprecipitate SUV39H1 → high concentration of SUV39H1


Incubate Immunoprecipitate with purified histones and S-adenosyl-[methyl-¹⁴C]-L-methionin as methyl donor



SET – domain is required for histone methyl transferases activity


- The SET domain of the SUV39H1 is required for histone methyltransferase activity and this enzyme methylates H3 at Lys9

S-adenosyl-L-methionine (AdoMet) or (SAM),




HISTONE LYSINES CAN BE MONO- DI- AND TRI-METHYLATED

lysine methylation




**ARE THERE KMTs THAT CREATE SPECIFIC METHYLATION LEVELS
(mono-methylation, di-methylation, tri-methylation?)**

**SUBSTRATE SPECIFICITY OF HISTONE METHYL TRANSFERASES:
AN EXAMPLE: THE HKMT SUV39H1**



Suv39dn cells



H3K9me1: increased and pattern similar to wt H3K9me3 (chromocenter)
H3K9me2: similar to wt
H3K9me3: strongly reduced; lost at chromocenters



Suv39dn cells

H3K9me1: increased compared to wt
H3K9me2: similar to wt
H3K9me3: strongly reduced

SUBSTRATE SPECIFICITY OF HISTONE METHYL TRANSFERASES: AN EXAMPLE: THE HKMT SUV39H1




The H3K9 specific KMTs G9a and GLP are the major H3K9me1 and H3K9me2 methyltransferases

The H3K9 specific KMTases Suv39h1 and Suv39h2 are the major H3K9me3 methyltransferases

Suv39h1 and Suv39h1 work best on H3K9me2

H3K9 methylation



EPIGENETIC READERS AN EXAMPLE: H3K9me3 and HP1





Table 1 Histone readers and their target PTMs

Recognition of	Reader	Histone PTM
Methyllysine	ADD	H3K9me3
	Ankyrin	H3K9me2, H3K9me1
	BAH	H4K20me2
	Chromo-barrel	H3K36me3, H3K36me2, H4K20me1, H3K4me1
	Chromodomain	H3K9me3, H3K9me2, H3K27me3, H3K27me2
	DCD	H3K4me3, H3K4me2, H3K4me1
	MBT	H3Kme1, H3Kme2, H4Kme1, H4Kme2
	PHD	H3K4me3, H3K4me2, H3K9me3
	PWWP	H3K36me3, H4K20me1, H4K20me3, H3K79me3
	TTD	H3K4me3, H3K9me3, H4K20me2
Methylarginine	Tudor	H3K36me3
	WD40	H3K27me3, H3K9me3
	zf-CW	H3K4me3
Methylarginine	ADD	H4R3me2s
	Tudor	H3Rme2, H4Rme2
	WD40	H3R2me2
	Acetyllysine	Bromodomain
DBD		H3KacKac, H4KacKac
DPF		H3Kac
Double PH		H3K56ac
Phosphoserine or phosphothreonine	14-3-3	H3S10ph, H3S28ph
	BIR	H3T3ph
	Tandem BRCT	H2AXS139ph
Unmodified histone	ADD	H3un
	PHD	H3un
	WD40	H3un


EPIGENETIC READERS – IN VIVO EVIDENCE
 AN EXAMPLE: HP1 has high affinity for H3K9me3



A chromodomain (chromatin organization modifier) is a protein structural domain of about 40-50 amino acid residues commonly found in proteins associated with the remodeling and manipulation of chromatin. The domain is highly conserved among both plants and animals, and is represented in a large number of different proteins in many genomes, such as that of the mouse. Chromodomain-containing proteins also bind methylated histones and appear in the RNA-induced transcriptional silencing complex.

YELLOW: histone tail


RODs: Interacting aminoacids of HP1



Loss of *Suv39h1/2*:
 reduced
 (2 slides earlier)
 and delocalized
 (this slide) H3K9me3.


Consequence:
 HP1 is also
 delocalized!!!!

HISTONE LYSINE AND ARGININE METHYL TRANSFERASES (HKMETs and HRMETs))



HISTONE ARGININE METHYL TRANSFERASES (HRMETs)



Family of PRMTs: Protein Arginine (R) methyl-transferases



PRMTs have a MTase domain that is Different from the SET domain!!!

Conserved core:

- MTase domain: catalyzes methylation of R
- Beta barrel domain: Important for dimerization of PRMTs



PRMTs

- Type I PRMTs: need to dimerize to be functional
- Type II PRMTs: form larger complexes – dimers interact to form tetramers, other proteins can interact

THE BIOCHEMISTRY OF HISTONE ARGININE METHYLATION

Methyl transfer reactions catalyzed by AdoMet-dependent PRMTs.

Example: PRMT1


The reacting arginine substrate acts by nucleophil attack on the methyl group present on AdoMet.

The reaction has been proposed to involve 3 key conserved residues in the active site of PRMT1: Arg-54, Glu-144, and Glu-153.

Arg-54 and **Glu-144** help to properly position the substrates for the nucleophilic attack


Glu-153 is hypothesized to play a role in increasing the nucleophilicity of the guanidinium moiety of the substrate via enhanced electronic effects.

Glu-144 has also been postulated to act as the active site base, abstracting a proton from the reacting arginine either during or immediately after methyl transfer.




THE BIOCHEMISTRY OF HISTONE ARGININE METHYLATION

PRMT activity requires :
substrate containing R,
AdoMet (SAM) as enzymatic cofactor
PRMT



S-adenosyl-L-methionine (AdoMet) or (SAM),



PRMTs CATALYZE MONO and DIMETHYLATION
- Not trimethylation -

PRMT SUBSTRATES AND BIOLOGICAL ACTIVITY

PRMTs can act as activators and repressors of gene expression

PRMTs:		Type	Histone substrate	Biological Function
PRMT1		I	H4R3	<u>NR, chromatin dynamic, transcription activation</u>
PRMT2		?		Coactivator for ER, Cellular proliferation
PRMT3		I		ribosomal biosynthesis
PRMT4		I	H3R2, H3R17 (Rare)	<u>NR, transcription activation, epigenetic reprogram in embryos</u>
PRMT5		II	H4R3; H3R8	<u>Stem cell function, transcription repression, repressive chromatin</u>
PRMT6		I	H3R2	<u>Repressive chromatin, supression of H3K4 methylation</u>
PRMT7		II	H2A, H4R3	Potentiating DNMT3 binding, regulation of imprinting genes
PRMT8		I	H4?	?
PRMT9 Isoform 4		II	H4, H2A	?
PRMT10		?		?
PRMT11		?		?

PRMTs epigenetic writers, are substrate specific and can result in gene repression but also gene activation

→→→

The epigenetic reader that binds to the modified histone R residue at the individual histone tail makes the difference

HISTONE MODIFICATIONS AND EPIGENETIC READERS

Protein domains that bind to histone modifications




Figure 1 Readers of histone PTMs. Recognition of the methylated (me) lysine, methylated (me) arginine, acetylated (ac) lysine and phosphorylated (ph) serine and threonine residues of the N-terminal histone H3 tail by indicated readers.


A large number of proteins contain these protein domains:
 → High complexity in gene regulation that
 → Creation of large numbers of EPIGENOMES

Table 1 Histone readers and their target PTMs

Recognition of	Reader	Histone PTM
Methyllysine	ADD	H3K9me3
	Ankyrin	H3K9me2, H3K9me1
	BAH	H4K20me2
	Chromo-barrel	H3K36me3, H3K36me2, H4K20me1, H3K4me1
	Chromodomain	H3K9me3, H3K9me2, H3K27me3, H3K27me2
	DCD	H3K4me3, H3K4me2, H3K4me1
	MBT	H3Kme1, H3Kme2, H4Kme1, H4Kme2
	PHD	H3K4me3, H3K4me2, H3K9me3
	PWWP	H3K36me3, H4K20me1, H4K20me3, H3K79me3
	TTD	H3K4me3, H3K9me3, H4K20me2
	Tudor	H3K36me3
WD40	H3K27me3, H3K9me3	
zf-CW	H3K4me3	
Methylarginine	ADD	H4R3me2s
	Tudor	H3Rme2, H4Rme2
	WD40	H3R2me2
Acetyllysine	Bromodomain	H3Kac, H4Kac, H2AKac, H2BKac
	DBD	H3KacKac, H4KacKac
	DPF	H3Kac
	Double PH	H3K56ac
Phosphoserine or phosphothreonine	14-3-3	H3S10ph, H3S28ph
	BIR	H3T3ph
Unmodified histone	Tandem BRCT	H2AXS139ph
	ADD	H3un
	PHD	H3un
	WD40	H3un

AN EXAMPLE FOR PRMT FUNCTION IN TRANSCRIPTIONAL ACTIVATION


p53 is a tumor-suppressor that acts as transcription factor



- H4R3me stimulates subsequent acetylation of H4
- p53 binds transcription factor binding site in promoter and brings p300.
- p300 acetylates promoter regions
- PRMT4 has affinity for acetylated histones H3R17me
- (note: PRMT1, 4 do not contain a bromo domain → other proteins mediate affinity to acetylated histones)


→ FULL TRANSCRIPTIONAL ACTIVATION

LYSINE AND ARGININE METHYLATION IN HISTONES IS REVERSIBLE




Nature Reviews | Genetics


The **Jumonji N (JmjN)** and **Jumonji C (JmjC)** domains are two non-adjacent domains which have been identified in the jumonji family of transcription factors. Although it was originally suggested that the JmjN and JmjC domains always co-occur and might form a single functional unit within the folded protein, the JmjC domain was later found without the JmjN domain in organisms from bacteria to human. The JmjC domain is the best studied domain that mediated histone demethylation - is conserved from yeast to human




1. LSD1 (KDM1A): demethylation by oxidation
2. Big family of Jumonji domain containing proteins: hydroxylation



LYSINE AND ARGININE METHYLATION IN HISTONES IS REVERSIBLE



Arginine-demethylases are less well studied



LECTURE 4


DNA METHYLATION

DNA METHYLATION CONTROLS GENE EXPRESSION

FACTS:


1. DNA methylation is created at CpG di-nucleotide motifs
2. An accumulation of CpG is called "CpG island" (CGI)
3. CpG islands are enriched at promoters and sequence elements that are important for gene expression control. In some cases, CpG islands can be also located in distant locations.
4. *CpG methylation (=“DNA methylation”) is directly linked with stable gene silencing*

ADVANTAGE OF DNA METHYLATION AT CpG



CpGs are self-complementary
Di-nucleotide in paired stand also contains methylation
Methylation patterns can be maintained during DNA replication

Epigenetic modification that is imposed on genomic DNA




CYTOSINE

METHYL-CYTOSINE

DNA METHYLTRANSFERASES CATALYZE DNA METHYLATION

DNA methyltransferases transfer a methyl-group from AdoMet (SAM) to Cytosine located in a CpG dinucleotide



S-adenosyl-L-methionine (AdoMet) or (SAM),



The source of the methyl group is S-adenosyl-L-methionine (AdoMet) or (SAM), which is converted to S-adenosyl-L-homocysteine (AdoHcy) in the reaction.

S-Adenosyl methionine is a common cosubstrate involved in methyl group transfers, transsulfuration, and aminopropylation.

SAM = enzymatic cofactor


SAM is after ATP the most commonly used cofactor used by the cell

Although these anabolic reactions occur throughout the body, most SAM-e is produced and consumed in the liver. More than 40 methyl transfers from SAM-e are known, to various substrates such as nucleic acids, proteins, lipids and secondary metabolites. It is made from adenosine triphosphate (ATP) and methionine by methionine adenosyltransferase. SAM was first discovered in Italy by Giulio Cantoni in 1952.




Mapping DNA methylation at CpG islands BISULFITE SEQUENCING


Methylation of cytosine at CpG dinucleotides is an important epigenetic regulatory modification in many eukaryotic genomes. **DNA methylation was found to be located genome-wide with a pattern of low methylation in proximity to promoters and high genebody-methylation in highly-expressed genes → methylation pattern can identify transcribed DNA (gene)**



Bisulfite conversion: C → U conversion





DMS = Dimethyl sulfate



Treatment of DNA with bisulfite converts cytosine residues to uracil, but leaves 5-methylcytosine residues unaffected. Thus, bisulfite treatment introduces specific changes in the DNA sequence that depend on the methylation status of individual cytosine residues, yielding single-nucleotide resolution information about the methylation status of a segment of DNA.


Mapping DNA methylation at CpG islands BISULFITE SEQUENCING

Genomic DNA



Bisulfite conversion

DNA for Sequencing



Compare with genomic sequence
C→U sequence change = DNA methylation
C→C no sequence change = no DNA methylation

Mapping DNA methylation at CpG islands BISULFITE SEQUENCING


An example:

hTERT encodes the telomerase gene
hTERT elongates telomeres thereby
protecting cancer cells from
replicative senescence


→ Cancer cells do not age and can proliferate
forever!!

→ Therefore 90% of human cancers express
telomeres; but telomerase is silenced by
DNA methylation in differentiated cells.

→ Bisulfite sequencing showed that cancer
cells have a de-methylated CpG island
located at the hTERT promoter. CpG islands
can overlap with the 1st intron of the
gene!!!!



CANCER CELL
CpG
island
not methylated



NORMAL CELL
CpG
island
is methylated (**)

Mapping DNA methylation at CpG islands of individual genes

BISULFITE SEQUENCING

Prepare DNA from normal cell and cancer cell

Purify DNA and perform bi-sulfite conversion (DMS)
(Unmethylated C → U; Methylated C → C)

Amplify your region of interest = CpG islands
in the TERT promoter

Purify DNA fragment obtained by PCR



Clone fragment into Plasmid

Transform bacteria with plasmid (one bacteria receives only one plasmid!!!)

Purify amplified plasmids from 10-15 individual bacterial cultures

Sequence inserts using a primer that anneals to the vector DNA,
adjacent to the insertion site of the PCR product
REMEMBER: only a single type of plasmid is sequenced
This refers to a single type of molecule.

To have a good representation, you need to
sequence at least 10-15 clones



Mapping DNA methylation at CpG islands BISULFITE SEQUENCING

Prepare DNA from normal cell and cancer cell

Purify DNA and perform bi-sulfite conversion

Amplify your region of interest = CpG islands in the TERT promoter

Purify DNA fragment obtained by PCR

Clone fragment into Plasmid

Transform bacteria with plasmid

Purify amplified plasmids

Sequence inserts using a primer that anneals to the vector


DNA, adjacent the insertion site of the PCR product

REMEMBER: only a single type of plasmid is sequenced

This refers to a single type of molecule.

To have a good representation, you need to sequence at least 10 clones

HOW DO I HAVE TO DESIGN THE PRIMER FOR PCR OF BISULFITE CONVERTED DNA??



C Cytosine not in CpG site C G CpG site

Compare with genomic sequence

Bisulfite conversion: C → U sequence change = DNA methylation

C → C no sequence change = no DNA methylation


Mapping DNA methylation at CpG islands BISULFITE SEQUENCING

An example:

hTERT encodes the telomerase gene
Tert elongates telomeres thereby
protecting cancer cells from
replicative senescence
→ Cancer cells do not age and can
proliferate forever!!

→ Therefore 90% of human cancers
express telomeres; but telomerase is
silenced by DNA methylation in
differentiated cells.

→ Bisulfite sequencing showed that
cancer cells have a de-methylated
CpG island located at the hTERT
promoter



Mapping DNA methylation at CpG islands

METHYLATED DNA IMMUNOPRECIPITATION: METHYL-DIP

Methyl-DIP works similar to ChIP:

2 experimental samples:

- Control (normal cells)
- Experimental sample (cancer cells)

Prepare DNA – carries DNA methylation marks of CpG

Sonicate DNA

Immunoprecipitation using an anti-methyl-CpG specific


Antibody (monoclonal, discriminates between CpG and met-CpG)

Washing of precipitate


DATA ACQUISITION

Main methods to analyse DNA methylation

1. PCR on specific CpG islands of interest
2. Differential labelling (Cy3-control; Cy5-cancer methyl-DNA) followed by hybridization to genome array
3. Massive parallel sequencing (different approaches)




Mapping DNA methylation genome wide: REDUCED REPRESENTATION BISULFITE SEQUENCING (RRBS)




Reduced representation bisulfite sequencing (RRBS) is an efficient and high-throughput technique used to analyze the genome-wide methylation profiles on a single nucleotide level. This technique combines restriction enzymes and bisulfite sequencing in order to enrich for the areas of the genome that have a high CpG content. Due to the high cost and depth of sequencing needed to analyze methylation status in the entire genome. **The fragments that comprise the reduced genome but still includes the majority of promoters, as well as regions such as repeated sequences that are difficult to profile using conventional bisulfite sequencing approaches.**

Mapping DNA methylation genome wide: REDUCED REPRESENTATION BISULFITE SEQUENCING (RRBS)



The color key from light blue to dark blue indicates the DNA methylation level from low to high, respectively. The white regions in the left panels indicate a lack of DNA methylation information. The red bars in the right panel represent the average DNA methylation level of the corresponding genomic region. The DNA methylation levels were calculated and presented based on 30 kb windows, only if these windows have more than 5 CpG sites covered.

Mapping DNA methylation genome wide: REDUCED REPRESENTATION BISULFITE SEQUENCING (RRBS)



Enzyme digestion: First, genomic DNA is digested using a methylation-insensitive restriction enzyme. It is integral for the enzymes to not be influenced by the methylation status of the CpGs (sites within the genome where a cytosine is next to a guanine) as this allows for the digestion of both methylated and unmethylated areas. *MspI* is commonly used. This enzyme targets 5'CCGG3' sequences and cleaves the phosphodiester bonds upstream of CpG dinucleotide. When using this particular enzyme, each fragment will have a CpG at each end. This digestion results in DNA fragments of various sizes.

End repair and A-tailing: Due to the nature of how *MspI* cleaves double stranded DNA, this reaction results in strands with sticky ends. End repair is necessary to fill in the 3' terminal of the ends of the strands. The next step is adding an extra adenosine to both the plus and minus strands. This is referred to as A-Tailing and is necessary for adapter ligation in the subsequent step. End repair and A-Tailing is done within the same reactions, with dCTP, dGTP and dATP deoxyribonucleotides. In order to increase the efficiency of A tailing, the dATPs are added in excess in this reaction.

Sequence adapters: Methylated sequence adapters are ligated to the DNA fragments. The methylated adapter oligonucleotides have all cytosines replaced with 5'methyl-cytosines, in order to prevent the deamination of these cytosines in the bisulfite conversion reaction. For reactions to be sequenced using Illumina sequencers, the sequence adapters are used to hybridize to the adapters on the flow cell.

Fragment purification: The desired size of fragments is then selected to be purified. The different sizes of the fragments are separated using gel electrophoresis and are purified using gel excising. According to Gu et al., DNA fragments of 40-220 base pair are representative of the majority of promoter sequences and CpG islands[2]

Bisulfite conversion: The DNA fragments are then bisulfite converted, which is a process that deaminates unmethylated cytosine into a uracil. The methylated cytosines remain unchanged, due to the methyl group protecting them from the reaction.

PCR amplification: The bisulfite converted DNA is then amplified using PCR with primers that are complementary to the sequence adapters.

PCR purification: Before sequencing, the PCR product must be free of unused reaction reagents such as unincorporated dNTPs or salts. Thus, a step for PCR purification is required. This can be done by running another electrophoresis gel or by using kits designed specifically for PCR purification.

Sequencing: The fragments are then sequenced. When RRBS was first developed, Sanger sequencing was initially used. Now, next generation sequencing approaches are used. For Illumina sequencing, 36-base single-end sequencing reads are most commonly performed.


Sequence alignment and analysis: Due to the unique properties of RRBS, special software is needed for alignment and analysis. Using *MspI* to digest genomic DNA results in fragments that always start with a C (if the cytosine is methylated) or a T (if a cytosine was not methylated and was converted to a uracil in the bisulfite conversion reaction). This results in a non-random base pair composition. Additionally, the base composition is skewed due to the biased frequencies of C and T within the samples. Various software for alignment and analysis is available, such as Maq, BS Seeker, Bismark or BSMAP. Alignment to a reference genome allows the programs to identify base pairs within the genome that are methylated.

DNA methyltransferases methylate DNA

FACTS:

1. DNA methylation is created at CpG di-nucleotide motifs
2. An accumulation of CpG genes is called CpG island
3. CpG islands are enriched at promoters and other, more distant sequence elements that are important for gene expression control
4. CpG methylation (=“DNA methylation”) is linked with Stable gene silencing

ADVANTAGE OF DNA METHYLATION AT CpG



CpG Di-nucleotide are self-complementary

PROPOSED MODEL:

Methylation patterns can be maintained during DNA replication





Figure 1. Cytosine methylation in DNA. (A) Addition of a methyl group, CH_3 (red), at the five position of the cytosine pyrimidine ring (black arrow) does not sterically interfere with GC base pairing (blue lines). DNA methyltransferases associate covalently with the carbon 6 position (straight green arrow) during methyl group transfer. (B) A model of B-form DNA methylated at cytosines in two self-complementary CpG sequences. The paired methyl moieties (magenta and yellow) lie in the major groove of the double helix.

De-novo DNMTs → place new DNA methylation

Maintenance DNMTs → propagate methylation after replication

DNA methyl transferases methylate DNA

Maintenance of DNA methylation patterns by DNMTs




Figure 2. De novo methylation and maintenance methylation of DNA. A stretch of genomic DNA is shown as a line with self-complementary CpG pairs marked as vertical strokes. Unmethylated DNA (*top*) becomes methylated “de novo” by Dnmt3a and Dnmt3b to give symmetrical methylation at certain CpG pairs. On semiconservative DNA replication, a progeny DNA strand is base-paired with one of the methylated parental strands (the other replication product is not shown). Symmetry is restored by the maintenance DNA methyltransferase, Dnmt1, which completes half-methylated sites, but does not methylate unmodified CpGs.

Biochemical discovery of maintenance DNMT1:

Cell extract + DNA containing CpG repeats + ^{14}C labelled -CH₃ in AdoMet (SAM) → radioactive -CH₃ transferred to DNA

NOTE:


This enzyme is 7– to 100-fold more active on hemimethylated DNA as compared with unmethylated substrate *in vitro*

Next step: Purification of enzymatic activity: 200kDa complex containing **DNMT1**

Discovery of de novo DNMTs:

Sequence of DNMT1 was used to look for genes with similar sequence (sequence homology)

- Discovery of de-novo DNMTs that work efficiently work on un-methylated DNA (DNMT3a, 3b)
- De-novo DNMTs cannot efficiently methylate hemi-methylated DNA



LOSS OF DNA METHYLTRANSFERASES IS LETHAL DURING EMBRYONIC MOUSE DEVELOPMENT

Table 1. Function of mammalian DNA methyltransferases

DNA methyltransferase	Species	Major activity	Major phenotypes of loss of function
Dnmt1	Mouse	Maintenance methylation of CpG	Genome-wide loss of DNA methylation, embryonic lethality at embryonic day 9.5 (E9.5), abnormal expression of imprinted genes, ectopic X-chromosome inactivation, activation of silent retrotransposon. In cancer cell lines, it leads to cell cycle arrest and mitotic defects.
Dnmt3a	Mouse	De novo methylation of CpG	Postnatal lethality at 4–8 wk, male sterility, and failure to establish methylation imprints in both male and female germ cells
Dnmt3b	Mouse	De novo methylation of CpG	Demethylation of minor satellite DNA, embryonic lethality around E14.5 days with vascular and liver defects. (Embryos lacking both Dnmt3a and Dnmt3b fail to initiate de novo methylation after implantation and die at E9.5.)
DNMT3B	Human	De novo methylation of CpG	ICF syndrome: immunodeficiency, centromeric instability, and facial anomalies. Loss of methylation in repetitive elements and pericentromeric heterochromatin.





Figure 4. Mammalian DNA methyltransferases. The catalytic domains of Dnmt1, Dnmt2, and the Dnmt3 family members are conserved (the signature motifs, I, IV, VI, IX, and X, are most conserved in all cytosine methyltransferases), but there is little similarity among their amino-terminal regulatory domains. Domain abbreviations: PCNA, PCNA-interacting domain; NLS, nuclear localization signal; RFT, replication foci-targeting domain; CXXC, a cysteine-rich domain implicated in binding DNA sequences containing CpG dinucleotides; BAH, bromo-adjacent homology domain implicated in protein-protein interactions; PWWP, a domain containing a highly conserved “proline-tryptophan-tryptophan-proline” motif involved in heterochromatin association; ATRX, an ATRX-related cysteine-rich region containing a C2-C2 zinc finger and an atypical PHD domain implicated in protein-protein interactions.

DNA METHYLATION IS ABUNANT IN THE GENOME AND ISSUBJECTED TO DRAMATIC ALTERATIONS DURING EMBRYOGENESIS



DNA methylation levels are high in fertilized Oocytes that contain the paternal and maternal genome (carries characteristic methylation patterns)

Paternal and maternal methylation patterns are rapidly erased (exception: imprinted genes maintain paternal and maternal methylation information). → the paternal and maternal methylation epigenome is cancelled

DNA methylation levels remain low during the first cell division events until the blastocyst stage

In the blastocyst stage cell differentiation programs are activated and genes need to be regulated on the epigenetic level → DNA methylation is increasing (loss of DNMT1, DNMT3a or DNMT3b is lethal → establishment and maintenance of DNA methylation is impaired)

70%- 80% of CpG dinucleotides are methylated in the genome

70%- 80% of CpG di-nucleotides are methylated in the human genome!

Remember only 2% of the genome encode for mRNAs

98% is noncoding DNA that contains a large proportion of transposable elements, repeat sequences, etc...

ON THE SINGLE GENE LEVEL:

CpG islands (CGIs) are short sequences stretches with variable DNA methylation that regulate promoter activity

NOTE: single CpGs are generally hyper methylated (60-90%)

CpG islands are differentially methylated, but are generally demethylate


CpG islands (CGIs) have a length of ca. 1kb

60% of human genes are controlled by CGIs containing promoters that allow tissue/cell specific gene expression

CpG islands can overlap with the first exon (methylation level in 1st exon is good predictor of gene expression)

CpG islands located <2kb from promoter: shores

CpG islands located <2-4kb from promoter: shores



TRANSCRIPTIONAL REGULATION BY METHYL-DNA BINDING PROTEINS

Interference with transcription factor binding





Figure 1. Cytosine methylation in DNA. (A) Addition of a methyl group, CH_3 (red), at the five position of the cytosine pyrimidine ring (black arrow) does not sterically interfere with GC base pairing (blue lines). DNA methyltransferases associate covalently with the carbon 6 position (straight green arrow) during methyl group transfer. (B) A model of B-form DNA methylated at cytosines in two self-complementary CpG sequences. The paired methyl moieties (magenta and yellow) lie in the major groove of the double helix.

Methylated DNA obtains different structure:
Transcription factors cannot bind anymore
→ DNA methylation sensitive transcription factors



Example: CTCF

Unmethylated DNA CTCF binds → activation of expression

Methylated DNA: CTCF does not bind → no activation

Note: CTCF is a major epigenetic regulator that is involved in controlling genomic imprinting, enhance activation,...

EPIGENETIC READERS OF DNA METHYLATION

Transcriptional regulation by methyl-dna binding proteins

Table 2. Functions of methyl-CpG binding proteins

MBP	Major activity	Species	Major phenotypes of loss-of-function mutations
MeCP2	Binds mCpG with adjacent run AT-rich Transcriptional repressor	Mouse	Delayed onset neurological defects including inertia, hind-limb claspings, nonrhythmic breathing, and abnormal gait. Postnatal survival ~10 wk.
MECP2	Binds mCpG with adjacent AT run Transcriptional repressor	Human	Heterozygotes suffer from Rett syndrome, a profound neurological disorder characterized by apraxia, loss of purposeful hand use, breathing irregularities, and microcephaly
Mbd1	Binds mCpG via MBD; Mouse a major splice form is also able to bind CpG via a CxxC domain	Mouse	No overt phenotype, but subtle defects in neurogenesis detected
Mbd2	Binds mCpG Transcriptional repressor	Mouse	Viable and fertile, but show reduced maternal nurturing behavior. Defective gene regulation in T-helper cell differentiation leading to altered response to infection. Highly resistant to intestinal tumorigenesis.
Mbd3	Core component of NuRD corepressor complex Does not show strong binding to mCpG	Mouse	Early embryonic lethal
Mbd4	DNA repair protein that binds mCpG and T:G mismatches at mCpG sitesThymine DNA glycosylase that excises T from T:G mismatches	Mouse	Viable and fertile. three- to fourfold increase in mutations at CpG sites. Increased susceptibility to intestinal cancer correlates with C to T transitions within the <i>Apc</i> gene. Mbd4 functions to minimize the mutability of 5-methylcytosine.
Kaiso	Binds mCGmCG and CTGCNA Transcriptional repressor	Mouse	No overt phenotype. Small but significant delay in tumorigenesis on Min background.

Several proteins were identified to have affinity to methylated CpG but do not have affinity to unmethylated CpG → mediate transcriptional silencing

→ CpG METHYL BINDING DOMAIN PROTEIN (MBD) FAMILY : MeCP1, MeCP2, Mbd1, Mbd2, Mbd2, Mbd4

→ Kaiso (unrelated protein)




Figure 8. Proteins that bind methyl-CpG. Five members of the MBD protein family are aligned at their MBD domains (purple). Other domains are labeled and include TRD; CXXC domains, which are zinc fingers, some of which are implicated in binding to nonmethylated CpG; GR repeats that may bind; a T:G mismatch glycosylase domain that is involved in repair of 5mC deamination. Kaiso lacks the MBD domain, but binds methylated DNA via zinc fingers (orange) and possesses a POB/BTB domain that is shared with other transcriptional repressors. Domain abbreviations: MBD, methyl-CpG binding domain; TRD, transcriptional repression domain; POZ, poxvirus and zinc finger, a protein-protein interacting domain.

How does MeCP2 effect the brain function?

- Through it's job as a reader of epigenetic bookmarks
- The wide array of functions that MeCP2 performs ALL contribute to Rett syndrome.
- The different mutations have different effects on the presentation of the disease.
- In addition since each person is different based on their personal epigenetics, the disease will be individual as well.





Figure 1 Composition of MeCP2: gene structure, splicing patterns and putative functional domains

(A) Splicing patterns generating the two mRNA isoforms of *MECP2*, *_e1* and *_e2*. The two isoforms generate two protein isoforms of MeCP2 with differing N-termini due to the use of alternative translation start sites (bent arrows) and the absence or presence of exon 2 in the transcript. **(B)** Apart from the N-terminus, both MeCP2 isoforms are identical and contain several functionally distinct domains: NTD, N-terminal domain; MBD, methylated DNA-binding domain; ID, interdomain; TRD, transcription repression domain; CTD, C-terminal domain; NLS; nuclear localization signals. Locations of seven of the most common point mutations in RTT are indicated (◆). Below each domain are indicated major (bold) and other (grey) interactors and functions. HP1, heterochromatin protein 1; TFIIIB, transcription factor IIB; YB1, Y-box-binding protein 1.

EPIGENETIC READERS OF DNA METHYLATION

Transcriptional regulation by methyl-dna binding proteins

Table 2. Functions of methyl-CpG binding proteins

MBP	Major activity	Species	Major phenotypes of loss-of-function mutations
MeCP2	Binds mCpG with adjacent run AT-rich	Mouse	Delayed onset neurological defects including inertia, hind-limb claspings, nonrhythmic breathing, and abnormal gait. Postnatal survival ~10 wk.
MECP2	Binds mCpG with adjacent AT run	Human	Heterozygotes suffer from Rett syndrome, a profound neurological disorder characterized by apraxia, loss of purposeful hand use, breathing irregularities, and microcephaly
Mbd1	Binds mCpG via MBD; a major splice form is also able to bind CpG via a CxxC domain	Mouse	No overt phenotype, but subtle defects in neurogenesis detected
Mbd2	Binds mCpG	Mouse	Viable and fertile, but show reduced maternal nurturing behavior. Defective gene regulation in T-helper cell differentiation leading to altered response to infection. Highly resistant to intestinal tumorigenesis.
Mbd3	Core component of NuRD corepressor complex Does not show strong binding to mCpG	Mouse	Early embryonic lethal
Mbd4	DNA repair protein that binds mCpG and T:G mismatches at mCpG sitesThymine DNA glycosylase that excises T from T:G mismatches	Mouse	Viable and fertile. three- to fourfold increase in mutations at CpG sites. Increased susceptibility to intestinal cancer correlates with C to T transitions within the <i>Apc</i> gene. Mbd4 functions to minimize the mutability of 5-methylcytosine.
Kaiso	Binds mCGmCG and CTGCNA	Mouse	No overt phenotype. Small but significant delay in tumorigenesis on Min background.


Methyl-CpG binding proteins are present in transcriptional co-repressor complexes

MeCP2: component of the Sin3A HDAC complex

Mbd3: component of the NuRD HDAC complex


Mbd1: interacts with HDAC3. Mbd1 and HDAC3 are recruited by the PML-RARalpha hybrid protein to silence gene expression in Acute promyelocytic leukemia

MBD1: interacts with the H3K9 HMTase SETDB1



Collaboration to repress genes

TRANSCRIPTIONAL REGULATION BY METHYL-DNA BINDING PROTEINS RECRUITMENT OF Methyl-CpG binding proteins and co-repressor complexes



MeCP2: components of the Sin3A HDAC complex

Mbd3: component of the NuRD HDAC complex

Mbd1: interacts with HDAC3.

Example: Mbd1 and HDAC3 are recruited by the PML-RARalpha hybrid protein (specialized transcription factor) to silence gene expression in cancer

MBD1: interacts with the H3K9me3 HMTase SETDB1

DNA methylation collaborates with other chromatin modifying complexes to repress gene expression





Figure 9. Recruitment of corepressors by methyl-CpG binding proteins. A hypothetical transition between an active, nonmethylated gene promoter and a repressed promoter whose silence is attributable to DNA methylation, as mediated by complexes containing an MBD protein such as MeCP2 (gray shading). The transition phase represents an intermediate step during which transcription is silenced and DNA methylation occurs. MeCP2 is envisaged to recruit the NCoR histone deacetylase (HDAC) complex and histone lysine methyltransferase (HKMT) activity to the methylated sites. I

DNA METHYLATION IS ABUNANT IN THE GENOME AND ISSUBJECTED TO DRAMATIC ALTERATIONS DURING EMBRYOGENESIS



DNA methylation levels are high in fertilized Oocytes that contain the paternal and maternal genome (carries characteristic methylation patterns)

Paternal and maternal methylation patterns are rapidly erased (exception: imprinted genes maintain paternal and maternal methylation information). → the paternal and maternal methylation epigenome is cancelled

DNA methylation levels remain low during the first cell division events until the blastocyst stage

In the blastocyst stage cell differentiation programs are activated and genes need to be regulated on the epigenetic level → DNA methylation is increasing (loss of DNMT1, DNMT3a or DNMT3b is lethal → establishment and maintenance of DNA methylation is impaired)

70%- 80% of CpG dinucleotides are methylated in the genome

70%- 80% of CpG di-nucleotides are methylated in the human genome!

Remember only 2% of the genome encode for mRNAs

98% is noncoding DNA that contains a large proportion of transposable elements, repeat sequences, etc...

DNA METHYLATION IS REVERSIBLE: DNA DEMETHYLATION BY Tet-family proteins





Figure 6. Model of Tet-initiated DNA demethylation pathways. DNA methylation (5mC) is established and maintained by DNMT. 5mC can be oxidized by Tet family of dioxygenases to generate 5hmC, 5fC, and 5caC. Because the oxidized 5mC derivatives cannot serve as substrates for DNMT1, they can be lost by replication-dependent passive demethylation. 5hmC can be deaminated by AID/APOBEC to become 5hmU, which together with 5fC and 5caC can be excised by glycosylases such as TDG, followed by DNA repair to generate C. Alternatively, a putative decarboxylase may convert 5caC to C.

Tet-family proteins mediate DNA demethylation



5mC, 5hmC and 5fC are abundant in the cell
5caC is present only at very low abundance

DNMT1 has exclusive specificity for 5mC

DNA METHYLATION IS REVERSIBLE: DNA DEMETHYLATION BY Tet-family proteins





Figure 6. Model of Tet-initiated DNA demethylation pathways. DNA methylation (5mC) is established and maintained by DNMT. 5mC can be oxidized by Tet family of dioxygenases to generate 5hmC, 5fC, and 5caC. Because the oxidized 5mC derivatives cannot serve as substrates for DNMT1, they can be lost by replication-dependent passive demethylation. 5hmC can be deaminated by AID/APOBEC to become 5hmU, which together with 5fC and 5caC can be excised by glycosylases such as TDG, followed by DNA repair to generate C. Alternatively, a putative decarboxylase may convert 5caC to C.




The protein encoded by this gene belongs to the TDG/mug DNA glycosylase family. Thymine-DNA glycosylase (TDG) removes thymine moieties from G/T mismatches by hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of DNA and the mispaired thymine. With lower activity, this enzyme also removes thymine from C/T and T/T mispairings. TDG can also remove uracil and 5-bromouracil from mispairings with guanine. Interestingly, TDG knockout mouse models showed no increase in mispairing frequency suggesting that other enzymes, like the functional homologue MBD4, may provide functional redundancy. This gene may have a pseudogene in the p arm of chromosome 12. Additionally, in 2011, the human thymine DNA glycosylase (hTDG) was reported to efficiently excise 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC), the key oxidation products of 5-methylcytosine in genomic DNA. Later on, the crystal structure of the hTDG catalytic domain in complex with duplex DNA containing 5caC was published, which supports the role of TDG in mammalian 5-methylcytosine demethylation.

Check textbooks: glycosilases cleave off bases from sugar → apyrimidic/apurinic site → BER pthway

DNA METHYLATION IS REVERSIBLE: ACTIVE AND PASSIVE DNA DEMETHYLATION

DNA de-methylation of the paternal and maternal genome has different kinetics



5hmC: 5 hydroxymethylcytosine

PASSIVE DNA DEMETHYLATION

Successive rounds of DNA methylation reduce the amount of 5mC. In this situation DNMT1 is excluded from the

Nucleus! (only transient presence of oocyte specific version of DNMT1 at the 8 cell stage)

MATERNAL GENOME: slow de-methylation of DNA

ACTIVE DNA DEMETHYLATION

Enzymatic activity rapidly de-methylates 5mC

PATERNAL GENOME: fast de-methylation of DNA

→ **In zygotes Tet3 is localized to the PATERNAL nucleus**

→ **Paternal DNA is demethylated**

→ **High levels of 5hmC: 5-hydroxymethylcytosine, 5fc: 5-formylcytosine and 5caC: 5-carboxylcytosine were detected at high levels in the paternal nucleus**

→ **BER machinery concentrated in pronucleus**

