

### CO<sub>2</sub> as a C<sub>1</sub> building block




### Carbon based products




CO<sub>2</sub> volumes required to cover the production of the main chemicals consumed in the EU (in MtCO<sub>2</sub>/yr)

### Carbon dioxide



$$\delta^{-}$$
  $2\delta^{+}$   $\delta^{-}$   $0$   $C$   $0$ 

- Non-polar
- Electrophilic at C (Lewis acid)
- Nucleophilic at O (Lewis base)



LUMO 2π<sub>u</sub>

## CO<sub>2</sub> Activation

#### STRENGTHEN POINTS

**WEAKNESS POINTS** 


Cheap;

Nontoxic;

Widely available (30 Gt/year (2014)).


High thermodynamic stability; Low reactivity.

#### CO<sub>2</sub> coordination modes to transition metals




## Hydrogenation of CO<sub>2</sub>

### REDOX POTENTIALs



#### **Involved reactions**



## Hydrogenation of CO<sub>2</sub> to

#### **FORMIC ACID**

It is endoergonic; it is entropically disfavored

$$\Delta H^0 = -31.2 \text{ kJ/mol } \Delta G^0 = +32.9 \text{ kJ/mol } T = 25 ^{\circ}C \text{ pH} = 0$$

With the addition of a base, like NH<sub>3</sub>

$$\Delta H^{0} = -84.3 \text{ kJ/mol } \Delta G^{0} = -9.5 \text{ kJ/mol } T = 25 ^{\circ}C$$

In polar solvents, like water


$$\Delta G^0 = -4.0 \text{ kJ/mol} \quad T = 25 \, ^{\circ}\text{C}$$

#### **METHANOL**

It is exoergonic; but kinetically difficult

$$\Delta G^0 = -17.3 \text{ kJ/mol} \quad T = 25 \text{ °C pH} = 0$$

# Methylation of amines with $CO_2$ and $H_2$



20 examples up to 94% yield

33 examples up to 99% yield

### Literature

- T. Cantat et al. Angew. Chem. Int. Ed. 2012, 51, 187.
- T. Cantat et al. Angew. Chem. Int. Ed. 2014, 53, 2543.
- T. Cantat et al. ACS Catal. 2017, 7, 2107.

### **Tutorial**

A. Azapagic et al. *J. of CO<sub>2</sub> Util.* 2015, 9, 82. CCS vs CCU review 3 studenti

M. Beller et al. *JACS* 2012, *134*, 20701. From CO<sub>2</sub> to HCOO articolo 2 studenti

W. Leitner et al. *Chem. Sci.* 2015, *6*, 693. From CO<sub>2</sub> to CH<sub>3</sub>OH articolo con DFT e meccanismo 3 studenti

M. Beller et al. *Angew. Chem. Int. Ed.* 2013, *53*, 12156.

From CO<sub>2</sub> to N-CH<sub>3</sub> articolo 2 studenti

O
W. Leitner et al. *Angew. Chem. Int. Ed.* 2013, *53*, 9554.

W. Leitner et al. *Angew. Chem. Int. Ed.* 2013, *53*, 9554. From CO<sub>2</sub> to N-CH<sub>3</sub> articolo 1 studente