



Università di Trieste
LAUREA MAGISTRALE IN GEOSCIENZE
Curriculum Geofisico
Curriculum Geologico Ambientale

Anno accademico 2018 – 2019

Geologia Marina

Parte III

Modulo 3.1 Continental Margin Depositional Processes: down-slope processes

Relatore
Angelo Camerlenghi con materiale di Renata G. Lucchi

rglucchi@ogs.trieste.it



Source-to-sink system

Siliciclastic continental shelves


Sedimentary processes on Continental Margins

Deep Marine Systems

Sedimentary processes on Continental Margins

Mass Transport Deposition >>> MTDs

Turbidity currents >>> Turbidites




Sedimentary Processes on Continental Margins

down-slope: driven by gravity forces

along-slope: driven by density forces

(thermo-haline or water mass accumulation)



the Source to Sink System



Continental shelf

Preferential area of sediment accumulation

High sediment accumulation

High isostatic subsidence




Continental slope sediment deposition and transfer toward deeper environments

Continental rise: sediment deposition (deep sea fans, sediment drifts)




Siliciclastic continental shelves

Wind/wave dominated shelves
e.g. Mediterranean margins (tides ≤ 3 m)



Other sedimentary/biological processes

- Storms sediment resuspension
- Surface and bottom turbidity currents associated to river output (Hypopycnal and Hyperpycnal flows on deltas and prodeltas)
- Incursions of surface ocean currents on the outer shelf
- sediment bioturbation



IL PICCOLO

EVENTI IN


[HOME](#) [CRONACA](#) [SPORT](#) [TEMPO LIBERO](#) [NORDEST ECONOMIA](#) [ITALIA MONDO](#) [FOTO](#) [VIDEO](#)SI PARLA DI [CASAPOUND](#) [BASKET](#) [MALTEMPO](#) [TRIESTINA](#) [BALCANI](#) [FERIERA](#) [REGENI](#)Sei in: [TRIESTE](#) > [CRONACA](#) > LO SCIROCCO TRASFORMA IL CORNO D'ORO...

DALMAZIA

Lo scirocco trasforma il Corno d'oro in un uncino sull'isola di Brazza

Il maltempo di questi giorni ha determinato un fenomeno alquanto curioso e insolito lungo le coste della Croazia


04 novembre 2018 |


4
Commenti647
8⁺

Deep-Marine Systems

Down-slope
processes

Along-slope
processes





Sedimentary processes on Continental Margins

Depositional process → **Deposit**

down-slope processes:
driven by gravity forces

- » Mass Transport Deposition → **MTDs**
- » Turbidity currents → **Turbidites**
- » Riverine outflows → **Hyper (Hypo)- picnites**
- » Turbid meltwaters → **Plumites**
- » Brine-related deposition

along-slope: driven by density forces (thermo-haline origin)

- » Contour currents → **Contourites**



Classification of MTDs (Mass Transport Deposits):

- Reology (sediment deformation)
- Sediment mass mechanism of support (gravity, flow turbulence, grains interaction)
- Physical properties of the mass flow and deposit (sediment disturbance, shear strength, etc.)
- Morphological characteristics of the deposit


References:

- Dott R. H., 1963. Dynamics of subaqueous gravity depositional processes. AAPG Bulletin, 47, 1, pp. 104-128.
- Lowe, D.R., 1982, Sediment gravity flows II. Depositional models with special reference to the deposits of high-density turbidity currents: Journal Sed. Petrology, 52, pp. 279-297.
- Prior, D.B. (1984). Submarine landslides. Proceedings of the IV International Symposium on Landslides, Toronto, Vol. 2, pp. 179-196.
- Norem, H., Locat, J. and Schieldrop, B. (1990). An approach to the physics and the modelling of submarine landslides. Mar. Geotech., 9, 93-111.
- Martinsen, O. (1994). Mass movements. in: The geological deformation of sediments, (A. Maltman Ed.), Chapman and Hall, London, pp. 127-165.
- Mulder, T. and Cochonat, P. (1996). Classification of offshore mass movements. J. Sediment. Res., 66, 43-57.
- Masson, D.G., Harbitz, C.B., Wynn, R.B., Pedersen, G., Lovholt, F. (2006). Submarine Landslides: processes, triggers and hazard prediction. Phil. Trans. R. Soc. A, 364, pp 2009-2039.



cohesive material
distinct boundaries
few displacements
pore fluid not important

- mass slide
 - no failure surface
low deformation rate
 - distinct failure surface
 - isolated blocks or aggregates of blocks
rapid movement



indistinct boundaries
pore fluid important
in triggering and motion

- gravity flow
 - laminar flow → mass flow
 - turbulent flow

creep

slide

debris avalanche

$h/l < 0.15$
translational slide

$h/l > 0.15$
rotational slide

matrix supported → debris flow

fluid supported → grain flow


turbidity current → low density

high density



Submarine slides/slumps

They are **ubiquitous** features of submarine slopes in all geological settings and at all water depths, particularly in areas where fine grained sediments predominate.



Rift & Transform Margins

- Moderate earthquakes
- **Submarine/subaerial landslides**
- Debris flows, turbidity currents
- Tsunamis
- Methane emissions

Subduction Margins

- Large earthquakes
- **Submarine/subaerial landslides**
- Explosive eruptions, lahs
- Debris flows, turbidity currents
- Tsunamis
- Methane emissions

Oceanic Volcanoes

- **Submarine/subaerial landslides**
- Debris flows, turbidity currents
- Volcanic eruptions
- Earthquakes
- Tsunamis
- Methane emissions


Passive Margins

- **Submarine landslides**
- Debris flows, turbidity currents
- Modest earthquakes
- Tsunamis
- Methane emissions

Complexity:


Once failure initiates, the event may **progress by means of a number of mass movement processes**. Although various subdivisions and classification schemes for these processes exist, each process represents part of a continuum, whereby one type may evolve into or trigger another.

Many submarine slope failures are likely to have involved a number of processes, possibly active at different stages of failure. Therefore, it is common that the depositional units resulting from submarine mass movements are defined as '**Mass-Transport Complexes (MTC)**'.







Submarine slides/slumps



Rotational landslide



Translational landslide



Number of Skempton
height of slide/length of slide

<0.15 SLIDE



>0.15 SLUMP




Small slump
In sediment core



Pleistocene Submarine Landslides in the Boso Peninsula, Japan




Multibeam



Sub-bottom

Ana submarine landslide Ibiza Channel Western Mediterranean




Lastras et al., 2004 Sedimentology



STOREGGA SUBMARINE LANDSLIDE, NORWAY


8000 y BP
3500 km³ of debris




Courtesy Petter Bryn



Norsk Hydro
E&D Norway
Geophysical Operations





STOREGGA SUBMARINE LANDSLIDE






Goleta landslide, California






Goleta landslide, California



Deep penetration seismics
2D Sparker

Lee et al., 2009. GSA Special Paper



Morgan et al., 2009. Scientific Drilling




Submarine debris avalanches


Volcanic Island Margins Hawaii

Moore et al., 1994. JGR




Volcanic Island Margins Canarie





Volcanic Island Margins Stromboli, Lipari Islands, Italy




Stromboli Sciara di Fuoco 100.000 y

Romagnoli et al., 2009. Marine Geology





cohesive material
distinct boundaries
few displacements
pore fluid not important

- mass slide
 - no failure surface
low deformation rate
 - distinct failure surface
 - isolated blocks or aggregates of blocks
rapid movement





indistinct boundaries
pore fluid important
in triggering and motion

- gravity flow
 - laminar flow → mass flow
 - turbulent flow



Gravity flows




Debris flows

Laminar flux supported by the water-rich muddy matrix


Debris flow: mud/sand >1; pebbles >5%

Mud flow: mud/sand <1; pebbles <5%


strongly coherent debris flows




A




weakly/moderately coherent debris flows



B



Gravity-Driven Downslope Processes in Deep Water









Turbidity flows

Density currents in which the granular support is maintained by the vertical component of the turbulent flux

A






Turbidity flows

Density currents in which the granular support is maintained by the vertical component of the turbulent flux

A



TYPE OF EVENT



Long steady flow (e.g. river fed)
Short surge-type (e.g. river floods,
slope instability)

FLOW DENSITY

High density (higher velocity) $>1.1 \text{ g/cm}^3$
Low density (lower velocity) $<1.1 \text{ g/cm}^3$

FLOW TRANSFER

Confined (canyon, channel, levee,
deep-sea fan)
Unconfined






upwards decrease of flow velocity/energy

	GRAIN SIZE	BOUMA (1962) DIVISION	INTERPRETATION
100 cm	Mud	E - Laminated to homogeneus mud	Deposition from low-density tail or turbiditi current settling of pelagic or hemipelagic particels
	Silt	D - Upper mud/silt laminae	Shear sorting of grains and flocs
	Sand	C - ripples, climbing ripples wavy or convolute laminae	Lower part of lower flow regime of Simons et al. (1965)
		B - plane laminae	Upper flow regime plane beds
100 cm	Coarse sand	A - structurless or graded sand to granule	Rapid deposition with no traction transport possible quick (liquefied) beds

(after Pickering et al., 1989)




sandy turbidite

silty turbidite

Turbidite facies

Coarse-Grained Turbidites


Lowe (1982) Divisions



← High-Density Turbidity Currents →

Classic Turbidites

Bouma (1962) Divisions



Fine-Grained Turbidites

Stow and Shanmugam (1980) Divisions

	(Hemi) Pelagite Bioturbation
	Ungraded Mud, Mirabiloburated
	Ungraded Mud, ±Silt, Pseudonodules
	Graded Mud, ±Silt Lenses
	Wispy, Convolute Lamination
	Indistinct Lamination
	Thin, Regular Lamination
	Thin, Irregular Lam., Low Amplitude Climbing Ripples
	Convolute Lamination
	Basal Lenticular Lamination

← Low-Density Turbidity Currents →




LOW DENSITY turbidity flows

Stow and Shanmugam (1980)

Bouma (1962)

Lowe (1982)



- Shanmugam, G., 2000. 50 years of the turbidite paradigm (1950s-1990s): deep-water processes and facies models – a critical perspective. *Marine and Petroleum Geology* 17, 285-342.
- Kevin Pickering, Richard Hiscott, 2014. Deep Marine Systems: Processes, Deposits, Environments, Tectonic and Sedimentation. Wiley-Blackwell, ISBN: 978-1-4051-2578-9, 776p.



UNIVERSITÀ DEGLI STUDI DI TRIESTE

Dipartimento di Matematica e Geoscienze




i Geologia Marina 2018-19



silty turbidites



sandy turbidite



muddy turbidites



silty turbidite




UNIVERSITÀ DEGLI STUDI DI TRIESTE

Dipartimento di Matematica e Geoscienze

Corso di Geologia Marina 2018-19





MOST COMMON FEATURES

- « Sharp base characterized by sharp grain size change often with sharp color change (careful with sediment oxidation)
- « Planar laminations
- « Bioturbated top

INDICATION OF SHEAR SORTING


Grain size and compositional sorting through the deposit. Sorting occurs according to size and specific weight (e.g. large forams with medium-size quartz with small-size pyroxene)

COMPOSITION


Presence of allocthonous particles
e.g. shelf derived particle in deep-sea environments (typically bryozoa, autogenic glauconite)



Size - Velocity Diagram



Bouma Sequence






SMALL-SCALE BED FORMS: $\lambda < 20 \text{ cm}$				
	Symmetric small ripples (SSR) regular, 2D, symmetrical, sharp crests, straight flanks, broad troughs	SSR + asymmetric small ripples (ASR) more irregular, 2-3D, still symmetrical rounded crests, some straight and some bimodal flanks	ASR + asymmetric large ripples irregular, 3D, asymmetrical, larger λ and height, round bimodal profiles, pronounced occur on lower end of slopes	Current ripples very irregular, 3D, sharp crests, steep and straight lee, convex-up slopes 5-10 (Yokokawa 1995)
Symmetry index	-1.2		-1.5	- angle of repose (30-35°)
Dip of lee side		11-18°	"24-27°" dip of lee side increases with increasing U_u	0.5-0.6 (Yokokawa 1995)
Roundness index	0.44	-0.50	> 0.50	12-22 (Harms 1989) 7-20 (Allen 1985a) 6-11, fca (Yokokawa 1995) -20, fca (Boggs 2001)
Ripple index	generally between 8-12 for all bed forms			N/A
Orbital diameter/ wavelength	8-15	-8-15	8-15	
LARGE-SCALE BED FORMS: $\lambda > 100 \text{ cm}$				
	Symmetric large ripples (SLR) SLR: 2D, symmetrical, sharp discontinuous crests = to brink, straight flanks	Hummocky (HM) + SLR + ALR HM: 3D, symmetrical, no brink point, broad round crests, domal, convex-up flanks	Asymmetric large ripples (ALR) ALR: 2D-3D, asymmetrical, brink not always = to crest, round stoss with break in slope, can have scour pits on lower end of stoss	Dunes regular (2D) to irregular (3D), sharp crests, steep and straight lee, straight to convex-up stoss
Symmetry index	-1.0 (-1.5)	-2	> 2	-
Dip of lee side	14-24° (SLR); 15-29° reverse large ripples (RLR)		"23-31°" dip of lee side increases with increasing U_u	- angle of repose (30-35°)
Roundness index	~ 0.40-0.50 highest for HM bed forms	-0.45-0.60	-0.55-0.75 (up to 0.95)	-
Ripple index	generally between 8-12 for all bed forms			12-22 (Harms 1989) 20-40 (Allen 1985a) - 6, fca (Boggs 2001)
Orbital diameter/ wavelength	1-2	1-2	1-2	N/A





antidune formation



bedforms



Lucchi, 1997. PhD Thesis, University of Cardiff



Turbidite facies


Fine-Grained Turbidites

Stow and Shanmugam (1980) Divisions

	(Hemi) Pelagite Bioturbation
	Ungraded Mud, Mirabiloburated
	Ungraded Mud, ± Silt, Pseudonodules
	Graded Mud, ± Silt Lenses
	Wispy, Convolute Lamination
	Indistinct Lamination
	Thin, Regular Lamination
	Thin, Irregular Lam., Low Amplitude Climbing Ripples
	Convolute Lamination
	Basal Lenticular Lamination


Coarse-Grained Turbidites

Lowe (1982) Divisions



Classic Turbidites

Bouma (1962) Divisions




← High-Density Turbidity Currents →




Low-Density Turbidity Currents



HIGH DENSITY turbidity flows




The *linked debrite*







Talling et al., 2007. Nature 450, 541-544.

Confined systems: Canyons and associated deep see fans




Reflected turbidites and Multi-sources turbidites



Contourites

or

Fine-grained turbidites



Stow and Shanmugam (1980) Divisions

	(Hemi) Pelagite Bioturbation
T_8	Ungraded Mud, Microbioturbated
T_7	Ungraded Mud, \pm Silt Pseudonodules
T_6	Graded Mud, \pm Silt Lenses
T_5	Wispy, Convolute Lamination
T_4	Indistinct Lamination
T_3	Thin, Regular Lamination
T_2	Thin, Irregular Lam. Low Amplitude Climbing Ripples
T_1	Convolute Lamination
T_0	Basal Lenticular Lamination