



Università di Trieste
Corso di Laurea in Geologia

Anno accademico 2018 - 2019

Geologia Marina

Parte IV

**Modulo 4.2 Indicatori di movimento di fluidi: Vulcani di Fango,
chimneys, pockmarks, vents...**

Docente

A. Camerlenghi

Outline

Review of main mechanisms of fluid flow:

- **Mud diapirs and mud volcanoes**
- Gas chimneys
- Pockmarks
- Seafloor vents in general
- Polygonal fault systems
- Diagenetic fronts
- Gas hydrates

Mud volcanoes

Surface expressions of focused fluid flow inside hydrocarbon-bearing sedimentary basins. They can:

- indicate subsurface petroleum accumulations
- may react to or reveal precursor signals of earthquakes
- induce hazards for people and industrial facilities
- release large amounts of methane into the atmosphere.

Definition of Mud Volcano

stacks of debris flow deposits composed of fluid-rich, fine-grained sediments expelled on the Earth's surface or on the sea floor. During the ascent, the mud is able to carry litho-clasts of various size, shape, age, and composition.

Mud volcanoes are often associated to sedimentary diatremes and mud diapirs (shale diapirs, or clay diapirs), all generated by subsurface overpressure of sedimentary (high accumulation rate), tectonic, or diagenetic origin following a state of under-consolidation in low-permeability sediments.

Although mud volcanoes occur in both divergent and convergent margins, they play an important role in the evolution of accretionary wedges, where they too participate in the world wide controversy about the origin and significance of mélanges.



Olistostromes, or sedimentary mèlanges: uplifted and at times deformed **chaotic** sedimentary bodies (Cretaceous to Pliocene) originated by subaqueous mass gravitational processes, such as debris-flows, and submarine slides and/or mud volcanoes/diapirs.

Tectonosomes, or broken formations: strongly deformed up to stratally disrupted Ligurian units, which retain their original stratigraphic coherence. They represent fossil, uplifted portions of the offscraping complexes of the Cretaceous-Eocene paleo-Apennine accretionary wedge.


- ARGILLE SCAGLIOSE -
 - OLISTOSTROMES - Sedimentary melanges
 - TECTONOSOMES - Tectonic melanges
- MASS WASTING DEPOSITS
- MUD VOLCANOES
- MUD DIAPIRS

**mud volcanoes
overlying a diapir**


diatreme




Kopf (2002)



Hovland et al. (1998)



Seep structures and deposits on dormant mud volcanoes. A Salse A at the crater field of the Dashgil mud volcano, with the gryphon field to the west (B). C Hydrocarbons (black mud) in a gryphon at Bakhar. D Burning hydrocarbon gas in the vent at Lokbatan. The fire has been burning for more than a year since the October 2001 eruption (Figs. 2 and 3)





NATURAL FIRES OF AZERBEIJAN

Marco Polo(?)

Images courtesy of Luis Piñero




“The appearance of Zoroastrans in Azerbaijan and their cult of the eternal flame in the Temple of Fire of the Magi might be related to the fires from the mud volcanoes”




Planke et al. (2003)

Lokbatan Mud Volcano, Azerbaijan,
25 October 2001





Eruption Piparo, 22/2/1987



Piparo, Trinidad 22 February 1987





CHECKING IT OUT: BEHIND: down, Opposition leader Patrick Manning, a trained pilot, and Prime Pandey, closely examine the road which was covered from the Piparo mud volcano last Saturday, with his

Sunday Guardian

• 25519 • ESTABLISHED SEPTEMBER 5, 1917 • SUNDAY, FEBRUARY 23, 1997 • CANTONMENT CORNER (9887 THE TRINIDAD PUBLICATION CO. LTD.) • \$3

Teekwondo champion Chery-Anni Sankar in flying high — PAGE 19

SUNSHINE'S Danella Callender cuddles up with teddy bear

Courtney Walsh is Windies captain against Indians — PAGE 6

Jeanine Bonterra is a shining star in ZONE

Volcano erupts



Mountain of mud leaves 31 Piparo families homeless

By LOUIS B HOMER

A MUD volcano erupted in Piparo at 5:30 am, yesterday leaving 31 families homeless and 15 injured and 20 are listed under a 100-foot high mountain of mud. Villagers, who live on the edge of the newly formed crater, were yesterday, basically remaining household items as well as farm animals and even parts of houses, buried in another eruption.

The officials who visited the area have issued a warning for people to stay away from the village since the volcano is still erupting.

A normally quiet area, Lightfoot Town, the scene of disaster, was a hub of activity with hundreds of tourists people coming to see the volcano and listen to the villagers tell of the horror of the eruption.

Alan Khan, one of the Piparo villagers who lost his house to the volcano, was lucky to be awake when the eruption started.

"It was about midnight after five a 'clock when I heard the electricity wires crackling, then it started to spurt. Afterwards I heard the whole place crackling and the earth started to move, then I heard a rattle and there an explosion."

"When I looked outside I saw a set of mud going up in the air for about 100 feet high. This lasted for about 15 minutes, then it quieted down and started again and lasted for over one hour". He took his family to stay with relatives who live further away from the volcano in Piparo.

Another villager, Hulsan Sankar, believed a storm was coming on. Going outside to check, he saw mud spouting into the sky like fountains of smoke from a chimney.

"The mud was about 60 feet in diameter. I had to run 100 feet because the mud from the volcano was

The



Trinidad Guardian

• 25520 • ESTABLISHED SEPTEMBER 5, 1917 • MONDAY, FEBRUARY 24, 1997 • CANTONMENT CORNER (9887 THE TRINIDAD PUBLICATION CO. LTD.) • \$1.25

Volcano takes first life

Piparo resident dies after helping villagers

BY LOUIS B HOMER

TOBY SPAIN has been the life in the aftermath of the mud volcano which has killed one of the 30 people in Piparo, South Trinidad, on Saturday.

Formerly one of the most famous and respected men in the area, Toby Spain, who lived about 100 feet from the mud volcano, died yesterday after he was found by mud which he had dug up for his wife's funeral.

Spain, a beloved by his family, made sure he had been buried in the mud volcano, which was the only way to get to his home.

Spain was found by mud which he had dug up for his wife's funeral.

Spain, a beloved by his family, made sure he had been buried in the mud volcano, which was the only way to get to his home.

Spain was found by mud which he had dug up for his wife's funeral.



Mud mass

Piparo volcano erupts Page 3



BELOW: A HOUSE is left in ruins in the aftermath of the eruption of a mud volcano in Piparo early yesterday morning. This house was pushed over by the wall of mud on the left.

RIGHT: MEMBERS of a Piparo family salvage what is left of their belongings from this second house which was pushed over and partially covered by mud following the early morning eruption.

Photo: STEPHEN KJZ



TESTING THE MUD: Stepping lightly, Prime Minister Basdeo Pandey examines the mud while visiting the site of the mud volcano which almost wiped out the Piparo Village in Central Trinidad last Saturday morning. Photo by TONY HOWELL.

Piparo villagers to be relocated




By LOUIS B HOMER

Mosque, Prime Minister Basdeo Pandey, who travelled by helicopter to the area, said the volcano eruption could be seen as another opportunity "to unite our people as one."

25 miles of roadway. Those listed for immediate attention are Piparo, Guaracara Tabaquite Rd, Sisters Rd, Cho Road, Hoseeene Road, and Nar Mayaro Road.





THE ENVIRONMENTAL DISASTER OF THE MUD VOLCANOE TRIGGERED BY DRILLING FOR OIL IN JAVA: ISOLA DI GIAVA, MAY 29 2006





MUD VOLCANOES **SALSE DI NIRANO, ITALY**



How to recognize submarine mud volcanoes

1. Core samples showing ‘mud breccia’ containing sediments with a range of different ages, compositions and structures.
2. Strong backscatter on side-scan sonar records representing topographic features (craters, cones, mud flows, etc.).
3. Evidence of gas seepage and associated features (bacterial mats, cold-seep communities or methane derived authigenic carbonate – MDAC).
5. Seismic evidence of feeder channels and/or mud diapirs.

Classification of mud breccia from Mediterranean Sea mud volcanoes according to sedimentary facies

Lithotype or sedimentary facies	Description
A - MASSIVE	Matrix-supported clasts of soft to indurated marls. No size sorting observed in clasts and matrix.
MASSIVE A1	centimetric to pluri-centimetric clasts. Stiff matrix.
MASSIVE A2	millimetric clasts. Stiff matrix.
MASSIVE A3	mousse-like texture of the matrix produced by gas micro-vesicles
B - ORGANIZED	The mud breccia shows internal textural changes. The breccia can be either matrix- or clast-supported.
ORGANIZED B1	sub-horizontal (in sediment cores) bedding produced by thin layers of millimetric clasts sorted by size. No embricate structures observed.
ORGANIZED B2	upward graded grain-supported mud breccia. The matrix/clasts ratio increases upwards.
ORGANIZED B3	matrix supported mud breccia with patches (clouds) of different colors and composition.


(adapted from Camerlenghi et al., 1992 and Staffini et al., 1993).



Dimitrov, 2002



Clasts




slumps




Mud-breccia oxidized

Photos by Renata G. Lucchi







Mousse facies





Organized facies

Fig. 4 Seafloor and sediment images from DMV (4A-C): **A** recent mud flow sheets from a seafloor fissure; **B** small vent sites from an area of seepage on DMV; **C** white bacterial mat in a seafloor crack on DMV; **D** fractured gas hydrate slabs in sediments from Odessa mudflow core M52/1-18





FOSSIL MUD VOLCANO, OLIMPIC PENINSULA



Degree of Overpressure


$$\lambda = (P_f - P_{hy}) / (P_d - P_{hy})$$

P_f = Pore pressure

P_{hy} = Hydrostatic Pressure

P_d = Total Stress

SEDIMENT COMPACTION AND INFLATION GRAIN PRESSURE AND OVERPRESSURE






$$\lambda = 0 \text{ if } P_f = P_{hy}$$

$$\lambda = 1 \text{ if } P_f = P_d = \text{fluid movement (liquid mud)}$$




Mud diapirs move when $0 < \lambda < 1$

MUD VOLCANOES IN THE WORLD










© Ifremer 2003





Mud volcanoes in the Gulf of Mexico







Geological setting of mud volcanoes









Shale tectonics, Offshore Brunei







classical interpretation based on 2D seismic






new interpretation based on 3D seismic




Mud volcanoes offshore Nigeria






Mechanisms of emplacement



Mechanisms of emplacement



Diapir initiation in buoyant shales with potential deep fluids migration along structural highs (e.g. anticline axes) or fault networks

Fluids migration from different units and overpressure increase, diapiric structure development and brecciation during its growth

Overpressured diapir reaches critical depth. Overburden cannot contain fluids rich diapir. System in unstable conditions ready for triggering

Blast of gas. The sudden pressure release allows large amount of fluidized and gas saturated sediments to reach the surface

THE DISCOVERY OF SUBMARINE MUD VOLCANOES IN THE MEDITERRANEAN SEA

- **1981** Mud volcanoes were first reported in the Eastern Mediterranean by M.B. Cita, W.B. Ryan and L. Paggi.




The Prometheus dome was identified according to:


Morphology: wrinkled surface of small concentric ridges;

Acoustic character: no penetration, no coherent reflections

Lithologic composition : **MUD BRECCIA**, structureless pebbly mud with dominantly angular semi-indurated clasts of various, non carbonatic composition. The matrix contains foraminiferal species dating to the Aptian-Cenomanian.

It was interpreted as a SHALE DIAPIR, and a comparison between the chaotic sedimentary facies of the Prometheus dome and the Argille Scagliose was immediately presented to the public.






References:

- Bohrmann, G., Ivanov, M., Foucher, J.-P., Spiess, V., Bialas, J., Greinert, J., Weinrebe, W., Abegg, F., Aloisi, G., Artemov, Y., Blinova V., Drews, M., Heidersdorf, F., Krabbenhöft, A., Klaucke, I., Krastel, S., Leder, T., Polikarpov, I., Saburova, M., Schmale, O., Seifert, R., Volkonskaya, A. and Zillmer, M. (2003) Mud volcanoes and gas hydrates in the Black Sea: new data from Dvurechenskii and Odessa mud volcanoes. *Geo-Mar Lett.*, 23, 39–249
- Camerlenghi, A., and Pini, G.A., in press. Mud volcanoes, olistostromes, and argille scagliose in the Mediterranean Region. In: J.A. McKenzie, D. Bernoulli, M.B. Cita (Eds.), *Major Discoveries in Sedimentary Geology in the Mediterranean Realm from a Historical Perspective to New Developments*. IAS Special Publications, Blackwell Publishing.
- Costa, E., Camerlenghi, A., Polonia, A., Cooper, C. Fabretti, P., Mosconi, A., Murelli, P., Romanelli, M., Sormani, L. and Wardell, N. (2004) Modeling deformation and salt tectonics in the Eastern Mediterranean Ridge accretionary wedge. *Geol. Soc. Am. Bull.*, 116, 880-894.
- Davies, R.J. and Stewart, S.A. (2005) Emplacement of giant mud volcanoes in the South Caspian basin: 3D seismic reflection imaging of their root zone. *J. Geol. Soc. (London)*, 162, 1-4.
- Dimitrov, L.I. (2003) Mud volcanoes—a significant source of atmospheric methane. *Geo-Mar. Lett.*, 23, 155-161.
- Davies, R.J., Swarbrick, R.E., Evans, R.J. and Huuse M., 2006. Birth of a mud volcano: East Java, 29 May 2006. *GSA-Today*, v. 17, no. 2, doi: 10.1130/GSAT01702A.1
- Kopf, A. (2002) Significance of mud volcanism. *Rev. Geophys.*, 40, 1-51.
- Diaz del Rio, V., Somoza, L., Martinez Frias, J., Mata, M.P., Delgado, A., Hernandez Molina, F.J., Lunar, R., Martin Rubi, J.A., Maestro, A., Fernandez Puga, M.C., Leon, R., Llave, E. and Medialdea, T. (2003) Vast fields of hydrocarbon-derived carbonate chimneys related to the accretionary wedge/olistostrome of the Gulf of Cadiz. *Mar. Geol.*, 195, 177-200.
- Hovland, M., Hill, A. and Stokes, D. (1997) The structure and geomorphology of the Dashgil mud volcano, Azerbaijan. *Geomorphology* 21, 1-15.
- Krastel, S., Spiess, V., Ivanov, M., Weinrebe W., Bohrmann G., Shashkin P. and Heidersdorf, F. (2003) Acoustic investigations of mud volcanoes in the Sorokin Trough, Black Sea. *Geo-Mar. Lett.*, 23, 230–238.
- Loncke, L., Mascle, J. and Fanil Scientific Party (2004) Mud volcanoes, gas chimneys, pockmarks and mounds in the Nile deep-sea fan (Eastern Mediterranean): geophysical evidences. *Mar. Petr. Geol.*, 21, 669–689.
- Planke, S., Svensen, H., Hovland, M., Banks, D. A. and Jamtveit, B. (2003) Mud and fluid migration in active mud volcanoes in Azerbaijan. *Geo-Mar. Lett.*, 23, 258–268.
- Robertson, A.H.F. and Kopf, A. (1998) Tectonic setting and processes of mud volcanism on the Mediterranean Ridge accretionary complex: evidence from Leg 160. In: *Proc. ODP Sci. Results*, 160 (Eds. A.H.F. Robertson, K.-C. Emeis, C. Richter and A. Camerlenghi), pp. 665-680. Ocean Drilling Program, Texas A&M University, College Station, TX.
- Somoza, L., Diaz del Rio, V .D., Leon, R., Ivanov, M., Fernandez Puga, M.C, Gardner, J.M., Hernandez Molina, F.J., Pinheiro, L.M., Rodero J., Lobato A., Maestro, A., Vazquez, J.T., Medialdea, T. and Fernandez Salas, L.M. (2003) Seabed morphology and hydrocarbon seepage in the Gulf of Cadiz mud volcano area: Acoustic imagery, multibeam and ultra-high resolution seismic data. *Mar. Geol.*, 195, 153-176.
- Sumner, R.H. and Westbrook, G.K. (2001) Mud diapirism in front of the Barbados accretionary wedge: the influence of fracture zones and North America-South America plate motions. *Mar. Petr. Geol.*, 18, 591-613.
- Westbrook, G.K. and Reston, T.J. (2002), The accretionary complex of the Mediterranean Ridge: tectonics, fluid flow and the formation of brine lakes - an introduction to the special issue of *Marine Geology*. *Mar. Geol.* 186, 1-8.
- Woodside, J.M., Mascle, J., Zitter, T.A.C., Limonov, A.F., Erguün, M. and Volkonskaia, A. and shipboard scientists of the PRISMED II Expedition (2002) The Florence Rise, the Western Bend of the Cyprus Arc. *Mar. Geol.*, 185, 177-194.


Outline

Review of main mechanisms of fluid flow:

- Mud diapirs and mud volcanoes
- **Gas chimneys**
- **Pockmarks**
- **Seafloor vents in general**
- **Polygonal fault systems**
- **Diagenetic fronts**
- Gas hydrates





G. Bohrmann, Initial results from M52/1 **MARGASCH**



Judd and Hovland, 2007. *Seabed Fluid Flow*.

Parametric echo
sounder image of a
'flare' (intense water
column target caused
by vigorous gas
seepage) rising 850 m
from the seabed in
the NW BlackSea.





Methane concentration in the
seawater from methane sensors
on CTD casts

Gas chimneys are vertical zones in some way or other have been 'disturbed' by previous or on-going gas migration.


Exactly what has caused this acoustically-detected disturbance is still unknown, although it is believed that small (metre-sized) parcels of trapped gas and slightly displaced sediments may be involved. In many cases, rather than a distinct chimney, gas may be present as an amorphous cloud.


Gas accumulations provoke a high acoustic impedance contrast
>>>> High reflectivity.



- **Enhanced seismic reflections**
- **Bright spots**
- **Flat spots**
- **Acoustic blanking**
- **Columnar disturbances, gas chimneys, pipes**
- **BSR (in case of gas hydrates)**


Seismic section across the Tommeliten Delta structure, a salt piercement diapir. The noisy zone is interpreted as a gas chimney through which gas rises vertically (as indicated by the large arrow). Some gas escapes laterally to produce brightening of adjacent reflectors, and reducing the acoustic velocity (v_p) to produce 'pull down' (examples are in the rectangles).



Judd and Hovland, 2007. *Seabed Fluid Flow.*











Pockmarks are shallow seabed depressions, typically several tens of metres across and a few metres deep.

Generally, they are formed in soft, fine-grained seabed sediments by the escape of fluids (gas or water, but mainly methane) into the water column.

Judd and Hovland, 2007. *Seabed Fluid Flow.*




Figure 2.6: Assymmetric pockmarks, Witch Ground Basin, UK North Sea. Multi-beam echo sounder image [Image acquired by the UK government (Department of Trade and Industry) as part of the Strategic Environmental Assessment process.]

Judd and Hovland, 2007. *Seabed Fluid Flow.*




Figure 2.3: Typical North Sea pockmarks, Witch Ground Basin, UK North Sea. Multi-beam echo sounder image [Image acquired by the UK government (Department of Trade and Industry) as part of the Strategic Environmental Assessment process.]

Judd and Hovland, 2007. *Seabed Fluid Flow*.




Figure 2.40: MBES image of the *Scanner* pockmark, Block UK15/25, North Sea. [Image acquired by the UK government (Department of Trade and Industry) as part of the Strategic Environmental Assessment process.]

Judd and Hovland, 2007. *Seabed Fluid Flow*.




Figure 2.8: MBES image of pockmark strings in the Norwegian Sea. These strings have no preferred orientation, and some lead to (or from) large standard pockmarks. The 26 inch Haltenpipe pipeline is visible on the lower part of the image.

Judd and Hovland, 2007. *Seabed Fluid Flow*.




Figure 2.16: Pockmarks in the northern part of the South Fladen Pockmark Study Area; MBES survey, 2001. The pipeline (Scott-Forties Unity pipeline; 24 inch, 61 cm, diameter) gives an idea of the scale. [Image acquired by the UK government (Department of Trade and Industry) as part of the Strategic Environmental Assessment process.]

Judd and Hovland, 2007. *Seabed Fluid Flow.*




Figure 2.19: Boomer profile across the Witch's Hole, an unusual pockmark in the South Fladen area. [Reproduced by permission of the British Geological Survey. © NERC. All rights reserved. IPR/67-34C.]

Judd and Hovland, 2007. *Seabed Fluid Flow.*







Figure 2.44: Seep plumes from the Scanner (left), Scotia (centre), and Challenger (right) pockmarks, Block UK15/25, North Sea acquired during the Heincke 180 cruise, October 2002 (Alfred Wegener Institute) using the parametric sediment echo sounder system (SES-2000DS) developed at Rostock University, Germany; this scan shows depths from 140 to 190 m. [courtesy of Gerdt Wendt, University of Rostock.]

Judd and Hovland, 2007. *Seabed Fluid Flow*.



Judd and Hovland, 2007. *Seabed Fluid Flow*.





Bathymodiolus heckeriae mussel beds. (A) Juvenile and adult mussels at Marker 'E'. (B) Dead mussels and octopus. (C) Extensive bed of live mussels of relatively uniform size, partially covered by bacterial mats, at Marker 'B'. (D) Dead mussels at the eastward periphery of Marker 'B'. (E) Mussels with a chiridotid holothurian and Alvinocaris sp. (F) Mussels with Alvinocaris sp. And ophiuroids. Scale bars: A–D : 10 cm, E; F : 5 cm.

Van Dover et al. (2003). *Deep Sea Research*

Chemosynthetic organisms at cold seeps


Free-living filamentous sulfur bacteria: *Beggiatoa*



tube worms: *Lamellibrachia*



clams: *Calyptogena*




mussels: *Bathimodiolus*




mussels: *Acharax / Solemya*



Gradient of chemosynthetic communities





Authigenic Carbonates:

Isotopically light

Organic markers indicate presence of methane oxidizers



Serve as habitat for bottom fish (e.g. rock fish)





Jensen *et al.*, 1992

Fossil carbonate chimneys (PobitiKamani, Bulgaria) Judd &, Hovland 2006




Courtesy, Domenico Rio, University of Parma

FOSSIL DEWATERING CHIMNEYS IN PLIOCENE MARLS, CROTONE BASIN

Pipe-like chimneys in Gulf of Cadiz Mud Volcanoes

Composed of authigenic carbonates with iron oxides.



Carbonates are moderately depleted in ^{13}C , ranging from - 46‰ to -20 ‰ PDB



Diaz del Rio al., Marine Geology 195 (2003) 177-200

Judd and Hovland, 2007. Seabed Fluid Flow.






Coupled Sulfate / Barium Profiles

1. High concentrations of barite in pelagic sediments underlying high productivity waters are thought to result from **biologically mediated precipitation of barium sulfate within the water column.**
2. In organic-rich, rapidly accumulating sediment, sulfate is consumed by microbial reduction of organic carbon. **Barite is dissolved under conditions of sulfate depletion leading to high barium concentrations in the pore fluids.**
3. **When barium-rich fluids discharge at the seafloor, barite forms by reaction with seawater sulfate,** forming “cold-seep barite” deposits.

***Paleozoic Bedded barites* from Nevada, Arkansas, Mexico and South China are associated with:**

- Organic shales
- Chert and phosphorite
- Some carbonate



***Cold seep barites* along continental margins are associated with:**

- Organic-rich facies
- Opal-rich sediments
- Phosphorites
- Some carbonates



Jewell and Stallard, J. Geol., 99, 1991

ODP Leg 112


Polygonal Faults




'an array of layer-bound extensional faults within a mainly fine-grained stratigraphic interval that exhibit a diverse range of fault strikes which partially or fully intersect to form a polygonal pattern in map view'


- This type of fault has only been recognized in packages that are predominately composed of fine-grained sediments.
- The local stress regime operative at the time when polygonal faults grow can exert a significant influence on strike and on the organisation of the fault array as a whole.

Polygonal Faults. Geometry


- Polygonal faults systems develop in tiers. Faults in one tier may partially interconnect with those in adjacent tiers by cross-propagation of a sub-set of the total fault population, but the majority of the faults in the separate tiers are contained wholly within individual tiers.
- They range in fault trace length from 100 m to several kilometres and extend vertically across discrete layers from a few tens of metres to over one kilometre in thickness.
- Polygonal faults can be planar or listric
- Faults are characterized by a large range of fault strikes. Where strikes are almost randomly oriented, a classical polygonal plan form geometry results.
Variations in the basic polygonal plan form can arise from regional slope, tectonic context, or basement topography, or from intrinsic variation in the physical properties or the thickness of the deforming interval.
- Three-dimensional geometry of polygonal fault systems is invariably complex and difficult to appreciate from simple 2D cross-sections.









0 ————— 3 km



2km





Polygonal Faults. Genetic mechanisms

Gravity sliding-collapse NO

Sliding down a slope, with a basal detachment at the boundary would imply iso-orientation, and basal contraction.


Density Inversion (Such as Henriët)

Density inversion should produce folding (like in salt diapirs), which are not always observed.

Syneresis


- Syneresis is a spontaneous contraction (shrinkage) without evaporation, but is a process that is specifically restricted to gels.
- Gels are a framework of colloidal particles, and the primary condition for gel formation is the very fine size range of the constituent particles (clay size range).
- Ultra-fine grained sediments in which all polygonal fault systems form, fall into the range of colloidal materials

Density inversion model For polygonal fault systems Henriet et al., 1989.



Deposition of clays
Downward flow into aquifer

Formation of top and bottom seals by preferential compaction adjacent to sands



Density inversion –
Pressure build-up –
Rayleigh-Taylor instability

Overpressure relaxation
hydrofracturing
Roof faulting of top seal
Collapse
Undercompacted claystone becomes compacted

References:

- Cartwright, J.A., 1994a. Episodic basin-wide hydrofracturing of overpressured Early Cenozoic mudrock sequences in the North Sea Basin. *Marine and Petroleum Geology*, 11, 587-607.
- Cartwright, J.A. 1994b. Episodic collapse of geopressured shale sequences in the North Sea Basin. *Geology*, 22, 447-450.
- Cartwright, J.A. 1996. Polygonal Fault Systems: a new type of geological structure revealed by 3D seismic data. In: WEIMER, P. (ed.) *Application of 3-D seismic Data to Exploration and Production, American Association of Petroleum Geologists, Studies in Geology*, 42, 225-231.
- Cartwright, J.A. 1997. Polygonal Fault Systems in thick shale sequences. In: Yardley, B, & Jamtveit, B. (Eds) *Fluid Flow Processes in the Continental Crust*. Chapman & Hall, London, 81-104.
- Cartwright, J.A. and DeWhurst, D. 1998. Layer-bound compaction faults in fine-grained Sediments. *Bulletin of the Geological Society of America*, 110, 1242-1257.
- Cartwright, J.A. and Lonergan, L. 1996. Volumetric contraction during the compaction of mudrocks: a mechanism for the development of regional-scale polygonal fault systems. *Basin Research*, 8, 183-193.
- Cartwright, J. and Lonergan, L. 1997. Polygonal fault systems in the Eromanga and North Sea Basins: a comparison. *Exploration Geophysics*, 28, 323-331.
- Davies, R. Cartwright, J.A. & Rana, J. 1999. Polygonal density inversion structures from the Faerøe-Shetland Trough. *Geology*, 27, 798-802.
- DeWhurst, D., Cartwright, J.A. & Lonergan, L. 1999a. The development of polygonal fault systems by the syneresis of fine-grained sediments. *Marine and Petroleum Geology*, 16, 793-810.
- DeWhurst, D., Cartwright, J.A. & Lonergan, L. 1999b. Three-dimensional consolidation of clay-rich sediments. *Canadian Geotechnical Journal*, 36, 355-362.
- Gouly, N.R. 2001. Polygonal fault networks in fine grained sediments - an alternative to the syneresis mechanism. *First Break*, 19, 69-73.
- Henriet, J.P., Debatist, M., Van vaerenbergh, W. & Verschuren, M. 1989. Seismic facies and clay tectonic features in the southern North Sea. *Bulletin of the Belgian Geological Society*, 97, 457-472.



FOSSILIFEROUS SOILS




- They trap water and introduce a significant bias on index properties, and diatoms microfossils in particular can play a significant role on physico-chemical properties of soil because of their potentially large specific surface area.
- They can provide delayed compressibility or a sudden increase in compressibility once the yield strength of the microfossil is exceeded.
- They influence the frictional behavior of soils by their size and shape.

Calcareous




Siliceous







Distribution of diatoms in a consolidated sediment taken at a depth of 383 m at the site of Kansai airport.


Intra-skeletal (Is, a) and skeletal (Sq, b) porosity of microfossils.




Rebesco et al. (1998),
Terra Antarctica, 5(4),
715-725









SITE 1095



SITE 1096







— 0,1 mm



— 10 μm




— 40 μm




— 50 μm

A



— 50 μm


B




— 10 μm



C


Site 1095



Site 1096







Conclusions:

- Due to their shape and strengths, microfossils affect significantly the physical properties and mechanical behavior of marine sediments:
 - Microfossil-rich sediments retain porosity with depth.
- They resist consolidation until a threshold value of applied stress, exceeded which the rigid structure of the sediment collapses.
- Micro-structural collapse may trigger overpressure in the pore fluids, and weakness of the sediment (decreased effective stress).
 - Diagenesis may act in two ways:
 - _ Cementation contributed to strengthening of the sediment.
 - _ Opal A to C/T transformation acts as a micro-structural collapse
- Microfossil rich sediments, and oozes in particular, are candidate sediments to provide weakness surfaces in submarine slopes.

References:

- Davies, r. J., and Cartwright j., 2002. A fossilized Opal A to Opal C/T transformation on the northeast Atlantic margin: support for a significantly elevated Palaeogeothermal gradient during the Neogene? *Basin Research*, 14, 467-486.
- Handwerger, D.A., Cooper, A.K., O'Brien, P.E., Williams, T., Barr, S.R., Dunbar, R.B., Leventer, A., and Jarrard, R.D., 2004. Synthetic seismograms linking ODP sites to seismic profiles, continental rise and shelf of Prydz Bay, Antarctica. In Cooper, A.K., O'Brien, P.E., and Richter, C. (Eds.), *Proc. ODP, Sci. Results*, 188 [Online]. Available from World Wide Web: http://www-odp.tamu.edu/publications/188_SR/010/010.htm .
- Lodolo, E., and Camerlenghi, A., (2000). The occurrence of BSRs on the Antarctic margin. In: *Natural Gas Hydrate in Oceanic and Permafrost Environments*, (Ed. by M.D. Max), pp.199-213. Kluwer Academic Publ., Dordrecht.
- Locat, J., and Tanaka, H., 2001. A new class of soils: fossilifereous soils ? Une nouvelle classe de sols: les sols fossilifères ? In: *Proceedings of the 15th International Conference on Soil Mechanics and Geotechnical Engineering*, Istanbul, 27-31 August 2001, Vol. 3, pp.: 2295-2300.
- Marine Geology*, 189:343-370.
- O'Brien, P.E., Cooper, A.K., Richter, C., et al., 2001. *Proc. ODP, Init. Repts.*, 188 [Online]. Available from World Wide Web: http://www-odp.tamu.edu/publications/188_IR/188ir.htm.
- Rebesco, M., Larter, R.D., Camerlenghi, A., and Barker, P.F., 1996. Giant sediment drifts on the continental rise west of the Antarctic Peninsula. *Geo-Marine Letters*, 16, 65-75.
- Rebesco, M., Camerlenghi, A., and Zanolla, C., 1998. Bathymetry and morphogenesis of the continental margin West of the Antarctic Peninsula. *Terra Antarctica*, 5(4), 715-725.
- Rebesco M., Pudsey C., Canals M., Camerlenghi A., Barker P., Estrada F., and Giorgetti A., 2002. Sediment Drift and Deep-Sea Channel Systems, Antarctic Peninsula Pacific Margin. In Stow D. A. V., Pudsey C. J., Howe J.A, Faugeres J. C. and Viana A.R., (Eds.), *Deep-Water Contourite Systems: Modern Drifts and Ancient Series, Seismic and Sedimentary Characteristics*. Geological Society, London Memoirs, 22, 353-371.
- Tanaka, H., and Locat, J., 1999. A microstructural investigation of Osaka Bay clay: the impact of microfossils on its mechanical behavior. *Canadian Geotechnical Journal*, 36, 493-508.
- Tribble, J.S., Mackinkie, F.T., Urmos, J., O'brien, D.K., & Manghnani, M.H., 1992. Effects of biogenic silica on acoustic and physical properties of clay-rich marine sediments. *Am. Ass. Petr. Geol. Bull.*, 76(6), 792-804.
- Volpi, V., Camerlenghi, A., Moerz, T., Corubolo, P., Rebesco, M., & Tinivella, U., 2001. Data report: Physical properties relevant to seismic stratigraphic studies, continental rise Sites 1095, 1096, and 1101, ODP Leg 178, Antarctic Peninsula. In: *Proc. ODP, Sci. Results*, Vol. 178 (Ed. by P.F. Barker, A. Camerlenghi, G.D. Acton & A.T.S. Ramsay). Available from World Wide Web: http://www-odp.tamu.edu/publications/178_SR/chap_17/chap_17.htm
- Volpi, V., Camerlenghi, A., Hillenbrand, A.-D., Rebesco, M., & Ivaldi, R., 2003. Effects of biogenic silica on sediment compaction and slope stability on the Pacific Margin of the Antarctic Peninsula. *Basin Research*, 15, 339–363.