


LECTURE 5

COORDINATION OF HISTONE AND DNA METHYLATION


Linking de-novo DNA methylation to histone methylation



De-novo DNMT family has 2 enzymatic active members (DNMT3a, b) and one regulatory factor DNMT3L


PWWP (Pro-Trp-Trp-Pro) domain: protein or DNA interaction domain

ADD (ATRX-DNMT3-DNMT3L) domain: highly similar between DNMT proteins: **CAN INTERACT WITH HISTONE TAILS**



*ADD domain of DNMT3a
Interacts with the **unmethylated** Histone H3
tail → a link of de-novo DNMTs
with histone methylation*

Linking de-novo DNA methylation to histone methylation




- DNMT3L forms a complex with DNMT3a → tetramer: 2x DNMT3L; 2x DNMT3a (best studied); DNMT3L also interacts with DNMT3b
- Phenotype of DNMT3L Knock-out = phenotype of DNMT3a = DNMT3a and DNMT3L are functionally linked
- Deletion of interaction domains that link DNMT3a to DNMT3L results in enzymatic inactivation = DNMT3a function depends on tetramer formation and DNMT3L!!
- Histone H3 tails interact with ADD domains (only DNMT3L shown); red circles: interaction with DNA


Linking de-novo DNA methylation to histone methylation

How can we find out whether there is a functional link between histone modifications and DNA methylation???

Linking de-novo DNA methylation to histone methylation



DNMT3L links histone methylation to DNA methylation




Note:

- Olig1/2 have a bivalent status in mouse embryonic stem (ES) cells → bivalent (H3K27me3/H3K4me3) → not expressed

- Olig1/2 are monovalent active: no H3K27me3 but H3K4me4 → expression in NPCs

DNMT3L links histone methylation to DNA methylation



H3K4me0: DNA METHYLATION IN CpG ISLANDS

H3K4me1,2,3: NO DNA METHYLATION IN CpG ISLANDS




De novo DNA methyl-
transferases
translate patterns of
H3K4methylation
into heritable patterns of
gene expression

HOW???

DNMT3L links histone H3K4 methylation to DNA methylation

B



DNMT3L ADD domain binds with high affinity to un-methylated Histone H3 tails

DNMT3L in tetramer binds unmethylated histone H3 → CpG methylation by DNMT3a/DNMT3b


Mutated DNMT3L does not bind to unmethylated H3K4 → no DNA methylation at CpG islands!!

De novo DNA methyl-transferases
translate patterns of H3K4methylation
into heritable patterns of gene expression

H3K4 HKMTs have an important role in defining CpG methylation levels

H3K4 methylation and CpG island methylation

H3K4 HKMTs and CpG methylation




MLL1 and SET1 HKMTs are most relevant


H3K4 specific HKMTs are important for the activation of gene expression

- MLL proteins are required to activate Hox gene during differentiation
- MLL proteins are often involved in translocations in myeloid and lymphoid leukemias (→ MLL hybrid gene results HKMTase activation at inappropriate genes)

Is there a link between H3K4me and DNA methylation to coordinate gene expression



The CXXC domain binds un-methylated CpG islands




MLL1 binds to UNMETHYLATED CpGs



MLL HKMTs mediate H3K4methylation and prevent CpG methylation

MLL1 is essential to activate the expression of Hox genes; Hox genes are essential for embryoid developments

This Study: MLL1 knock-out mice → use primary mouse embryonic fibroblasts to study Hox gene expression



H3K4me ChIP at the *Hox c8* locus Analysis by PCR




MLL1 is essential to
Impose H3K4 methylation

(control: H3K9me: MLL1 does
not change H3K9me)

Note: ChIP does not discriminate
Between H3K4me1, H3K4me1,
H3K4me3

MLL1 is central for H3K4 methylation at Hox genes

H3K4 specific MLL HKMTs prevent CpG methylation



+: wild-type; -: MLL1 null

DNA methylation sensitive restriction enzymes:

B: BstUI:

Ha: HaeIII

Hh: HhaI

Hp: Aval

Cut when CpG unmethylated; do not cut when CpG methylated





Prepare genomic DNA from MLL1 wt and MLL1 knock-out cells


Digest with DNA methylation sensitive restriction enzymes

Make southern blot and hybridize with probes that recognize the *Hox c8* locus

General result: in MLL1^{-/-} cells there is 1 band more (red bxx)





SET1 AND CFP1 LINK H3K4 methylation and DNA methylation



The CXXC domain binds
un-methylated CpG islands

SET1 H3K4 HKMT binds CFP1
and is recruited by
CFP1 to un-methylated CpG islands



SET1 H3K4 HKMT

Figure: SET1 in vicinity to histone H3 tail.

Note: CFP1 is an interacting partner of SET1. CFP1 has CXXC domain that binds unmethylated CpG

Scenario: unmethylated CpG;
CFP1 binds and recruits SET1 →
H3K4me increases

CFP1 is enriched at peaks of H3K4me3 that overlap with CpG islands




Figure 2. Genome-wide ChIP sequencing shows a tight association between Cfp1 and H3K4me3 at CGIs


a, Typical Cfp1 ChIP-Seq profiles from whole mouse brain. For comparison, we also carried out H3K4me3 ChIP-Seq. The data were aligned with non-methylated CGIs mapped in mouse brain using a CXXC affinity column²⁹. The panel shows a typical region of the genome from chromosome 4 (nucleotides 126,333,759–127,054,849) demonstrating the coincidence of Cfp1 and H3K4me3 peaks with CGIs. A subset of genes is labelled (RefSeq). Two CGIs that lack H3K4me3 and Cfp1 coincide with sites of H3K27me3 binding (red rectangles; data of ref. 30 for mouse brain). **b**, Venn diagram showing strong overlap

ChIP seq on brain cells:
CpG islands that show high H3K4me3 but are unmethylated (see earlier slides) and are enriched for CFP1 (interacts with SET1 H3K4 HKMT)

CFP1 CXXC domain is required to bind to unmethylated CpG islands

CFP1 is enriched at peaks of H3K4me3 that overlap with CpG islands

LOSS OF CFP1 RESULTS IN A REDUCED H3K4me3 AT unmethylated CpG ISLANDS → LOSS OF SET1 RECRUITMENT!!!!
= CFP1 is essential to recruit SET1 to CpG islands





ChIP qPCR using Cfp1, H3K4me3 and H3K27me3 antibodies at selected loci in vector-only control and Cfp1-depleted NIH3T3 cells. The results were replicated with an independent clone expressing the same shRNA combination (data not shown) and with each of two individual shRNA constructs (see Supplementary Fig. 3).

CFP1 is enriched at peaks of H3K4me3 that overlap with CpG islands

Experimental model system:

- Embryonic stem cells are stably transfected with a DNA fragment that contains puromycin and EGFP: both sequences are enriched in extremely CG rich (but are protein coding)
- The fragment does NOT contain a promoter
- A) bisulfite sequencing: the inserted CpG rich DNA sequence is NOT METHYLATED
- B) ChIP seq using CFP1, H3K4me3 and RNA PolII



CFP1 and H3K4me3 is enriched at un-methylated CpG islands


BUT: RNA Pol II is not recruited; why? → fragment does not contain promoter.

RESULT: un-methylated CpG are sufficient to recruit CFP1 + SET1


To increase H3K4me3, also in the absence of transcription

THAT MEANS THAT THE UNMETHYLATED CpG SEQUENCE IS SUFFICIENT TO DIRECT H3K4me3

H3K36 de-methylation and CpG island methylation



CXXC domains mediate binding to unmethylated CpGs: H3K36 specific KDM2A/JHDM1A (K-Histone De-methylase)




JHDM1A/KDM2A is a histone de-methylase that ensures low H3K36me2/me1 levels at CpG islands

CXXC domain binds un-methylated CpG islands

Tet1 has CXXC zinc finger domain. However, the CXXC domain of TET1 has no DNA binding activity and is dispensable for its catalytic activity in vivo. Other interacting proteins recruit Tet1 to DNA


PRIMARY OBSERVATION THAT JHDM1A IS LINKED TO DNA METHYLATION



WT ES cells: **chromocenters are DNA methylated (a large block of methylated DNA)**: Flag-tagged JHDM1a is randomly distributed in the nucleus



DNMT1^{-/-} ES cells: No DNA methylation at chromocenters = a large block of DNA are without DNA methylation. In this case KDM2A/JHDM1a localizes to chromocenters = **is attracted by unmethylated CpGs**

CXXC domains mediate binding to unmethylated CpGs: H3K36 specific KDM2A/JHDM1A (K-Histone De-methylase)




ChIP seq on ES cells: JHDM1a/KDM2A
Concentrate on un-methylated CpG islands

Interaction is dependent on the CXXC domain



H3K36me2 is high in gene body but low at promoter
High levels of KDMA2 at unmethylated CpG islands is paralleled by low **H3K36me2** levels


H3K4 de-methylation and CpG island methylation



The H3K4 de-methylase LSD1 (KDM1A) is essential for establishing DNA methylation


LSD1 is a H3K4 specific demethylase: oxidizes H3K4me_{2,1} → H3K4me₀

LSD1 konditional knock-out mice die early in embryogenesis (E5.5) and show strongly reduced DNA methylation



Loss of LSD1 results in
Reduced DNA methylation

Recruitment of LSD1 eliminates H3K4me_{1,2} resulting in H3K4me₀
This creates a binding site for DNMT3L → thus recruiting the
DNMT3L-DNMT3a tetramer





Southern blotting using CH3 sensitive restriction enzymes: a probe for the H19 and IAP imprinted gene locus are used. These are classic loci are controlled by DNA methylation

Note: Loss of DNA methylation results in Efficient restriction digest (more small fragment). This means that DNA methylation is strongly reduced Situation is similar to DNMT1 knock-out cells

DNA methylation sensitive restriction enzyme –
does not cut when CpG is methylated


H3K9 methylation and CpG island methylation

The relation of H3K9me and DNA methylation



The role of the G9a/GLP heterodimer in controlling DNA methylation

G9a HMTase and GLP HMTase form dimer and methylate H3K9m1; H3K9me2




Immunoprecipitation:

- Cells transiently transfected with Flag-tagged G9a and Dnmt3a
- IP anti-flagG9a: DNMT3a interacts
- IP anti-flagG9a: DNMT3b interacts

The role of the G9a/GLP heterodimer in controlling DNA methylation

H3K9 methylation



The role of the G9a/GLP heterodimer in controlling DNA methylation

Embryonic stem cells:

Self-renewing mESCs
(pluripotent)

Oct4, Stk10, Gpr54
Nanog, Dnmt3L, Tnfrsf8
ON
NO DNA METHYLATION


retinoic
acid

Differentiated mESCs

Oct4, Stk10, Gpr54
Nanog, Dnmt3L, Tnfrsf8
OFF
DNA METHYLATION



G9a – knock-out cells in differentiation

Show a loss of DNA methylation
As detected by bisulfite sequencing
In CpG islands of several genes



G9a: Required for silencing of transposable elements, repeat elements, retroviral insertions, imprinting centers but also in **gene expression control**

The role of the G9a/GLP heterodimer in controlling DNA methylation



Loss of Suv39h1 HMTases is linked to reduced DNA methylation at pericentric repeats

Southern blot using genomic DNA that was digested with methylation sensitive restriction enzyme. DNA on blot was hybridized using probes for minor satellite, major satellite and C-type retroviral DNA

Histone methylation and DNA methylation

