# Condensed Matter Physics I

(M. Peressi)

Exercises - Drude Model

Hint: normally the Ashcroft-Mermin book uses CGS units; however, it can be convenient to use MKS(SI) units in presence of eqs. containing the resistivity (use  $\rho$  in  $\Omega$ -meters, m in Kg, n in electrons/ $m^3$ , e in Coulomb.

#### Exercise 1

1. Consider typical values of the magnetic field (H(Earth) $\approx$ 0.5 Gauss; H(lab) $\approx$  1-10 T). Evaluate the product  $\omega_c \tau$ , where  $\omega_c$  is the cyclotron frequency and  $\tau$  the electron relaxation time. On average, can an electron in a metal make many revolutions between collisions or not?

## Exercise 2

1. Give a numerical estimate of the mean electronic velocity in case of a current density of 0.1 A/mm<sup>2</sup> flowing in a copper wire ( $n_{el} = 8.47 \ 10^{22} \ \text{cm}^{-3}$ ). [ $v=7 \ 10^{-4} \ \text{cm/s}$ ]

# Exercise 3

Consider Al at room temperature. Its electron density is  $n=18.1 \cdot 10^{22}/cm^3$  and its electrical resistivity is  $\rho=2.45~\mu\Omega\cdot cm$ .

- 1. Find its electron relaxation time  $\tau$  and electron mean free path  $\ell$  in the Drude model. [  $\tau$ =8.01  $10^{-15}$  s;  $\ell$  = 9 Å ]
- 2. Consider AC conductivity. At which frequency w the real part of the conductivity  $\sigma(\omega)$  will be 1/10 of its zero-frequency value? [  $\nu = \omega/2\pi = 59.7 \ 10^{12} \ \mathrm{Hz}$  ]

### Exercise 4

Sodium (Na) in standard temperature and pressure conditions is a metal with BCC structure, density of about  $0.97 \text{ g cm}^{-3}$  and mass number = 23.

- 1. Calculate the atomic density (number of atoms per unit volume) of solid sodium. [  $n_{at}$ =2.54 10<sup>22</sup> cm<sup>-3</sup> ]
- 2. Calculate the electron density. [  $n_{el}=n_{at}$  ]
- 3. Calculate the plasma frequency. [  $\omega_p$ =8.99 10<sup>15</sup> s<sup>-1</sup> ]
- 4. If the electron gas is treated as a classical one, which is the average kinetic energy of an electron at  $T=0^{\circ}$  K? and at room temperature? [0; 3.88  $10^{-2}$  eV]
- 5. Given the resistivity in DC at room temperature,  $\rho$ = 4.2  $\mu\Omega$  cm, calculate the relaxation time  $\tau$ . [  $\tau$ =3.3  $10^{-14}$  s ]