

Tight binding bands - DOS - cyclotron orbits

Exercise 1: *Tight binding bands in the square lattice*


Consider a square lattice with one s orbital per site, only nearest-neighbor interactions and neglect overlap.

1. Show that the expression of the tight binding s -band is:

$$E(\mathbf{k}) = E_{1s} - 2\gamma (\cos k_x a + \cos k_y a)$$

where $\gamma = -\int \phi_{1s}(\mathbf{r}) \Delta U(\mathbf{r}) \phi_{1s}(\mathbf{r} - \mathbf{R})$; \mathbf{R} are Bravais lattice vectors joining the nearest-neighbour sites to the origin, and $\gamma > 0$.

2. Show that the band has remarkable symmetry properties: the band $E(\mathbf{k})$ is symmetric with respect to E_{1s} , in the sense that every state at $E - E_{1s}$ has a corresponding state at $-(E - E_{1s})$, and, more precisely: $E(\mathbf{k}) - E_{1s} = -\left[E\left(\mathbf{k} - \frac{\pi}{a}(1, 1)\right) - E_{1s}\right]$
3. Show that the Fermi energy coincides with E_{1s} in case of half-filled band.
4. Plot the Brillouin zone and the Fermi surface in case of half-filled band.
5. Plot $E(\mathbf{k})$ along Γ - M - X - Γ , where $M = \frac{\pi}{a}(1, 0)$ and $X = \frac{\pi}{a}(1, 1)$.
6. Show that: Γ corresponds to the minimum, X to the maximum, M to a saddle point.
7. Plot the density of states $g(E)$ and show that it diverges at E_F (set to 0 in the figure below) with a logarithmic divergence. The figure also shows the occupied states in case of half filling of the band.



Exercise 2: Tight binding bands of “cubium”


Similarly as above, consider now the 3D case. Consider a simple cubic lattice with one s orbital per site, only nearest-neighbor interactions and neglect overlap (a toy model known also as *cubium*).

1. Show that the expression of the tight binding s -band is:

$$E(\mathbf{k}) = E_{1s} - 2\gamma (\cos k_x a + \cos k_y a + \cos k_z a)$$

where $\gamma = -\int \phi_{1s}(\mathbf{r}) \Delta U(\mathbf{r}) \phi_{1s}(\mathbf{r} - \mathbf{R})$; \mathbf{R} are Bravais lattice vectors joining the nearest-neighbour sites to the origin, and $\gamma > 0$.

2. Plot $E(\mathbf{k})$ along Γ - X - M - Γ - R , where $X = \frac{\pi}{a}(1, 0, 0)$, $M = \frac{\pi}{a}(1, 1, 0)$ and $R = \frac{\pi}{a}(1, 1, 1)$.
3. Plot the density of states $g(E)$.



Exercise 3: Semiclassical theory of electron dynamics: cyclotron orbits

Consider the problem of a cyclotron orbit in the (k_x, k_y) plane (suppose an external uniform magnetic field \mathbf{H} applied in the z direction, and $k_z(t = 0) = k_z(t = \infty) = 0$) for a SC crystal with band structure:

$$\mathcal{E}(\mathbf{k}) = \mathcal{E}_0 - 2A[\cos(ak_x) + \cos(ak_y) + \cos(ak_z)]$$

1. Show that the equation for the orbit with a given energy \mathcal{E} close to the minimum \mathcal{E}_{min} is: $\mathcal{E} = \mathcal{E}_{min} + Aa^2(k_x^2 + k_y^2)$.
2. Calculate the period of the orbit as a function of the parameter A .
3. Describe the orbit in real space.
4. Generalize your result to a crystal with band structure:

$$\mathcal{E}(\mathbf{k}) = \mathcal{E}_0 - 2[A \cos(ak_x) + B \cos(ak_y) + C \cos(ak_z)]$$