



Università di Trieste
LAUREA MAGISTRALE IN GEOSCIENZE
Curriculum Geofisico
Curriculum Geologico Ambientale

Anno accademico 2019 – 2020

Geologia Marina

Parte II


Modulo 2.3 **Metodi diretti: Sondaggi superficiali ed
analisi dei sedimenti**

Relatore

Dr. Renata G. Lucchi


rglucchi@ogs.trieste.it

BOTTOM SAMPLING SYSTEMS




Gravity corer

- classic gravity corer (Emery and Dietz, 1941; Hvorslev and Stetson, 1946)
- box corer
- kastenlot corer
- multi-corer



Piston corer

- classic piston corer (Kullenberg, 1947; 1955)
- long piston corer
 - Calypso piston corer (e.g. R/V Marion Dufresne, G.O. Sars)
 - Jumbo piston corer (e.g. R/V Araon)






Drilling systems


- ocean floor drilling systems (e.g. IODP-drilling vessels & semi-automated MeBo system)
- ice drilling systems (e.g. EPICA-European Project for Ice Coring in Antarctica & NorthGRIP- North Greenland Ice Core Project)

Classic gravity core system (Emery and Dietz, 1941)

It is the simplest coring device in which the weight of the coring equipment is used to force the barrel into the sea bottom. This system can work with or without a triggering system (sistema di sgancio)




Classic gravity core system: corer head



weight 600-800 kg
6000 kg

Classic gravity core system: core catcher and cutter



core catcher
(cestello)





core cutter
(scarpa)



core cutter
and catcher

Additional gravity core systems: **Kastenlot corer**


The kastenlot (kastens) corer was originally designed by Kögler (1963) it was improved and modified by Zangger and McCave (1990). The barrel, of variable lengths, is square in section (15x15 cm) and it contains a base plate that can be raised to reveal a new cleaned core surface



Square section
of barrel



Core cutter and catcher




Head of
kastens
corer



The core catcher has a shutter-like closure consisting of two square doors held under tension and blocked in a retracted position by two levers located on the outside. During the corer pullout, the pressure of the surrounding sediments pushes down the two levers closing the doors.

- (a) removal of the barrel lid to reveal the core surface;
- (b) two PVC trunkings are pushed into the sediments
- (c) the inner base is moved upward to expose the trunkings that are cut at the base and removed from the main core using a cheese wire;
- (d) sampling with x-ray slabs, and a further set of trunkings (e). Each time the sediments are lifted upward and withdrawn with a cheese wire

Sampling of Kastens cores



Additional gravity core systems: **BOX-corer**

Designed for minimum disturbance of the sediment surface, ideal for coarse/stiff sea floor sediments, it allows the recovery of bottom waters.



lateral surface



coral sampling

sampling of
glacigenic
sediments





Box core sampling



Additional gravity core systems: **Multi-corer**



Especially designed for the sampling of sea bottom sediments-water interface, it permits to recover low disturbance sediment. Ideal for geochemical and biological sediment and water analysis.



Eurofleets
BURSTER
PS99-1
RV POLARSTERN
LTER HAUSGARTEN
Bremerhaven – Longyearbyen
13.06.2016 – 23.06.2016

Kullenberg piston corer system


Standard assemblage for piston coring. The core barrel penetration is maximised by the action of a piston located in the lower barrel (or into the lower plastic liner if present) that helps to overcome the friction between sediments and the coring tube by generating Vacuum behind the cutter. The sediment cores obtained are less compacted and distorted than gravity cores. This system is always used coupled with a trigger mechanism.






Long piston corers:
Claypso, Jumbo etc.


Specifications	Kullenberg piston corer	Long piston corer
headweight	600 kg	6000 kg
barrel length	6 m	13 m
barrel inner diameter	65 mm	140 mm
barrel thickness	5 mm	5 mm
plastic liner outer diameter	63 mm	113 mm
plastic liner thickness	3 mm	5 mm
maximum cable length	5000 m	10000 m
cable diameter	12 mm	30 mm
freefall	4-5 m	1.5 m

Grab (benna)



Dredge (draga)





CORE ON
DECK!



Video Core logging lab


CUT INTO SECTIONS



SECTIONS' LABELING



The plastic liner is extracted from the barrel and cut into sections 1-1.5 m-long






The sections are numbered consecutively from bottom to top of the core or *vice versa* depending on the total length of the core.

Short cores are numbered consecutively from top to bottom.

Long cores are numbered consecutively on removal from the barrel from bottom to top of core.

Each section is labeled with a code indicating the name of the project and/or the name of the research vessel; the core number; the section number, and stratigraphic orientation (top-bottom)



CORE OPENING AND SEDIMENTS ANALYSIS

The plastic liner of each section is cut longitudinally. The plastic liner is cut by means of an electric saw/microvibro saw, while the sediments are cut using a “cheese wire”. The two splitted half-sections are labeled as **working section** and **archive section** and will undergone a different analytical process:


ARCHIVE SECTIONS: not destructive analyses

- X-radiographs
- multi-sensor core logger
- XRF core-scan
- photographs

WORKING SECTIONS: visual logging and sub-sampling



Half-sections' labeling




CORE OPENING AND SEDIMENTS ANALYSIS

The plastic liner of each section is cut longitudinally. The plastic liner is cut by means of an electric saw/microvibro saw, while the sediments are cut using a “cheese wire”. The two splitted half-sections are labeled as **working section** and **archive section** and will undergone a different analytical process:


ARCHIVE SECTIONS: not destructive analyses

- X-radiographs
- multi-sensor core logger
- XRF core-scan
- photographs

WORKING SECTIONS: visual logging and sub-sampling



Half-sections' labeling



CORE DESCRIPTION



Visual core description FORM



Project: _____

Observer(s) Date

samples		SEDIM. STRUCT.	LITHOLOGY (cm from top of section)	texture clay silt fine sand medium coarse grains- pebbles	LITHOLOGIC DESCRIPTION	COLOUR
forams	nanos diatoms s. slides					
			0			
			10			
			20			
			30			
			40			
			50			
			60			
			70			
			80			
			90			
			100			




length of section (cm)

total length of core (cm)

remarks:

LEGEND

— sharp boundary	🐚 pteropod rich
..... gradational boundary	🐚 forams rich
..... irregular boundary	🐚 shell's fragments
▲ normal and reverse grading	🐚 soupy sediment
▬ planar laminations	🐚 bioturbation
▬ cross laminations	
▲ slump	
V V V tephra	
— dark layer	



Visual core description 2nd step sediment composition

Mud composition - Smear slides

(view <https://www.youtube.com/watch?v=2sDejrpwxD4&feature=youtu.be>)

- A smear slide is a thin layer of unconsolidated sediment embedded on a glass slide for petrographic microscopic examination;
- Smear slides are a useful tool to quickly assess the **compositional content** of clay-silty sediment samples;
- Smear slides are a powerful method for rapidly evaluating tiny quantities of sediment (mineralogy, components, form, size) as the basis for **sediment classification**, and for ascertaining the presence of microfossils.






Foto: IODP Exp. 307




Quarz




IODP Exp .320


Plagioclase



Clay fraction





Composition/Quantification




Rothwell R.G., 1988. Minerals and mineraloids in marine sediments. An Optical Identification Guide. Elsevier Science Publishers, 279 pp

Diatoms





オホーツク海・中心部, 水深 1107m, XP98, PC1, Sec.1, 20cm
200倍, スケールは横 100 μ m

Foraminifers



太平洋・熊野トラフ, 水深 2190m, KT02-1, KK2PC
400倍, スケールは横 50 μ m

Calcareous nannoplankton




東地中海・キプロス沖, ODP Leg160, 967D, 1H, Sec.1, 0-5cm
1000倍, スケールは横 20 μ m

Sand composition

- wet sievings at 63 microns are used to investigate the composition of sediment coarse fraction, the grains form and roundness, in order to define the sediment provenance. The sand fraction is also used for biostratigraphic purposes.

Ice Rafted Debris
(Antarctica)



Beach sands
(Menorca-Spain)



Glacigenic
sediments
(Arctic)



Tephra
(volcanic glass)
(Tyrrhenian)





SEDIMENT SAMPLE DESCRIPTION

Project:

lat. N long. E water depth (m)
core section cm cm from core top

sediment type lithology
colour remarks

DESCRIPTION OF RESIDUE

textural characteristics

sorting	very good good moderately poor very poor	sphericity	high low	angularity	very angular angular sub-angular/rounded rounded well rounded
remarks				

residue composition

(AA=very abundant; A=abundant; O=occurring; R=rare; RR=very rare)

terrigenous components: predominant not predominant

quartz calcite K-feldspar gypsum glauconite
mica pyrite dark minerals volcanic glass micronodules
rock fragments others

bioclastic components: predominant not predominant

benthic forams	planktonic forams	pteropods	bryozoa
echinoid spines	sponge spicules	ostracods	corals
bivalves	gastropods	corals	algae
radiolarians	diatoms	spores	fish teeth
plant debris	coccoliths	discoasters	organic matter
micrascidites of tunicates	others	

Foraminifera: list of the most characteristic taxa

<i>G. bulloides</i>	<i>G. tenellus</i>	<i>H. siphonifera</i>	<i>G. coriacoenensis</i>
<i>G. trilobus</i>	<i>N. eggeri duttertrei</i>	<i>G. quinqueloba</i>	<i>G. inflata</i>
<i>N. pachyderma</i>	<i>G. ruber</i>	<i>G. scitula</i>	<i>O. universa</i>
<i>G. gomitulus</i>	<i>G. sacculifer</i>	<i>G. glutinata</i>	<i>G. truncatulin. exc.</i>
<i>G. conglobatus</i>	<i>G. quadrilobatus</i>	<i>H. pelagica</i>	<i>G. digit./preadigit.</i>
<i>S. ionica</i>	others	

AGE

ZONE

FACIES

REMARKS

SMEAR SLIDES DESCRIPTION

Project:

lat. N long. E water depth (m)
core section cm cm from core top

sediment type lithology
colour remarks

DESCRIPTION OF SMEAR SLIDES

Calcareous nannofossil: list of the most characteristic taxa and their abundance
(AA=very abundant; A=abundant; O=occurring; R=rare; RR=very rare)

<i>E. huxleyi</i>	<i>G. oceanica</i>	<i>G. caribbeanica</i>	<i>small Gephyrocapsa</i>
<i>H. carteri</i>	<i>C. leptopus</i>	<i>S. pulchra</i>	<i>S. histrica</i>
<i>C. pelagicus</i>	<i>S. recurvata</i>	<i>P. scutellum</i>	<i>O. antillarum</i>
<i>R. clavigera</i>	<i>R. stylifer</i>	<i>S. fossilis</i>	<i>B. bigelowi</i>
<i>U. tenuis</i>	<i>C. jonesii</i>	<i>T. saxea</i>	<i>C. rugosus</i>
<i>C. cristatus</i>	<i>P. multipora</i>	<i>P. lacunosa</i>	<i>C. macintyreii</i>
<i>D. broweri</i>	<i>H. sellii</i>	<i>Discoaster sp.</i>	

others.....

Sediment composition

(AA=very abundant; A=abundant; O=occurring; R=rare; RR=very rare)

terrigenous component: predominant not predominant

quartz calcite K-feldspar gypsum glauconite
mica dark minerals pyrite volcanic glass micronodules
dolomite aragonite Fe oxides zeolite plagioclase
rock fragments clay minerals.....
others

bioclastic component: predominant not predominant

benthic forams	planktonic forams	pteropods	bryozoa
echinoid spines	sponge spicules	ostracods	corals
bivalves	gastropods	corals	algae
radiolarians	diatoms	spores	fish teeth
plant debris	coccoliths	discoasters	fish remain
shell fragments	silicoflagellates	organic matter
others		







AGE

ZONE


FACIES

REMARKS

SEDIMENT CORE ANALYSES




X-RAY




SEDIMENT
SURFACE

SVAIS CORE-02 vert.



SVAIS CORE-02



CORING DISTURBANCE

