


**University of Trieste: GLOBAL CHANGE ECOLOGY a.a.
2019-2020**

BIODIVERSITY AND ECOSYSTEM FUNCTIONING
Dr. Stanislao Bevilacqua (sbevilacqua@units.it)

**The role of disturbance in
marine community dynamics**



Definition(s)

Disturbance is...

Any discrete event able to determine killing / removal from the substratum of one or more individuals, with the consequence of providing direct or indirect opportunities to new individuals for settlement or development **Sousa 1984**



Before the event



After the event

Disturbance

(e.g., storm)

It refers to the damage itself, that is, the effect (impact) of some external agent or force.

Sousa 2001

Definition(s)

Disturbance is...

Any discrete event able to change the structure of ecosystems, communities, or populations, limiting resources, modifying the substrate or the environment.

Pickett & White 1985



Before the event



After the event


(e.g., storm)

Disturbance

Disturbance is seen as a physical external force able to modify the system, for example removing organisms and opening patches. It refers to the physical agent that determine the biological consequences.

Perturbations

More generally, a perturbation is any interference with processes and structure characterizing a given system, or any event that change the state beyond its natural variation.



We intend **disturbance** as any event, which is caused or originates from a physical, chemical or biological **agent**, able to produce directly or indirectly changes (**impact**) to the system or its components.

The nature of disturbance

Physical

Physical disturbance refers to physical (or chemical) agents. For instance, hydrodynamic forces from intense wave action.



Biological

Biological disturbance is caused by organisms, though finally ascribable to physical actions. For instance, the whiplash of large algae.




Others?

In a wider sense, even predation could be considered as a disturbance, since it is able to remove large number of individuals and opens free space available for other organisms. However, it is internal to the system and we should exclude it from disturbance. The same for diseases, invasions, etc.

Types of disturbance

Sediments



Abrasion,
burial
Injuries,
suffocation,
death

Volcanic activity



Burning,
burial
Killing,
death



Storm wave and currents

Substrate
modifications,
physical action
Killing,
displacement

Temperature extremes

Salinity extremes

Anoxia



Oxygen
depletion,
osmotic and
metabolic
stress
Killing,
death



Ice scouring

Abrasion
Killing,
breaking,
death

Landslides



Abrasion,
burial
Killing,
breaking

Types of disturbance

Bioturbation



Whiplash




Debris accumulation



Burial,
displacement
Injuries,
suffocation,
death

Abrasion,
injuries
Killing, death

Shading, burial
Suffocation



Characteristics of disturbance


Intensity:

the strength of disturbance



Frequency:

the reoccurrence of disturbance



Spatial variability:

Variations in the extent of areas affected and distribution of disturbance

Ecological traits of organisms are important for the impact of disturbance and recovery potential


Regularity of disturbance – adaptation

Catastrophes




Expected effects under different scenarios

LOW-NO
DISTURB.



REG.
DISTURB.




REP.
DISTURB.



Temporal variability


LOW
FREQUENCY



Variations in timing of
disturbance occurrence

$$F_L = \frac{4}{12}$$


HIGH
FREQUENCY




$$F_H = \frac{6}{12}$$

$$I_L < I_H$$

LOW VARIANCE



HIGH VARIANCE




$$I_L = I_H$$


$$V_L < V_H$$

Effects of temporal variance...


LOW VARIANCE



MID VARIANCE



HIGH VARIANCE




■ = DISTURBANCE (6 / 24 m)


T_1 - T_7 = Sampling dates (!)



...and intensity



Results



Changes in temporal variability

- a) increasing intensity lead to increasing temporal variability in assemblage structure
- b) Increasing variance in disturbance lead to decreasing temporal variability

Recovery dynamics are affected differently by intensity and variance

Effects of temporal variance and intensity


Manipulative
transplanting experiment



Drilling




T
R
A
N
S
P
L






Experimental design




Timing of manipulation



Results

+ Aerial exposure
- Temp. variance
(+ variance)




+ barnacles
(drastically decrease)




Filamentous algae
C. branched algae

- Aerial exposure
- Temp. variance
(+ variance)



Reduced effects



Enhanced by high
variance
Irrespective of
intensity, whereas
regular disturbance
decrease cover

**Temporal variance may drastically change
the effect of disturbance intensity**

IDH

The intermediate disturbance hypothesis was formulated by S.J. Connell (1978) to explain the high diversity of rain forests and coral reefs.

1 – when disturbance is rare (low frequency) and weak (low intensity), strong competitors win. Species richness is therefore reduced. (the assumption is that a hierarchy of competitors exists, and strong competitors occupy the space efficiently).



1	1	1	1	1	3
1	1	1	1	3	3
1	1	1	5	5	5
1	1	1	2	2	2
4	1	6	6	8	8
4	4	6	7	8	8




1	1	1	1	1	3
1	1	1	1	3	3
1	1	1	1	3	3
1	1	1	1	2	2
4	1	1	1	1	8
4	4	6	6	8	8

2 – when disturbance is very intense and frequent, strong competitors are reduced or excluded, and new settlers among weak competitors colonize the space. Species richness is again reduced because some species lack, and only few species tolerate high level of disturbance



3 – finally, when disturbance regime has intermediate strength and frequency, strong weak competitors coexist, since disturbance is not so high to cause the local extinction of the former, but sufficient to create patches available for the latter.



1	1	1	1	1	3
1	1	1	1	3	3
1	1	1	5	5	5
1	1	1	2	2	2
4	1	6	6	8	8
4	4	6	7	8	8




13	1	1	11	11	3
1	1	1	11	3	3
10	10	10	5	5	5
4	10	2	2	2	2
4	6	6	6	8	8
4	4	6	7	8	8

Competitive
exclusion


High stress
Extinction

Richness




Patch dynamics

isolated



Recolonization: Arrival of drifting propagules from the water column

non-isolated



Recolonization: Vegetative growth from neighbours

discrete pieces of substratum that were surrounded by water (isolated patches), and areas that were cleared within a background of other sessile organisms (nonisolated patches).

Non-isolated


Rate of
recolonization



Keough 1984

Isolation and size

Isolated



Recovery after disturbance


High intertidal zone (N Pacific coast of USA)
Manipulation of diversity and different level of disturbance (Allison, 1997)

Diversity Treatment Code	Diversity Treatment level	Algal groups manipulated			Average species richness (SE)
		Fucoids	Foliose Reds	Low abundance species	
H:+F+R+M	high	+	+	+	27.4 (1.81)
M1:-F+R+M	moderate	-	+	+	24.3 (2.00)
M2:+F+R-M	moderate	+	+	-	18.9 (0.43)
L1:+F-R-M	low	+	-	-	15.0 (1.02)
L2:-F+R-M	low	-	+	-	13.3 (0.75)





Experimental removal of different groups (Fucoid, Red Algae, other Macroalgae)

Simulation of thermal stress and dessiccation following heat wave




Results



Reduction in biomass increased with disturbance intensity in general, but depended on group composition of assemblages. For example, reduction was lower for red algae.

Recovery depended on the initial diversity and the intensity of disturbance, but this dependence was strongly related to the characteristics of the species removed. No additional variations were related to the differences in the number of species

Interactions with biological processes




Three sites × three water depths (6, 12, 18 m). Y-axes on the left indicate cover values (mean ± SE) of the sessile communities: reef-building corals and other benthic components. Y-axes on the right indicate densities (mean ± SE) of coral-predators: populations of the outbreaking seastar *Acanthaster* and butterflyfish assemblages. Arrows on the x-axes indicate the occurrence of the tropical cyclone Oli.

Kayal et al 2012




(A) An *Acanthaster planci* observed on a living tabular coral from the genus *Acropora*.



(B) A partially-killed coral from the genus *Acropora* bearing feeding-scars left by successive predation events by *Acanthaster*:

- 1) live portion of the colony bearing the pigmented coral tissue,
- 2) freshly killed portion of the colony deprived of its pigmented living tissue (<1 day post-predation),
- 3) recently killed portion of the colony covered by early colonizing algae and cyanobacteria (~10 days post-predation),
- 4) dead portion of the colony killed long ago and covered by turf algae (>3 weeks post-predation).



(A) Corals dominate the healthy reef (coral cover >40%).

(B) Algae have colonized dead coral skeletons following severe predation by the seastar *Acanthaster* (~10% coral cover).

(C) Mostly dead and weakened coral skeletons were swept away by a cyclone occurring at the end of the seastar outbreak and colonizing algae once again dominate the devastated reef (~5% coral cover).

Summary

- Disturbance is an important factor interacting with biological processes such as competition and predation in shaping marine community assembly
- Disturbance begets spatial, temporal and environmental heterogeneity, and this sustain biodiversity within certain levels
- Effects of disturbance depend on its features, such as intensity and frequency, but spatio-temporal variance of perturbations plays also a crucial role
- Recovery after disturbance, and sometimes the effect of disturbance itself, are strongly related with biological and ecological traits of species composing disturbed assemblages or the available diversity pool
- Biological processes, such as predation, though not being proper disturbance, act similarly and may interact with disturbance
- Recovery after disturbance is related to size of disturbed patches and the potential mechanisms of recolonization or reoccupation