

Condensed Matter Physics I
Final written test
academic year 2010/11
July 12, 2011

(Time: 3 hours)

Exercise 1: Free electrons - Sommerfeld model

Copper (Cu) in normal temperature and pressure conditions is a metal with FCC structure and mass density of about 8.96 g cm^{-3} . It has an average mass number of 63.55 and Fermi energy of 7.00 eV.

1. From the given Fermi energy, derive the density n of the free electrons.
2. Using the Sommerfeld expansion, calculate the electronic contribution to the specific heat at room temperature.
3. Using now its mass density and its mass number, calculate the *atomic* density n_{at} . Using this result and the result in (1), calculate the average number of free electrons per atom. Is it what you would expect?

Exercise 2: Bravais lattices with basis

The copper-oxide high-Tc superconductors contain CuO_2 planes that can be idealized as shown in Figure (a).

1. Write a possible choice of basis vectors, describe and sketch the primitive cell.
2. Assuming that the atomic form factors are such that $f_{Cu} = 2f_O$, calculate the geometrical structure factor $S(\mathbf{K})$ on the reciprocal lattice vectors, and specify for which \mathbf{K} it is maximum and for which ones it vanishes.
3. There are other compounds having a similar structure, but with non equivalent atoms in the oxygen positions, as indicated in Figure (b). Specify in this case which is the primitive cell, the lattice parameter and the basis vectors of the reciprocal space.

Exercise 3: Band structures for free electrons

Consider the BCC lattice.

1. Plot the contours of the 1st Brillouin zone in the $k_x k_y$ plane, specifying the relevant point.
2. Write explicitly the expression of the first energy band $\mathcal{E}_1(\mathbf{k})$ for free electrons from Γ to $N = \frac{2\pi}{a}(1/2, 1/2, 0)$ as a function of the modulus of \mathbf{k} and plot it in the reduced zone scheme.
3. Calculate and plot the other three lowest bands, $\mathcal{E}_n(\mathbf{k})$ for $n = 2, 3, 4$.

Exercise 4: Tight binding

Consider a square lattice with one atom and one s orbital on each site. Neglect overlap between orbitals on different sites. Neglect also hopping integrals between atoms that are not nearest neighbors (NN) or next nearest neighbors (NNN) and let's indicate with γ_{NN} and γ_{NNN} these two non zero integrals.

1. Write the explicit expression for the dispersion of the energy band $E(\mathbf{k})$.
2. Plot $E(\mathbf{k})$ in the first Brillouin zone along the line ΓAB , where $\Gamma = (0,0)$, $A = (\pi/a, 0)$ and $B = (\pi/a, \pi/a)$ for $\gamma_{NNN}/\gamma_{NN} = 1/4$.
3. Calculate the elements of the inverse mass tensor for $\gamma_{NNN}/\gamma_{NN} = 1/4$ at the minimum and at the maximum of the band. Where is the effective mass larger?

NOTE:

- Give all the steps necessary to understand in detail the solution procedure. Answers with the final result only or with insufficient details will not be considered valid.
- When required, numerical evaluations should be given exactly with 3 significant figures, if not otherwise indicated.