

SYNAPTIC PLASTICITY

ROBERT C. MALENKA

The most fascinating and important property of the mammalian brain is its remarkable plasticity, which can be thought of as the ability of experience to modify neural circuitry and thereby to modify future thought, behavior, and feeling. Thinking simplistically, neural activity can modify the behavior of neural circuits by one of three mechanisms: (a) by modifying the strength or efficacy of synaptic transmission at preexisting synapses, (b) by eliciting the growth of new synaptic connections or the pruning away of existing ones, or (c) by modulating the excitability properties of individual neurons. *Synaptic plasticity* refers to the first of these mechanisms, and for almost 100 years, activity-dependent changes in the efficacy of synaptic communication have been proposed to play an important role in the remarkable capacity of the brain to translate transient experiences into seemingly infinite numbers of memories that can last for decades. Because of its fundamental importance, there has been an enormous amount of work describing the many forms of synaptic plasticity and their underlying mechanisms.

Synaptic transmission can either be enhanced or depressed by activity, and these alterations span temporal domains ranging from milliseconds to enduring modifications that may persist for days or weeks and perhaps even longer. Transient forms of synaptic plasticity have been associated with short-term adaptations to sensory inputs, transient changes in behavioral states, and short-lasting forms of memory. More lasting changes are thought to play important roles in the construction of neural circuits during development and with long-term forms of memory in the mature nervous system. Given these diverse functions, it is not surprising that many forms and mechanisms of synaptic plasticity have been described. In this chapter, I provide a brief overview of some of the forms of synaptic plasticity found at excitatory synapses in the mammalian brain, focusing on long-term potentiation (LTP) and long-term depression (LTD).

SHORT-TERM SYNAPTIC PLASTICITY

Virtually every synapse that has been examined in organisms ranging from simple invertebrates to mammals exhibits numerous different forms of short-term synaptic plasticity that last on the order of milliseconds to a few minutes (for detailed reviews, see 1 and 2). In general, these result from a short-lasting modulation of transmitter release that can occur by one of two general types of mechanisms. One involves a change in the amplitude of the transient rise in intracellular calcium concentration that occurs when an action potential invades a presynaptic terminal. This occurs because of some modification in the calcium influx before transmitter release or because the basal level of calcium in the presynaptic terminal has been elevated because of prior activity at the terminal. A second mechanism occurs downstream of calcium elevation in the presynaptic terminal and involves some modulation of the biochemical processes involved in synaptic vesicle exocytosis.

Paired-Pulse Facilitation and Depression

When two presynaptic stimuli are delivered within a short interval, the synaptic response to the second stimulus can be either enhanced or depressed relative to the first stimulus. Paired-pulse depression is commonly observed at all synapses at short (less than 20 milliseconds) interstimulus intervals. It may result from inactivation of voltage-dependent sodium or calcium channels or from a transient depletion of the synaptic vesicles that are “docked” adjacent to the presynaptic plasma membrane, waiting to be released. Many synapses at longer interstimulus intervals (20 to 500 milliseconds) exhibit paired-pulse facilitation that is thought to result from the influx of calcium that occurs in response to the first action potential. One simple idea is that the “residual” calcium left over from the first action potential combines with the calcium influx during the second action potential, and because the relationship between calcium concentration in the terminal and release is highly nonlinear, this small increase in resting calcium may cause substantial facilitation. However, with a single action potential, the increase in resting calcium concentration is very small, and

thus additional mechanisms are likely involved. Currently, there is much interest in the possibility that transient modulation, by activation of protein kinases, of some of the presynaptic phosphoproteins that are known to be involved in the control of transmitter release may play an important role in very short-term synaptic plasticity. For example, knockout mice lacking one or more of the synapsins (3, 4), or lacking the small guanosine triphosphate-binding protein rab3A (5,6), exhibit abnormal short-term synaptic plasticity.

Whether a specific synapse displays paired-pulse facilitation or depression depends on the initial state of the synapse and its recent history of activation. Because these forms of plasticity largely result from changes in the probability of transmitter release, synapses that begin with a very high probability of release tend to show depression, whereas those with a low probability of release exhibit facilitation. Consistent with this idea, activation of presynaptic receptors that cause a decrease in transmitter release almost always causes an increase in the magnitude of paired-pulse facilitation (or even a conversion of paired-pulse depression to paired-pulse facilitation).

Facilitation and Depression Following Trains of Stimuli

Longer-lasting forms of plasticity are observed following repetitive or tetanic stimulation of synapses with prolonged (approximately 200-millisecond to 5-second) trains of stimuli applied at high frequencies (10 to 200 Hz). *Augmentation* and *posttetanic potentiation* refer to enhancements of transmitter release that can last anywhere from seconds (augmentation) to several minutes (posttetanic potentiation). They are thought to result in large part to the buildup of calcium concentration in the presynaptic terminal during the trains of stimuli. This residual calcium may both combine with the calcium influx elicited by subsequent single action potentials and lead to biochemical modifications of proteins in the presynaptic terminal. At some synapses, repetitive activation leads to depression that can last for several seconds or even minutes. As in paired-pulse depression, this generally occurs at synapses that exhibit a high probability of release and is thought to result, at least in part, from a transient depletion of the synaptic vesicles that are poised to be released by an action potential.

In large part because of these short-term forms of synaptic plasticity, the strength of communication between pairs of neurons can be modified even during short bursts of presynaptic activity (e.g., five to ten action potentials at 20 to 50 Hz) (7). The functional relevance of such short-term synaptic dynamics has received much less attention than long-lasting forms of synaptic plasticity and is just beginning to be explored (8). One potential role of these short-term forms of synaptic plasticity is to transform incoming information in the temporal domain into a spatially distrib-

uted code (9,10). Furthermore, given that presynaptic proteins that may be involved in short-term plasticity may be abnormal in neuropsychiatric disorders (11), it is not unreasonable to speculate that abnormal short-term synaptic dynamics in specific neural circuits may contribute to the pathophysiology of any number of mental illnesses.

LONG-TERM SYNAPTIC PLASTICITY

During the last decade, there was enormous interest in elucidating the mechanisms responsible for activity-dependent long-lasting modifications in synaptic strength. The great interest in this topic is largely based on the simple idea that external and internal events are represented in the brain as complex spatiotemporal patterns of neuronal activity, the properties of which result from the pattern of synaptic weights at the connections made between the neurons that are contributing to this activity. The corollary to this hypothesis is that new information is stored (i.e., memories are generated) when activity in a circuit causes a long-lasting change in the pattern of synaptic weights. This simple idea was put forth by Ramon y Cajal almost 100 years ago, but experimental support for such a process was lacking until the early 1970s, when it was demonstrated that repetitive activation of excitatory synapses in the hippocampus caused an increase in synaptic strength that could last for hours or even days (12,13). This long-lasting synaptic enhancement, LTP, has been the object of intense investigation because it is widely believed that LTP provides an important key to understanding the molecular mechanisms by which memories are formed (14,15) and, more generally, by which experience modifies behavior. Furthermore, the activity- and experience-dependent refinement of neural circuitry that occurs during development shares features with learning, and thus a role for LTP in this process has been proposed (16–18).

Long-Term Potentiation

No form of synaptic plasticity has generated more interest and has been more extensively studied than LTP in the CA1 region of the hippocampus. The excitement surrounding this phenomenon derives mainly from four sources. First, there is compelling evidence from studies in rodents and higher primates, including humans, that the hippocampus is a critical component of a neural system involved in various forms of long-term memory (19). Second, several properties of LTP make it an attractive cellular mechanism for information storage (20,21). Like memories, LTP can be generated rapidly and is prolonged and strengthened with repetition. It is also input specific in that it is elicited at the synapses activated by afferent activity and not at adjacent synapses on the same postsynaptic cell. This feature dramatically increases the storage capacity of individual neurons

TABLE 11.1. AREAS OF BRAIN IN WHICH LTP HAS BEEN DEMONSTRATED

Hippocampus	Amygdala
Dentate gyrus	Cerebellum
CA1	Thalamus
CA3	Striatum
Cerebral cortex	Nucleus acumbens
Visual	Ventral tegmental area
Somatosensory	
Motor	
Prefrontal	

that, because synapses can be modified independently, can participate in the encoding of many different bits of information. Third, LTP is readily generated in *in vitro* preparations of the hippocampus, thus making it accessible to rigorous experimental analysis. Indeed, much of what we know about the detailed mechanisms of LTP derives from studies of LTP at excitatory synapses on CA1 pyramidal cells in hippocampal slices. Fourth, LTP has been observed at virtually every excitatory synapse in the mammalian brain that has been studied. Table 11.1 gives a list of the brain regions in which LTP has been demonstrated, and it can be seen that regions thought to be particularly important for various forms of learning and memory are prominent. Although LTP is not a unitary phenomenon, most synapses appear to express a form of LTP that is identical or highly analogous to the LTP found at excitatory synapses on CA1 pyramidal cells. Thus, this form of LTP is the focus of the remainder of this section.

Triggering of LTP: A Critical Role for NMDA Receptors and Calcium

It is well established that the triggering of LTP requires synaptic activation of postsynaptic *N*-methyl-D-aspartate (NMDA) receptors, a subtype of ionotropic glutamate receptor (see Chapter 6) and postsynaptic depolarization, which is accomplished experimentally by repetitive tetanic stimulation of synapses or by directly depolarizing the cell while continuing low-frequency synaptic activation (a so-called “pairing protocol”). How do these requirements explain the properties of LTP? During basal low-frequency synaptic transmission, synaptically released glutamate binds to two different subtypes of ionotropic glutamate receptor, termed AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid) and NMDA receptors, which are often, but not always (see later), co-localized on individual dendritic spines. The AMPA receptor has a channel that is permeable to monovalent cations (Na^+ and K^+), and activation of AMPA receptors provides most of the inward current that generates the excitatory synaptic response when the cell is




FIGURE 11.1. Model for the induction of long-term potentiation (LTP). During normal synaptic transmission (*left*), synaptically released glutamate acts on both NMDA and AMPA receptors. Na^+ flows through the AMPA receptor channel but not through the NMDA receptor channel because of the Mg^{2+} block of this channel. Depolarization of the postsynaptic cell (*right*) relieves the Mg^{2+} block of the NMDA receptor channel and allows Na^+ and Ca^{2+} to flow into the cell. The resultant rise in Ca^{2+} in the dendritic spine is a necessary trigger for the subsequent events leading to LTP.

close to its resting membrane potential (Fig. 11.1). In contrast, as described in Chapter 6, the NMDA receptor exhibits a strong voltage dependence because of the block of its channel at negative membrane potentials by extracellular magnesium. As a result, NMDA receptors contribute little to the postsynaptic response during basal synaptic activity. However, when the cell is depolarized, magnesium dissociates from its binding site within the NMDA receptor channel and allows calcium as well as sodium to enter the dendritic spine (Fig. 11.1). The resultant rise in intracellular calcium is a necessary and perhaps sufficient trigger for LTP. This local source of calcium within the dendritic spine accounts for the input specificity of LTP.

The evidence in support of this model for the initial triggering of LTP is compelling. Specific NMDA receptor antagonists have minimal effects on basal synaptic transmission but block the generation of LTP (22,23). Preventing the rise in calcium by loading cells with calcium chelators blocks LTP (24,25), whereas directly raising intracellular calcium in the postsynaptic cell mimics LTP (25,26). Furthermore, imaging studies have demonstrated that NMDA receptor activation causes a large increase in calcium level within dendritic spines (see 23 for references). The exact properties of the calcium signal that is required to trigger LTP are unknown, but a transient signal lasting only 1 to 3 seconds appears to be sufficient (27). Whether additional sources of calcium, such as release from intracellular stores, are required for the generation of LTP is unclear. It is also uncertain whether additional factors provided by synaptic activity are required. Various neurotransmitters found in the hippocampus such as acetylcholine and norepinephrine can modulate the ability to trigger LTP, and such modula-

tion may be of great importance for the functional in vivo roles of LTP. However, there is no compelling evidence to suggest that any neurotransmitter other than glutamate is required to trigger LTP.

Signal Transduction Mechanisms in LTP

A bewildering array of signal transduction molecules has been suggested to play a role in translating the calcium signal that is required to trigger LTP into a long-lasting increase in synaptic strength (28). However, for only a few of these has compelling evidence of a mandatory role in LTP been presented. A major limitation of much of the work on this topic is that investigators have not adequately distinguished molecules that are key components of the signal transduction machinery absolutely required for LTP from biochemical processes that modulate the ability to generate LTP or play a permissive role. For example, any manipulation that modifies the activity of NMDA receptors may affect LTP. Therefore, several requirements must be met for a signaling molecule to be considered a key component of the biochemical machinery that triggers LTP. First, it must be activated or produced by stimuli that trigger LTP but not by stimuli that fail to do so. Second, inhibition of the pathway in which the molecule participates should block the generation of LTP. Third, activation of the pathway should lead to LTP.

Strong evidence indicates that calcium/calmodulin-dependent protein kinase II (CaMKII) fulfills these requirements and is a key component of the molecular machinery for LTP. Inhibiting its activity pharmacologically by directly loading postsynaptic cells with CaMKII inhibitors or genetic knockout of a critical CaMKII subunit blocks the ability to generate LTP (29–31). Conversely, acutely increasing the postsynaptic concentration of active CaMKII increases synaptic strength and occludes LTP (32,33). Furthermore, CaMKII undergoes autophosphorylation after the triggering of LTP (34,35). That this autophosphorylation is required for LTP was demonstrated by the finding that genetic replacement of endogenous CaMKII with a form lacking the autophosphorylation site prevented LTP (36).

Several other protein kinases have also been suggested to play roles in the triggering of LTP, but the experimental evidence supporting their role is considerably weaker than for CaMKII. Activation of the cyclic adenosine monophosphate-dependent protein kinase (PKA), perhaps by activation of a calmodulin-dependent adenylyl cyclase, has been suggested to boost the activity of CaMKII indirectly by decreasing competing protein phosphatase activity (37,38). This presumably happens by phosphorylation of inhibitor-1, an endogenous protein phosphatase inhibitor (see section on LTD later). Protein kinase C may play a role analogous

to CaMKII, whereas the tyrosine kinases Fyn and Src may indirectly modulate LTP by affecting NMDA receptor function (see 23 for references). The mitogen-activated protein kinase (MAPK) has also been suggested to be important for LTP, albeit in unknown ways.

Expression Mechanisms and LTP

In the 1990s, tremendous confusion and controversy surrounded the seemingly simple issue of whether LTP is caused primarily by presynaptic or postsynaptic modifications. The great challenge to answering this question largely stemmed from the great technical difficulties inherent in examining the changes that occur at individual synapses that are embedded in a complex network in which each cell receives 10,000 or more synapses. Most neurobiologists studying this question agree that the simplest postsynaptic change that could cause LTP would be a change in AMPA receptor function or number, whereas the simplest presynaptic change would be an increase in the probability of neurotransmitter release.

Most studies examining this issue have used electrophysiologic assays, and most of these are inconsistent with the hypothesis that the release of glutamate increases significantly during LTP (23,39). For example, changes in transmitter release probability invariably influence various forms of short-term synaptic plasticity such as paired-pulse facilitation, yet these phenomena are not affected by LTP. To measure glutamate release more directly, two approaches were used. One took advantage of the finding that glial cells tightly ensheath synapses and respond to synaptically released glutamate by activation of electrogenic transporters that generate a current directly proportional to the amount of glutamate released (40,41). The other took advantage of use-dependent antagonists of the NMDA receptor or of a mutant AMPA receptor that lacks the GluR2 subunit. These antagonists decrease synaptic currents at a rate that is directly proportional to the probability of transmitter release (42,43). LTP had no discernible effect on these measures, even though they were affected in the predicted fashion by manipulations known to increase transmitter release.

In addition to these negative findings, certain electrophysiologic and biochemical measures were found to increase during LTP. An increase in the amplitude of miniature electrophysiologic synaptic currents (mEPSCs), which represent the postsynaptic response to the spontaneous release of a single quantum of neurotransmitter, normally indicates an increase in the number or function of postsynaptic neurotransmitter receptors. Such an increase occurs during LTP (44), as well as after manipulations that load dendritic spines with calcium (45,46). A more direct way of monitoring changes in AMPA receptors is to measure the postsynaptic response to direct application of agonist,

and such responses have also been reported to increase, albeit gradually (47).

That LTP is caused by a modification of AMPA receptors is supported by the finding that LTP causes an increase in the phosphorylation of the AMPA receptor subunit GluR1 at the site that is known to be phosphorylated by CaMKII (as well as PKC) (35,48,49). Using expression systems, this phosphorylation has been shown to increase the single-channel conductance of AMPA receptors (50). Because an increase in single-channel conductance of AMPA receptors has been reported to occur during LTP (51), one mechanism that seems likely to contribute to LTP is CaMKII-dependent phosphorylation of GluR1. Consistent with this idea, genetic knockout of GluR1 has been found to prevent the generation of LTP (52).

Silent Synapses and Quantal Synaptic Transmission

Although the evidence presented thus far makes a strong case for postsynaptic changes contributing to LTP, there remained one reproducible experimental result that was difficult to reconcile with this idea. This result derived from experiments that took advantage of the finding that the action potential-dependent release of quanta of neurotransmitter at individual synapses is probabilistic, and therefore release occurs only 10% to 40% of the time. Therefore, if a single synapse or a very small number of synapses is activated once every few seconds, on some of the trials no postsynaptic response is recorded, that is, a so-called failure occurs. An extensively replicated finding is that LTP causes a decrease in the proportion of failures that occur (see 53 for review). Because these failures were assumed to result from failures of neurotransmitter release, it was concluded that LTP involves an increase in the probability of transmitter release.

How can this result be reconciled with all the evidence suggesting that LTP is caused by modulation of AMPA receptors and is not accompanied by an increase in glutamate release? One straightforward idea to explain this apparent discrepancy is the *silent synapse hypothesis* (54), which predicts that some synapses express only NMDA receptors, whereas others express both AMPA and NMDA receptors (Fig. 11.2). Synapses with only NMDA receptors would be functionally silent at hyperpolarized membrane potentials, and thus, when transmitter is released, they would not yield a response. However, LTP at such silent synapses could occur by the rapid expression of AMPA receptors, and such a mechanism would account for the apparent change in failure rate.

There is now strong evidence to support this model of LTP. First, it is possible to record EPSCs that are mediated solely by NMDA receptors, and applying an LTP induction




FIGURE 11.2. Diagram of the silent synapse hypothesis. A synapse is functionally silent when it expresses NMDA receptors but not AMPA receptors in its plasma membrane (bottom). The induction of LTP causes the insertion of AMPA receptors (top) from a putative cytosolic pool. To the right of each diagram are the synaptic currents (i.e., EPSCs) that would be recorded from the corresponding synapse.

protocol at such synapses causes the rapid appearance of AMPA receptor-mediated EPSCs (55,56). Second, immunocytochemical analysis demonstrates that AMPA receptors are not found at a significant percentage of hippocampal synapses, whereas all synapses appear to contain NMDA receptors (see 23 for references). Third, LTP has been shown to cause the delivery of green fluorescent protein (GFP)-tagged AMPA receptors to dendritic spines and the insertion of recombinant AMPA receptors into the synaptic plasma membrane (57,58). Fourth, AMPA and NMDA receptors interact with different proteins at the synapse (59), a finding suggesting that they are regulated independently. Fifth, interference with membrane fusion and presumably exocytosis in the postsynaptic cells impairs LTP (60) and AMPA receptors can interact with proteins involved in membrane fusion (61). These findings are consistent with the idea that membrane fusion may be an important mechanism for the delivery of AMPA receptors to the synaptic plasma membrane.

Virtually all the data presented thus far are consistent with the simple model that LTP, at least initially, is caused by the phosphorylation of AMPA receptors and the delivery or clustering of AMPA receptors within the synaptic plasma membrane (23). These events will presumably occur both at synapses that already contain functional AMPA receptors and ones that are functionally silent. As discussed later, LTD appears to involve the converse process, that is, the removal or endocytosis of AMPA receptors. At the end of this chap-

ter, I discuss how these changes in the number of AMPA receptors at individual synapses may lead to more permanent structural changes, which, in turn, may mediate long-lasting forms of experience-dependent plasticity.

Long-Term Depression

Like LTP, LTD has been demonstrated in a large number of different brain regions and comes in a variety of different forms (62–64). This section focuses on the NMDA receptor-dependent form of LTD found at excitatory synapses on CA1 pyramidal cells and that appears to result, in large part, from a reversal of the processes that mediate LTP.

Triggering of LTD: A Critical Role for NMDA Receptors and Calcium

LTD is normally generated by prolonged (3- to 15-minute) low-frequency (1- to 5-Hz) afferent stimulation or by a pairing protocol during which the cell is held at approximately -50 mV. It shares many features with LTP including input specificity, and it can completely reverse LTP, a process often termed *depotentiation*. Surprisingly, the triggering of LTD requires NMDA receptor activation and an increase in postsynaptic calcium concentration (65,66). This can occur because at resting membrane potentials, the voltage-dependent block of the NMDA receptor channel by magnesium is not 100% effective, and thus, each stimulus will cause a very small amount of calcium entry. Current evidence suggests that the specific properties of the intracellular calcium signal dictate whether LTP or LTD is generated by a specific

pattern of synaptic activity, with LTD requiring a modest rise in calcium (67) and LTP requiring a large rise beyond some critical threshold value (68). The temporal characteristics of this calcium signal may also be important.

Signal Transduction Mechanisms in LTD: A Role for Protein Phosphatases

If calcium is the critical triggering signal for LTD, it must be capable of activating biochemical processes that reverse LTP. Because LTP results, at least in part, from activation of protein kinases, a reasonable hypothesis is that LTD is caused by preferential activation of protein phosphatases, several of which are known to be found at excitatory synapses (69). This idea was first proposed in a theoretic article (70) that presented a specific model that accounted for the bidirectional control of synaptic strength by calcium (Fig. 11.3). It proposed that a balance between CaMKII activity and protein phosphatase 1 (PP1) influenced synaptic strength by controlling the phosphorylation state of unidentified synaptic phosphoproteins. Small rises in calcium favored activation of PP1, whereas large rises were required to increase CaMKII activity. Because PP1 is not directly influenced by calcium, a well-established calcium-dependent phosphatase cascade was invoked to translate the calcium signal into increased PP1 activity (69). This cascade (Fig. 11.3) begins with activation of the calcium/calmodulin-dependent phosphatase calcineurin (also known as protein phosphatase 2B or PP2B). PP2B then dephosphorylates inhibitor 1 (I1), a phosphoprotein that, in its phosphorylated state, is a potent inhibitor of PP1. Thus, activa-




FIGURE 11.3. Model for the signaling cascades responsible for LTD and LTP. LTD is generated when a low rise in Ca^{2+} binds to calmodulin (cam) and activates calcineurin (PP2B). Calcineurin then dephosphorylates inhibitor 1 (I1), which therefore no longer inhibits protein phosphatase 1 (PP1). Active PP1 may act on any number of substrates including camkii or AMPA receptors (*left*). LTP is generated when a high rise in Ca^{2+} activates camkii. Other protein kinases that may also be involved in triggering LTP are protein kinase C (PKC), cAMP-dependent protein kinase (PKA), the tyrosine kinase src (Src), and MAP kinase (MAPK)(*right*).

tion of PP2B causes an increase in PP1 activity through a mechanism of disinhibition. An attractive feature of this model is that the affinity of PP2B for calcium/calmodulin is significantly greater than that of CaMKII. Therefore, PP2B will be preferentially activated by small increases in synaptic calcium levels. Furthermore, a large rise in calcium will preferentially increase protein kinase activity not only by directly activating CaMKII but also by leading to the activation of PKA, which phosphorylates I1 and thereby further inhibits PP1.

Several experimental results are consistent with this model, which currently remains the leading hypothesis for the triggering of LTD. Pharmacologic inhibitors of PP1 or PP2B, when loaded directly into CA1 pyramidal cells, prevent the generation of LTD (71,72). Furthermore, loading cells with the phosphorylated form of I1 blocked LTD. However, although the results of such inhibitor studies are consistent with an important role for protein phosphatases in triggering LTD, other interpretations are possible, and more experimental work testing this hypothesis needs to be performed. Most notably, if PP1 plays a role in LTD analogous to that played by CaMKII in LTP, it should be possible to increase PP1 activity directly in postsynaptic cells and to mimic LTD.

Expression Mechanisms of LTD: A Role for Endocytosis of AMPA Receptors

The silent synapse hypothesis discussed earlier suggested that LTP involves the insertion of AMPA receptors into the synaptic plasma membrane. Consistent with the idea that LTD is a reversal of LTP, there is now considerable evidence that LTD involves removal (i.e., endocytosis) of synaptic AMPA receptors. The first direct evidence that the synaptic localization of AMPA receptors could be rapidly modified was the demonstration that, in cultured hippocampal neurons, short application of glutamate receptor agonists caused rapid a loss of synaptic AMPA receptors with no significant effect on the synaptic localization of NMDA receptors (73). This agonist-induced loss of synaptic AMPA receptors was subsequently shown to result from dynamin-dependent endocytosis (74). Perhaps more important, synaptically triggered LTD in the cultured neurons was accompanied by a decrease in the number of synaptic surface AMPA receptors with no discernible effect on the distribution of NMDA receptors (75). Consistent with these findings, loading CA1 pyramidal cells with inhibitors of endocytosis prevented the generation of LTD (76). These inhibitors also caused a gradual increase in the size of the synaptic responses, whereas inhibitors of exocytosis caused a gradual decrease (76). These results suggest that there is a pool of AMPA receptors that cycle into and out of the synaptic plasma membrane

fairly rapidly and that LTP and LTD may involve a modification of the kinetics of these processes.

Structural Changes and Long-Term Synaptic Plasticity

How are the changes in synaptic strength that occur following the triggering of LTP or LTD maintained for periods lasting weeks or perhaps even years? Although the answer to this question is unknown, recent evidence suggests that the mechanisms described previously may be the initial steps in a more profound anatomic restructuring of synapses, including perhaps the growth of new synapses and the pruning away of preexisting ones. Dendritic spines, the postsynaptic sites that presynaptic boutons contact, have a complex ultrastructure and come in a large variety of shapes (77). With technical advances in microscopy and the use of recombinant fluorescent proteins such as GFP, it has become possible to image individual spines in living neurons. Such experiments have shown that spines are not static but can undergo rapid shape changes (78,79) that are influenced by activity (80). Furthermore, strong synaptic activation of the type that triggers LTP causes an NMDA receptor-dependent growth of spines as well as filopodia, which may be the precursors of spines (81,82). Prolonged application of NMDA to cultured neurons also can cause the loss of spines (83), a finding indicating that, like synaptic strength, the growth and loss of dendritic spines may be under the control of NMDA receptors.

Changes in synapse structure in response to activity also have been extensively explored using more standard electron microscopic techniques. One specific morphologic modification repeatedly associated with increased neuronal activity involves a reorganization of the postsynaptic density (PSD), the electron-dense thickening that contains synaptic glutamate receptors. Specifically, it has been suggested that LTP is associated with an increase in the fraction of synapses that contain discontinuities in their PSDs, termed *perforated synapses* (see 84 for references). This idea is strongly supported by studies in which the synapses activated by strong tetanic stimulation were identified in electron microscopic sections and were found to have larger total PSD surface areas and a larger proportion of perforated synapses (85, 86). Several lines of evidence suggest that this growth of PSDs and their eventual perforation may be initiated by increasing the number of AMPA receptors in the PSD (see 84 for references).

The insertion of new AMPA receptors in the PSD and the generation of perforated synapses may also be early events in the generation of new synapses by a process of splitting or duplication of existing spines (87,88). Consistent with this hypothesis, LTP may be associated with an increase in spine density (84), as well as the frequency of multiple-spine synapses in which two adjacent spines arising from the same dendrite contact a single presynaptic bouton




FIGURE 11.4. Model for sequence of events leading to structural changes following triggering of LTP. Within 10 minutes of LTP induction, AMPA receptors are phosphorylated and inserted into the postsynaptic membrane. This process leads to an increase in the size of the postsynaptic density (PSD) and the production of perforated synapses within 30 minutes. By 1 hour, some perforated synapses split and form multispine synapses. Eventually, retrograde communication, perhaps involving cell-adhesion molecules, leads to presynaptic structural changes and the production of new synapses.

(86). These types of observations have led to a model (Fig. 11.4) (84) that proposes a sequence of events by which the insertion of AMPA receptors leads to a growth and perforation of the PSD and eventually to multiple-spine synapses. Subsequently, retrograde communication perhaps involving cell adhesion molecules would cause appropriate presynaptic structural changes such that a completely new, independent synapse is formed. An attractive corollary to this hypothesis is that LTD involves a shrinkage of the PSD and eventually leads to a complete loss of the dendritic spine and its corresponding presynaptic bouton. However, minimal work on the structural changes associated with LTD has been performed.

An attractive feature of incorporating structural changes into the mechanisms of long-term synaptic plasticity is that it provides a straightforward means by which the activity generated by experience can cause very long-lasting modifications of neural circuitry. Structural changes also may explain the well-known requirement of long-lasting forms of synaptic plasticity for new protein synthesis and gene transcription (see 89 to 91 for reviews).

CONCLUSIONS

This is a brief review of some of the most common forms of synaptic plasticity found at excitatory synapses throughout the mammalian brain. Although relatively little is known about the functional roles of these phenomena, such changes in synaptic function and structure remain the lead-

ing candidates for some of the fundamental mechanisms by which experiences of any type cause the reorganization of neural circuitry and thereby modify thoughts, feelings, and behavior. One hopes that it is also apparent why understanding the mechanisms of synaptic plasticity has important implications for many branches of clinical neuroscience. For example, the development of many pathologic behaviors, such as drug addiction, likely depends on the maladaptive use of neural mechanisms that normally are used for adaptive learning and memory (92). Similarly, the recovery of function following brain injury or the successful pharmacologic and behavioral treatment of mental illness also certainly must result from the reorganization of neural circuitry that is in part achieved by synaptic plasticity mechanisms. Thus further elucidation of the mechanisms of phenomena such as LTP and LTD will continue to have implications for all those interested in the neural basis of cognition and behavior.

ACKNOWLEDGMENTS

I am currently supported by grants from the National Institute of Mental Health, the National Institute of Drug Abuse, and the National Institute on Aging, and I also acknowledge past support from the National Alliance for Research on Schizophrenia and Depression, the McKnight Endowment Fund for Neuroscience, and the Human Frontier Science Program. Thanks to Roger Nicoll, Steven Siegelbaum, Christian Lüscher and Dominique Muller,

with whom I have written previous reviews that provided much of the material for the present chapter.

REFERENCES

1. Zucker RS. Short-term synaptic plasticity. *Annu Rev Neurosci* 1989;12:13–31.
2. Zucker RS. Calcium- and activity-dependent synaptic plasticity. *Curr Opin Neurobiol* 1999;9:305–313.
3. Rosahl TW, Geppert M, Spillane D, et al. Short-term synaptic plasticity is altered in mice lacking synapsin I. *Cell* 1993;75:661–670.
4. Rosahl TW, Spillane D, Missler M, et al. Essential functions of synapsins I and II in synaptic vesicle regulation. *Nature* 1995;375:488–493.
5. Geppert M, Goda Y, Stevens CF, et al. The small GTP-binding protein Rab3A regulates a late step in synaptic vesicle fusion. *Nature* 1997;387:810–814.
6. Geppert M, Südhof TC. RAB3 and synaptotagmin: the yin and yang of synaptic membrane fusion. *Annu Rev Neurosci* 1998;21:75–95.
7. Markram H, Tsodyks M. Redistribution of synaptic efficacy between neocortical pyramidal neurons. *Nature* 1996;382:807–810.
8. Tsodyks M, Pawelzik K, Markram H. Neural networks with dynamic synapses. *Neural Comput* 1998;10: 821–835.
9. Buonomano DV, Merzenich MM. Temporal information transformed into a spatial code by a neural network with realistic properties. *Science* 1995;267:1028–1030.
10. Buonomano DV. Decoding temporal information: a model based on short-term synaptic plasticity. *J Neurosci* 2000;20:1129–1141.
11. Glantz LA, Lewis DA. Reduction of synaptophysin immunoreactivity in the prefrontal cortex of subjects with schizophrenia: regional and diagnostic specificity. *Arch Gen Psychiatry* 1997;54:660–669.
12. Bliss TV, Lomo T. Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Physiol (Lond)* 1973;232:331–356.
13. Bliss TV, Gardner-Medwin AR. Long-lasting potentiation of synaptic transmission in the dentate area of the unanaesthetized rabbit following stimulation of the perforant path. *J Physiol (Lond)* 1973;232:357–374.
14. Eichenbaum H. Spatial learning: the LTP-memory connection. *Nature* 1995;378: 131–132.
15. Eichenbaum H. Learning from LTP: a comment on recent attempts to identify cellular and molecular mechanisms of memory. *Learning Memory* 1996;3:61–73.
16. Crair MC, Malenka RC. A critical period for long-term potentiation at thalamocortical synapses. *Nature* 1995;375:325–328.
17. Singer W. Development and plasticity of cortical processing architectures. *Science* 1995;270:758–764.
18. Katz LC, Shatz CJ. Synaptic activity and the construction of cortical circuits. *Science* 1996;274:1133–1138.
19. Zola-Morgan S, Squire LR. Neuroanatomy of memory. *Annu Rev Neurosci* 1993;16:547–563.
20. Nicoll RA, Kauer JA, Malenka RC. The current excitement in long-term potentiation. *Neuron* 1988;1:97–103.
21. Bliss TV, Collingridge GL. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 1993;361:31–39.
22. Collingridge GL, Kehl SJ, McLennan H. The antagonism of amino acid-induced excitations of rat hippocampal CA1 neurones *in vitro*. *J Physiol (Lond)* 1983;334:19–31.
23. Malenka RC, Nicoll RA. Long-term potentiation: a decade of progress? *Science* 1999;285:1870–1874.
24. Lynch G, Larson J, Kelso S, et al. Intracellular injections of EGTA block induction of hippocampal long-term potentiation. *Nature* 1983;305:719–721.
25. Malenka RC, Kauer JA, Zucker RS, et al. Postsynaptic calcium is sufficient for potentiation of hippocampal synaptic transmission. *Science* 1988;242:81–84.
26. Yang SN, Tang YG, Zucker RS. Selective induction of LTP and LTD by postsynaptic $[Ca^{2+}]_i$ elevation. *J Neurophysiol* 1999;81:781–787.
27. Malenka RC, Lancaster B, Zucker RS. Temporal limits on the rise in postsynaptic calcium required for the induction of long-term potentiation. *Neuron* 1992;9:121–128.
28. Sanes JR, Lichtman JW. Can molecules explain long-term potentiation? *Nat Neurosci* 1999;2:597–604.
29. Malenka RC, Kauer JA, Perkel DJ, et al. An essential role for postsynaptic calmodulin and protein kinase activity in long-term potentiation. *Nature* 1989;340:554–557.
30. Malinow R, Schulman H, Tsien RW. Inhibition of postsynaptic PKC or CaMKII blocks induction but not expression of LTP. *Science* 1989;245:862–866.
31. Silva AJ, Wang Y, Paylor R, et al. Alpha calcium/calmodulin kinase II mutant mice: deficient long-term potentiation and impaired spatial learning. *Cold Spring Harb Symp Quant Biol* 1992;57:527–539.
32. Pettit DL, Perlman S, Malinow R. Potentiated transmission and prevention of further LTP by increased CaMKII activity in postsynaptic hippocampal slice neurons. *Science* 1994;266:1881–1885.
33. Lledo PM, Hjelmstad GO, Mukherji S, et al. Calcium/calmodulin-dependent kinase II and long-term potentiation enhance synaptic transmission by the same mechanism. *Proc Natl Acad Sci USA* 1995;92:11175–11179.
34. Fukunaga K, Muller D, Miyamoto E. Increased phosphorylation of Ca^{2+} /calmodulin-dependent protein kinase II and its endogenous substrates in the induction of long-term potentiation. *J Biol Chem* 1995;270:6119–6124.
35. Barria A, Muller D, Derkach V, et al. Regulatory phosphorylation of AMPA-type glutamate receptors by CaM-KII during long-term potentiation. *Science* 1997;276:2042–2045.
36. Giese KP, Fedorov NB, Filipkowski RK, et al. Autophosphorylation at Thr286 of the alpha calcium-calmodulin kinase II in LTP and learning. *Science* 1998;279:870–873.
37. Blitzer RD, Connor JH, Brown GP, et al. Gating of CaMKII by cAMP-regulated protein phosphatase activity during LTP. *Science* 1998;280:1940–1942.
38. Makhinson M, Chotiner JK, Watson JB, et al. Adenylyl cyclase activation modulates activity-dependent changes in synaptic strength and Ca^{2+} /calmodulin-dependent kinase II autophosphorylation. *J Neurosci* 1999;19:2500–2510.
39. Nicoll RA, Malenka RC. Expression mechanisms underlying NMDA receptor-dependent long-term potentiation. *Ann NY Acad Sci* 1999;868:515–525.
40. Lüscher C, Malenka RC, Nicoll RA. Monitoring glutamate release during LTP with glial transporter currents. *Neuron* 1998;21:435–441.
41. Diamond JS, Bergles DE, Jahr CE. Glutamate release monitored with astrocyte transporter currents during LTP. *Neuron* 1998;21:425–433.
42. Manabe T, Nicoll RA. Long-term potentiation: evidence against an increase in transmitter release probability in the CA1 region of the hippocampus. *Science* 1994;265:1888–1892.
43. Mainen ZF, Jia Z, Roder J, et al. Use-dependent AMPA receptor

- block in mice lacking GluR2 suggests postsynaptic site for LTP expression. *Nat Neurosci* 1998;1:579–586.
44. Oliet SH, Malenka RC, Nicoll RA. Bidirectional control of quantal size by synaptic activity in the hippocampus. *Science* 1996; 271:1294–1297.
 45. Manabe T, Renner P, Nicoll RA. Postsynaptic contribution to long-term potentiation revealed by the analysis of miniature synaptic currents. *Nature* 1992;355:50–55.
 46. Wyllie DJ, Manabe T, Nicoll RA. A rise in postsynaptic Ca^{2+} potentiates miniature excitatory postsynaptic currents and AMPA responses in hippocampal neurons. *Neuron* 1994;12:127–138.
 47. Davies SN, Lester RA, Reymann KG, et al. Temporally distinct pre- and post-synaptic mechanisms maintain long-term potentiation. *Nature* 1989;338:500–503.
 48. Barria A, Derkach V, Soderling T. Identification of the Ca^{2+} /calmodulin-dependent protein kinase II regulatory phosphorylation site in the alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate-type glutamate receptor. *J Biol Chem* 1997;272: 32727–32730.
 49. Mammen AL, Kameyama K, Roche KW, et al. Phosphorylation of the alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor GluR1 subunit by calcium/calmodulin-dependent kinase II. *J Biol Chem* 1997;272:32528–32533.
 50. Derkach V, Barria A, Soderling TR. Ca^{2+} /calmodulin-kinase II enhances channel conductance of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate type glutamate receptors. *Proc Natl Acad Sci USA* 1999;96:3269–3274.
 51. Benke TA, Luthi A, Isaac JT, et al. Modulation of AMPA receptor unitary conductance by synaptic activity. *Nature* 1998;393: 793–797.
 52. Zamanillo D, Sprengel R, Hvalby O, et al. Importance of AMPA receptors for hippocampal synaptic plasticity but not for spatial learning. *Science* 1999;284:1805–1811.
 53. Kullmann DM, Siegelbaum SA. The site of expression of NMDA receptor-dependent LTP: new fuel for an old fire. *Neuron* 1995; 15:997–1002.
 54. Malenka RC, Nicoll RA. Silent synapses speak up. *Neuron* 1997; 19:473–476.
 55. Isaac JT, Nicoll RA, Malenka RC. Evidence for silent synapses: implications for the expression of LTP. *Neuron* 1995;15: 427–434.
 56. Liao D, Hessler NA, Malinow R. Activation of postsynaptically silent synapses during pairing-induced LTP in CA1 region of hippocampal slice. *Nature* 1995;375:400–404.
 57. Shi SH, Hayashi Y, Petralia RS, et al. Rapid spine delivery and redistribution of AMPA receptors after synaptic NMDA receptor activation. *Science* 1999;284:1811–1816.
 58. Hayashi Y, Shi SH, Esteban JA, et al. Driving AMPA receptors into synapses by LTP and CaMKII: requirement for GluR1 and PDZ domain interaction. *Science* 2000;287:2262–2267.
 59. O'Brien RJ, Lau LF, Haganir RL. Molecular mechanisms of glutamate receptor clustering at excitatory synapses. *Curr Opin Neurobiol* 1998;8:364–369.
 60. Lledo PM, Zhang X, Sudhof TC, et al. Postsynaptic membrane fusion and long-term potentiation. *Science* 1998;279:399–403.
 61. Braithwaite SP, Meyer G, Henley JM. Interactions between AMPA receptors and intracellular proteins. *Neuropharmacology* 2000;39:919–930.
 62. Linden DJ. Long-term synaptic depression in the mammalian brain. *Neuron* 1994;12:457–472.
 63. Bear MF, Abraham WC. Long-term depression in hippocampus. *Annu Rev Neurosci* 1996;19:437–462.
 64. Bear MF, Malenka RC. Synaptic plasticity: LTP and LTD. *Curr Opin Neurobiol* 1994;4:389–399.
 65. Dudek SM, Bear MF. Homosynaptic long-term depression in area CA1 of hippocampus and effects of N-methyl-D-aspartate receptor blockade. *Proc Natl Acad Sci USA* 1992;89:4363–4367.
 66. Mulkey RM, Malenka RC. Mechanisms underlying induction of homosynaptic long-term depression in area CA1 of the hippocampus. *Neuron* 1992;9:967–975.
 67. Cummings JA, Mulkey RM, Nicoll RA, et al. Ca^{2+} signaling requirements for long-term depression in the hippocampus. *Neuron* 1996;16:825–833.
 68. Malenka RC. Postsynaptic factors control the duration of synaptic enhancement in area CA1 of the hippocampus. *Neuron* 1991; 6:53–60.
 69. Shenolikar S, Nairn AC. Protein phosphatases: recent progress. *Adv Second Messenger Phosphoprotein Res* 1991;23:1–121.
 70. Lisman J. A mechanism for the Hebb and the anti-Hebb processes underlying learning and memory. *Proc Natl Acad Sci USA* 1989;86:9574–9578.
 71. Mulkey RM, Herron CE, Malenka RC. An essential role for protein phosphatases in hippocampal long-term depression. *Science* 1993;261:1051–1055.
 72. Mulkey RM, Endo S, Shenolikar S, et al. Involvement of a calcineurin/inhibitor-1 phosphatase cascade in hippocampal long-term depression. *Nature* 1994;369:486–488.
 73. Lissin DV, Carroll RC, Nicoll RA, et al. Rapid, activation-induced redistribution of ionotropic glutamate receptors in cultured hippocampal neurons. *J Neurosci* 1999;19:1263–1272.
 74. Carroll RC, Beattie EC, Xia H, et al. Dynamin-dependent endocytosis of ionotropic glutamate receptors. *Proc Natl Acad Sci USA* 1999;96: 4112–4117.
 75. Carroll RC, Lissin DV, von Zastrow M, et al. Rapid redistribution of glutamate receptors contributes to long-term depression in hippocampal cultures. *Nat Neurosci* 1999;2:454–460.
 76. Lüscher C, Xia H, Beattie EC, et al. Role of AMPA receptor cycling in synaptic transmission and plasticity. *Neuron* 1999;24: 649–658.
 77. Harris KM, Kater SB. Dendritic spines: cellular specializations imparting both stability and flexibility to synaptic function. *Annu Rev Neurosci* 1994;17:341–371.
 78. Kaech S, Brinkhaus H, Matus A. Volatile anesthetics block actin-based motility in dendritic spines. *Proc Natl Acad Sci USA* 1999; 96:10433–10437.
 79. Fischer M, Kaech S, Knutti D, et al. Rapid actin-based plasticity in dendritic spines. *Neuron* 1998;20:847–854.
 80. Lendvai B, Stern EA, Chen B, et al. Experience-dependent plasticity of dendritic spines in the developing rat barrel cortex in vivo. *Nature* 2000;404:876–881.
 81. Maletic-Savatic M, Malinow R, Svoboda K. Rapid dendritic morphogenesis in CA1 hippocampal dendrites induced by synaptic activity. *Science* 1999;283:1923–1927.
 82. Engert F, Bonhoeffer T. Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature* 1999;399: 66–70.
 83. Halpain S, Hipolito A, Saffer L. Regulation of F-actin stability in dendritic spines by glutamate receptors and calcineurin. *J Neurosci* 1998;18:9835–9844.
 84. Lüscher C, Nicoll RA, Malenka RC, et al. Synaptic plasticity and dynamic modulation of the postsynaptic membrane. *Nat Neurosci* 2000;3:545–550.
 85. Buchs PA, Muller D. Induction of long-term potentiation is associated with major ultrastructural changes of activated synapses. *Proc Natl Acad Sci USA* 1996;93:8040–8045.

86. Toni N, Buchs PA, Nikonenko I, et al. LTP promotes formation of multiple spine synapses between a single axon terminal and a dendrite. *Nature* 1999;402:421–425.
87. Edwards FA. Anatomy and electrophysiology of fast central synapses lead to a structural model for long-term potentiation. *Physiol Rev* 1995;75:759–787.
88. Carlin RK, Siekevitz P. Plasticity in the central nervous system: do synapses divide? *Proc Natl Acad Sci USA* 1983;80:3517–3521.
89. Frey U, Morris RG. Synaptic tagging: implications for late maintenance of hippocampal long-term potentiation. *Trends Neurosci* 1998;21:181–188.
90. Lisman JE, Fallon JR. What maintains memories? *Science* 1999;283:339–340. Kandel ER, Pittenger C. The past, the future and the biology of memory storage. *Philos Trans R Soc Lond B Biol Sci* 1999;354:2027–2052.
92. Berke JD, Hyman SE. Addiction, dopamine, and the molecular mechanisms of memory. *Neuron* 2000;25:515–532.