

# Histones

## Replication dependent, canonical histones

- Gene organization
- Gene expression control mechanisms in S-phase
- Histone chaperons

## Replication independent histones – histone variants

- A list of histone variants
- Histone variant at centromere
- Histone H3.3 a variant with multiple function
- Histone gammaH2AX and DNA damage
- Histone variants with chromosome specific accumulation

## Coordinated control of histone gene expression and incorporation into DNA

### 1. Histone and cell cycle regulation:

Maintaining a stable and balanced histone pool is of vital importance for appropriate gene regulation, cell cycle progression and genome stability. Excess of free histones has damaging effect → precise control of histone gene expression required → cell cycle

### 2. Controlled production of nucleosomes:

**A. REPLICATION DEPENDENT REGULATION: Canonical histone proteins:** mRNA non-polyadenylated; defined RNA processing mechanisms; expression regulated in histone locus body. Histone mRNA levels increase ca 40-fold increase at the level of histone mRNAs during S phase. At the end of S phase or DNA synthesis is interrupted, cells turned off histone transcription and histone mRNA levels declined rapidly. Replication dependent histone chaperons are linked to DNA polymerase and generate nucleosomes in S-Phase

**B. REPLICATION INDEPENDENT REGULATION: histone variants: Classic polyadenyated RNAs** Nucleosome need also be formed in G1, G2 phase when nucleosome arrays get disturbed (transcription, DNA damage, eviction, etc...). The second class of histones (non canonical histones) is composed of **histone variants that are expressed at a relatively low level throughout the cell cycle, and are therefore regulated in a DNA replication-independent manner. Histone variants have specialized functions!!**

# Histones

## Replication dependent, canonical histones

- Gene organization
- Gene expression control mechanisms in S-phase
- Histone chaperons

## Replication independent histones – histone variants

- A list of histone variants
- Histone variant at centromere
- Histone H3.3 a variant with multiple function
- Histone gammaH2AX and DNA damage
- Histone variants with chromosome specific accumulation

## Histone gene clusters in vertebrates – Canonical histones

Histone genes are organized in histone gene clusters

**Human:**


*Chr.6:* major histone gene locus: HIST1 cluster: 45 core histone genes ; 6 Histone H1 genes

*Chr.1:* minor histone gene locus HIST2 cluster: 6 histone genes

*Chr.1:* minor histone gene locus HIST3 cluster: 3 histone genes

**Yeast:**

2 copies for each core histone




The human and mouse major histone gene cluster. (A) The histone gene cluster on human chromosome 6p21–p22 is shown. The position and direction of transcription of the 55 histone genes in this region are indicated, with the genes for the five histone proteins indicated in the box. Only “real” genes are shown (defined as genes that contain the expected 3' end of histone mRNA). The portion of chromosome 6 is going (left to right) from the centromere to telomere. HISTH4A is the first H4 gene starting from the right and the same is true for the other core histone genes. The numbers are nucleotides from the arbitrary start of the cluster at 0. The regions where there are tightly grouped clusters of histone genes have been expanded. The scale of each section is indicated in kilobases (kb). The position of each of the histone H1 histone gene is indicated with the nomenclature H1a–e, H1t, and the symbols for the core histone genes are in the inset. The asterisks indicate the position of genes present in human and not in mouse.

## Histone genes are organized in clusters in vertebrates


### Gene group: H3 histones (H3) ?

A subgroup of @: "Histones"



**Histone H3:** Histone H3 is one of the five main histone proteins involved in the structure of chromatin in eukaryotic cells. Featuring a main globular domain and a long N-terminal tail, H3 is involved with the structure of the nucleosomes of the 'beads on a string' structure. Histone proteins are highly post-translationally modified however Histone H3 is the most extensively modified of the five histones. The term "Histone H3" alone is purposely ambiguous in that it does not distinguish between sequence variants or modification state. Histone H3 is an important protein in the emerging field of epigenetics, where its sequence variants and variable modification states are thought to play a role in the dynamic and long term regulation of genes. [Source: Wikipedia]

The mapped domains of P68431, encoded by the H3C1 gene, an example gene within the group. [Source: Pfam & UniProt]



HGNC ID (gene)	Approved symbol	Approved name	Previous symbols	Aliases	Chromosome
HGNC:4766	H3C1	H3 clustered histone 1	H3FA, HIST1H3A	H3/A	6p22.2
HGNC:4776	H3C2	H3 clustered histone 2	H3FL, HIST1H3B	H3/I	6p22.2
HGNC:4768	H3C3	H3 clustered histone 3	H3FC, HIST1H3C	H3/c, H3.1	6p22.2
HGNC:4767	H3C4	H3 clustered histone 4	H3FB, HIST1H3D	H3/b	6p22.2
HGNC:54427	H3C5P	H3 clustered histone 5, pseudogene			6p22.2
HGNC:4769	H3C6	H3 clustered histone 6	H3FD, HIST1H3E	H3/d, H3.1	6p22.2
HGNC:4773	H3C7	H3 clustered histone 7	H3FI, HIST1H3F	H3/i	6p22.2
HGNC:4772	H3C8	H3 clustered histone 8	H3FH, HIST1H3G	H3/h	6p22.2
HGNC:18982	H3C9P	H3 clustered histone 9, pseudogene	HIST1H3PS1	dJ45P21.6, H3F3AP1, p36	6p22.2
HGNC:4775	H3C10	H3 clustered histone 10	H3FK, HIST1H3H	H3/k, H3F1K	6p22.1
HGNC:4771	H3C11	H3 clustered histone 11	H3FF, HIST1H3I	H3/f, H3.1	6p22.1
HGNC:4774	H3C12	H3 clustered histone 12	H3FJ, HIST1H3J	H3/j	6p22.1
HGNC:25311	H3C13	H3 clustered histone 13	HIST2H3D		1q21.2
HGNC:20503	H3C14	H3 clustered histone 14	H3F2, H3FM, HIST2H3C	MGC9629, H3/m, H3, H3.2, H3/M	1q21.2
HGNC:20505	H3C15	H3 clustered histone 15	HIST2H3A	H3/n, H3/o	1q21.2
HGNC:43735	H3Y1	H3.Y histone 1		H3.Y, H3.Y.1	5p15.1
HGNC:43734	H3Y2	H3.Y histone 2		H3.X, H3.Y.2	5p15.1
HGNC:1851	CENPA	centromere protein A		CENP-A, CenH3	2p23.3
HGNC:32060	H3-2	H3.2 histone (putative)	HIST2H3PS2	p06	1q21.1
HGNC:4764	H3-3A	H3.3 histone A	H3F3, H3F3A	H3.3A	1q42.12
HGNC:4765	H3-3B	H3.3 histone B	H3F3B	H3.3B	17q25.1
HGNC:4778	H3-4	H3.4 histone	H3FT, HIST3H3	H3t, H3/g, H3.4	1q42.13
HGNC:33164	H3-5	H3.5 histone	H3F3C	H3.5	12p11.21

Canonical H3

Histone H3 variants with special function

In vertebrates, there are a total of 10–20 genes encoding each of the core histone proteins.

Each of these genes encodes a unique mRNA, with a **distinct 5' and 3'-UTR**, as well as **nucleotide changes** in the coding region.

All the histone **H4** genes encode the same protein, **histone H3** genes encodes variants with aa changes, there **10–12 different H2a and H2b** proteins are known

<https://www.genenames.org/cgi-bin/genefamilies/set/864>

## Coordinated control expression of replication dependent histones

*S. cerevisiae*

Structure of yeast  
histone variants  
H2AZ and CenH3.


Structure of yeast  
canonical histone genes

The core histone gene promoters contain specialized DNA elements that enable cis-regulation of histone gene expression: UAS (upstream activating sequence) and NEG (negative regulation of expression) **Divergent arrangement** of histone promoters allows coordinated gene expression to get equal amount of all four core histones.

*H. sapiens*

Structure of mammalian  
canonical  
histone genes

Histone promoters also contain specialized cis elements that required for histone gene expression. Histone H2B promoter contains an octamer element (5'-ATTTGCAT-3'), which is bound by transcription activator Oct-1 (octamer-binding factor 1). Histone H4 promoter contains subtype-specific regulatory elements (SSREs)



*HTA1-HTB1* and *HTA2-HTB2*:  
encode H2A-H2B pairs


*HHT1-HHF1* and *HHT2-HHF2*:  
encode H3-H4 pairs;

*SBF*: transcription factor that drives histone gene expression at G1/S transition

*MBF* is required for histone gene activation

## CONTROL OF HISTONE EXPRESSION IN S-PHASE (replication dependent histones)

### Histone synthesis is limited to S-Phase



The transcription of histone gene takes place in a subnuclear organelle termed the **histone locus body (HLB)**, containing factors required for the processing of histone pre-mRNAs which have an **unusual mRNA structure, with a 3'UTR that forms a stem-loop structure instead of a polyA tail**


#### G1 → S transition:

- Transcription increases 5-fold during the G1 to S phase transition
- RNA processing enhanced RNA levels 8-fold
- Total increase: 30–40-fold increase of mature histone mRNAs.

#### S → G2 transition:

At the end of S phase, the half-life of these histone mRNAs drops dramatically in response to a destabilization of the specific 3' ends.

Coilin = most abundant proteins in Cajal bodies = site for chemical RNA modifications and RNA:Protein complex assembly)




Human cells in S-phase  
IF staining for NPAT combined with  
DNA FISH for the HIST1 cluster on Chr.6

## CONTROL OF HISTONE EXPRESSION IN S-PHASE – TRANSCRIPTIONAL REGULATION

### NPAT – TRANSCRIPTIONAL ACTIVATION

#### Histone synthesis is limited to S-Phase

Activation of histone H2B. Oct-1 binds to octamer elements in H2B promoter. During S phase, activated cyclin E/CDK2 complex phosphorylates NPAT. In combination with NPAT, Oct-1 recruits OCA-S to H2B promoter to activate the expression of H2B.




#### CDK2 and the cell cycle:

Entry into S-phase is triggered by the activity of the G1-S Cyclin complex, CyclinE/Cdk2.

After CyclinE/Cdk2 activity has reached its peak in early S-phase, CyclinE/Cdk2 activity drops due to the degradation of the essential CyclinE component


Activation of histone H4. HiNF-P binds to SSRE within H4 promoters and recruits NPAT and RNA polymerase II to activate gene transcription. NPAT recruits the Tip60 histone acetyltransferase complex to acetylate histone H4 at the G1/S-phase transition. At the end of S phase, the tyrosine kinase WEE1 is recruited to histone promoters to phosphorylate H2B tyrosine 37, which evicts NPAT and RNA polymerase II and instead recruits HIRA to repress histone gene expression.



NPAT, Nuclear Protein Ataxia-Telangiectasia Locus; RNAPII, RNA polymerase II; TRRAP, transformation/transactivation domain-associated protein; SSRE, subtype-specific regulatory elements; OCA-S, Oct-1 co-activator in S-phase; HiNF-P, histone nuclear factor P; TBP, TATA-box binding protein.



## CONTROL OF HISTONE EXPRESSION IN S-PHASE – REPLICATION DEPENDENT HISTONE RNA PROCESSING (RHP)



### Schematic view of the replication-dependent histone RNA processing complex

2 conserved mRNA processing signals 50 nucleotides downstream of the histone ORF stop codon:

- stem-loop element: recognized by stem-loop binding protein (**SLBP**)
- purine-rich histone downstream element (**HDE**): paired by the **U7 snRNP**

A 100 kDa zinc finger protein (**ZFP-100**) forms a connection between these two factors and stabilizes the complex.

The U7 snRNP contains a ring-like structure composed of five Sm proteins and the U7-specific Sm-like proteins, Lsm10 and Lsm11.

A N-terminal extension of Lsm11 additionally binds the 68 kDa subunit of mammalian cleavage factor I (**CFIm68**) and **FLASH**. Together, Lsm11 and **FLASH** provide a docking platform for the **Heat Labile Factor (HLF)** which consists of proteins that are also involved in cleavage/polyadenylation (**CPA**): symplekin (**SYMPK**), Cleavage Stimulation Factor 64 kDa subunit (**CstF64**) and all six subunits of Cleavage and Polyadenylation Factor (**CPSF**).


**CPSF-73 is the endonuclease that cleaves the pre-mRNA after a CA dinucleotide.**

While **SLBP** binds to the 5' side of the hairpin, the 3' side is occupied by **3' hExo**, an exonuclease that trims the 3' end of the mRNA after cleavage and may also be involved in histone mRNA degradation.

If the canonical histone processing site is not recognized, a downstream **polyadenylation signal** can be used by the **CPA** machinery, resulting in polyadenylated histone transcripts. Poly-adenylated transcripts remain nuclear and get rapidly degraded by the nuclear exosome. Only a small fraction can be detected on polysomes.

## CONTROL OF HISTONE EXPRESSION IN S-PHASE – REPLICATION DEPENDENT HISTONE RNA PROCESSING (RHP)

### HBP (SLBP) - REGULATION OF RNA METABOLISM



**HBP (histone RNA binding protein; also called SLBP) protein itself is cell cycle regulated.**

SLBP mRNA is synthesized constantly throughout the cell cycle, but **HBP becomes translated just prior to S-phase entry and the protein is degraded at the end of S-phase**

Histone mRNA 3'-end processing requires the RNA-binding protein **HBP (also called SLBP)**, which binds to the conserved hairpin structure in histone pre-mRNA, and the U7snRNP, which binds to a sequence element downstream of the cleavage site.



Together with other factors they position the nuclease CPSF73 for cleavage to produce histone mRNA ending immediately after the stem loop.

After nuclear export, HBP interacts with SLIP1 and other translation initiation factors to form a closed-loop structure for efficient translation.

This structure is disrupted, presumably when histone mRNA decay is initiated, for example at the end of S-phase.

Addition of an oligo(U) tail by the terminal uridylyl transferase ZCCHC11 is an early step in decay, which involves decapping followed by 5'→3' degradation by Xrn1 or 3'→5' decay by the exosome.

## FORMATION OF SUBNUCLEAR STRUCTURE = HISTONE LOCUS BODY




**Histone locus bodies (HLBs)** are subnuclear sites containing the clustered RD histone genes and components of the corresponding gene expression and processing machinery.

**A. In G1, only two HLBs corresponding** to the major histone cluster on chromosome 6 can be stained by antibodies against HLB marker proteins such as FLASH. As NPAT is not phosphorylated, the interaction with FLASH is weak. The level of SLBP is low, and the U7 snRNP, interacting with hnRNP UL1, inhibits histone transcription. Transcription is minimal and RNA 3' processing is inefficient so that most transcripts from the Replication dependent histone mRNAs become polyadenylated at cryptic polyA signals downstream from their regular cleavage sites.

**B. In S phase, NPAT gets phosphorylated by cyclinE/Cdk2** which stabilizes the interaction with FLASH. The concentration of factors in HLBs increases, and two more HLBs corresponding to the minor cluster on chromosome 1 can be detected. SLBP is up-regulated, and FLASH interacts with the U7 snRNP, hence providing a platform for the recruitment of the now activated HLF.

## CONTROL OF HISTONE EXPRESSION IN S-PHASE

### DROSOPHILA: HERS and Su(var)3-9




**Drosophila:** The histone gene-specific epigenetic repressor in late S-phase (HERS) protein becomes phosphorylated by the late S-G2 Cyclin complex CyclinA/Cdk1, which localizes it to the histone genes where it acts to silence histone genes after S-phase.

Cdk-activated HERS silences histone gene expression in late S phase through recruitment of Su(var)3-9/HP1 repressor complex.

**Drosophila:** also presence of HLB

## REPLICATION COUPLED HISTONES



**Figure 4.** Distribution of old and new nucleosomes at a replication fork. Old nucleosomes (gray disks) are randomly distributed behind the replication fork and new nucleosomes (cyan disks) are deposited in the gaps. CAF-1-mediated nucleosome assembly is depicted on the leading and lagging strand in magnification. DNA polymerase (green); replication processivity clamp, PCNA (gray ring); histone H3-H4 tetramers (cyan); newly synthesized DNA (red lines).

### REPLICATION COUPLED (RCs) HISTONES:

#### H2A, H2B, H3, H4

**Are incorporated into new and old DNA strand during DNA replication**

Chromatin assembly factor 1 (CAF-1) is a HISTONE CHAPERON that is associated with PCNA. → Facilitates the formation of new nucleosomes  
The assembly of a nucleosome consists of the loading of an (H3-H4)<sub>2</sub> tetramer (tetrasome) that is followed by the addition of 2 H2A-H2B dimers.

### REPLICATION INDEPENDENT (RIs) HISTONES:

**Are incorporated independently of DNA replication**


RI histones require the displacement of a preexisting nucleosome unit (active displacement or “loss”)

RI histones can **reset the epigenetic state of a pre-existing nucleosome**

Histone variants with high similarity to “normal” histone are also incorporated into canonical octamers during replication. However concentration of variants is very low → no big effect on chromatin structure

**HOWEVER: HISTONE CHAPERONS EXIST THAT ENSURE CONCENTRATED INCORPORATION OF HISTONE VARIANTS AT DEFINED SITES → CONTROLLED LOCAL CONCENTRATION → ALTERATION OF CHROMATIN STRUCTURE**

## Incorporation of replication coupled histones by CAF-1 histone chaperone



Humans CAF-1 subunits: p150, p60, and p48,  
 Budding yeast CAF-1 subunits: Cac1, Cac2, and Cac3

CAF-1 p150 interacts with PCNA

- The nucleosome assembly mechanism of CAF-1 is activated by H3-H4 binding, which releases the WHD domain from an intramolecular interaction with the acidic region on Cac1.
- DNA binding promotes the association of two CAF-1•H3-H4 complexes to join the histones into a (H3-H4)<sub>2</sub> tetramer.
- In the presence of DNA of sufficient length, the (H3-H4)<sub>2</sub> histones are directly sequestered from CAF-1.
- (H3-H4)<sub>2</sub> are transferred to the DNA to form the **tetrasome**, and the WHD rebinds to the now free acidic region, resulting in its dissociation from DNA.

H2A-H2B can spontaneously associate with tetrasomes in vitro and because CAF-1 itself has significantly lower affinity for H2A-H2B compared to H3-H4, it appears that the **primary role of CAF-1 is to promote the formation of an ordered (H3-H4)<sub>2</sub>•DNA complex, the tetrasome**

# Histones


## Replication dependent, canonical histones

- Gene organization
- Gene expression control mechanisms in S-phase
- Histone chaperons

## Replication independent histones – histone variants

- A list of histone variants
- Histone variant at centromere
- Histone H3.3 a variant with multiple function
- Histone H2AX and DNA damage
- Histone variants with chromosome specific accumulation

## REPLICATION INDEPENDENT (RIs) HISTONES: - HISTONE VARIANTS



**Figure 1.** Histone variants. Protein domain structure for the core histones (H3, H4, H2A, and H2B), linker histone H1, and variants of histones H3 and H2A. The histone-fold domain (HFD) is where histone dimerization occurs. Regions of sequence variation in histone variants are indicated in red. WHD, winged-helix domain.

**REPLICATION INDEPENDENT (RIs) HISTONES:**  
**Are incorporated throughout the cells cycle and independently of DNA replication**

RI histones require the displacement of a pre-existing nucleosome unit

RI histones can reset the epigenetic state of a pre-existing nucleosome

Assure nucleosome organization in interphase cells or terminally differentiated cell (transcription, remodeling)


Used in defined biological process

Histone variants with high similarity to "normal" histone can also be incorporated into canonical octamers during replication. However, concentration of variants is very low  $\rightarrow$  no big effect on chromatin structure

**HOWEVER: HISTONE CHAPERONS EXIST THAT ENSURE CONCENTRATED INCORPORATION OF HISTONE VARIANTS AT DEFINED SITES  $\rightarrow$  CONTROLLED LOCAL CONCENTRATION  $\rightarrow$  ALTERATION OF CHROMATIN STRUCTURE**



## REPLICATION INDEPENDENT (RIs) HISTONES: - HISTONE VARIANTS



**Figure 1.** Histone variants. Protein domain structure for the core histones (H3, H4, H2A, and H2B), linker histone H1, and variants of histones H3 and H2A. The histone-fold domain (HFD) is where histone dimerization occurs. Regions of sequence variation in histone variants are indicated in red. WHD, winged-helix domain.


**Humans:**

**Many isoforms of H2A and H3 exist; H2B and H4 have not diversified**


**Incorporation of variants of histones into nucleosomes dramatically altered chromatin structure**

**Some histone variants are deposited by specialized nucleosome assembly complexes**

**Variation to classic histone is very small (except macroH2A/CENPA)**




# 1. H3 VARIANTS AT THE CENTROMERE: CENP-A: A HISTONE VARIANT THAT DEFINES EUKARYOTIC CENTROMERES



Centromere protein **C 1** is a centromere autoantigen and a component of the inner kinetochore plate. The protein is required for maintaining proper kinetochore size and a timely transition to anaphase.

Centromere protein **B** is a highly conserved protein that facilitates centromere formation. It is a DNA-binding protein that is derived from transposases of the pogo DNA transposon family. It contains a helix-loop-helix DNA binding motif at the N-terminus, and a dimerization domain at the C-terminus. The DNA binding domain recognizes and binds a 17-bp sequence (CENP-B box) in the centromeric alpha satellite DNA. This protein is proposed to play an important role in the assembly of specific centromere structures in interphase nuclei and on mitotic chromosomes. It is also considered a major centromere autoantigen recognized by sera from patients with anti-centromere antibodies.



**CENP-A: RI histone variant, incorporated by the cenH3 specific histone HJURP histone chaperone**

**Kinetochore assembles on CENP-A (Mechanism unknown)**

**CENP-C and CENP-B are non histone centromere founder proteins**

# Histones

## Replication dependent, canonical histones


- Gene organization
- Gene expression control mechanisms in S-phase
- Histone chaperons

## Replication independent histones – histone variants

- A list of histone variants
- Histone variant at centromere
- **Histone H3.3 a variant with multiple function:**
- Histone H2AX and DNA damage
- Histone variants with chromosome specific accumulation

## H3.3: MAINTAINING NUCLEOSOME STRUCTURE IN NON S-PHASE CELLS


### H3.3 INCORPORATION INTO GENOME IS HIGHLY DYNAMIC: AN EXAMPLE



**H3.3 is essential to fill nucleosome gaps in terminally differentiated cells**

**Gene expression is not altered → H3.3 compensates for H3**


## Histone Chaperons: Pathways if RI incorporation of H3.3 into chromatin



DAXX/ATRX →  
H3.3 → repetitive  
elements

**Figure 2.** During RC assembly, the ASF1 chaperone is thought to transfer newly synthesized soluble H3.1-H4 dimers to CAF-1 through direct interactions with its p60 subunit (Tyler et al. 2001; Mello et al. 2002). CAF-1 would then facilitate the assembly of a central (H3.1-H4)<sub>2</sub> tetramer to which two H2A-H2B dimers are juxtaposed by other chaperones to complete a core nucleosomal unit. Similarly, during RI chromatin assembly, ASF1-bound H3.3-H4 would be transferred to HIRA for tetramer formation (Green et al. 2005). Note, however, that the deposition of H3.3 in the *Drosophila* male pronucleus is HIRA-dependent, but ASF1-independent (Bonney et al. 2007) in the histone-rich fertilized egg. Novel alternate pathways for H3.3-H4 deposition include the DAXX chaperone coupled to the ATRX ATP-dependent chromatin remodeler (Drané et al. 2010; Goldberg et al. 2010), as well as targeted H3.3 deposition to regulatory elements by DEK (Sawatsubashi et al. 2010). Boxes highlight queries to consider in future studies.

# 1. H3 VARIANTS: H3.3 A HISTONE VARIANT MARKING, TRANSCRIBED, ACTIVE CHROMATIN



**Figure 7.** H3.3 preferentially localizes to actively transcribed regions of *Drosophila* polytene chromosomes. DAPI staining (red) shows the DNA banding pattern (left), and H3.3-GFP (green) localizes to interbands (middle), which are sites of RNA Pol II localization. The merge (Schwartz and Ahmad 2005) is shown on the right. In each image, the shorter arrow points to a decondensed interband that is enriched in H3.3, and the longer arrow points to a condensed band that lacks H3.3.

**H3:** incorporated during replication (RC histone); co-purifies with CAF-1

**H3.3.:** incorporated during replication (RC histone); Co-purifies with DAXX1


also incorporated by histone chaperones (RI histone)


H3.2: canonical H3 (CAF-1, S-Phase)

H3.1: canonical H3 1aa exchange to H3.2 (CAF1, S-Phase)

H3.3: isoform, 4 aa exchange (DAXX/ATRX, RI histone)

**Only 4 aa exchanges**





**Figure 7.** H3.3 preferentially localizes to actively transcribed regions of *Drosophila* polytene chromosomes. DAPI staining (red) shows the DNA banding pattern (*left*), and H3.3-GFP (green) localizes to interbands (*middle*), which are sites of RNA Pol II localization. The merge ([Schwartz and Ahmad 2005](#)) is shown on the *right*. In each image, the shorter arrow points to a decondensed interband that is enriched in H3.3, and the longer arrow points to a condensed band that lacks H3.3.

H3.3 can form heterodimer with H4 → H3.3-H4 is incorporated by RI assembly

Chromatin is in flux during in G1/G2 and DNA is transcribed → frequent RI chromatin reassembly → enrichment of H3.3 chromatin **in active genes**


Active genes are defined by epigenetic writers (i.e. H3K4 HMTse) that impose active histone marks on H3.3.

Additional positive effect: eventual repressive modifications (H3K9me3) are eliminated by exchanging H3-mod for H3.3 unmodified

**H3.3 carries same modification like classic H3**

## Gene induction triggers histone replacement of H3 with H3.3

Polytene chromosome  
 Heat shock  
 Heatshock response genes  
 activated  
 Formation of "Puffs"  
 (aligned genes are activated  
 Opening of chromatin visible)




Brian E. Schwartz, and Kami Ahmad *Genes Dev.*  
 2005;19:804-814

Gene induction triggers histone replacement. Heat-shock induction triggers rapid puffing and transcription of the HSP70 genes at polytene bands 87A and 87C. Polytene chromosomes from larvae that contain a constitutively expressed H3-GFP construct (A–D, J, K), or a constitutively expressed H3.3core-GFP construct (E–I) were induced for the indicated times (in minutes) and stained with antibodies against **phosphorylated RNA polymerase II (blue)**, which marks activated puffs.


Arrows in A and E indicate the positions of the HSP70 loci at 87A and 87C before induction, while the puffs (B–D, F–H) are bracketed. Asterisks indicate chromosome bands that were used as internal standards for quantifying the summed intensities of histone-GFP signals in puffs. (B, C) **Puffs contain some H3 in the first 5 min of induction, but have much less by the time puffs reach their maximal size (D).** (F–H) **H3.3core-GFP in expanding puffs rapidly increases.** (I) **After 20 min of heat-shock induction, many active HSP puffs (arrows) have large amounts of H3.3core-GFP, while staining throughout the arms is undiminished.** (J, K) Histone H3 modifications that mark heat-shock puffs do not overlap with H3-GFP. **DNA is in red, and histone-GFP in green.**



## Reprogramming of epigenetic marks by dynamic replacement of histone H3 with H3.3 variant




(A) Fertilization induces the transition from metaphase II to anaphase II, leading to completion of oocyte meiosis and emission of a second polar body (which usually degenerates). The sperm nucleus decondenses, so the fertilized egg (zygote) contains two haploid nuclei (male and female pronuclei). In mammals, the pronuclei replicate DNA as they migrate toward each other. They then initiate mitosis, with male and female chromosomes aligning on a common spindle. Completion of mitosis and cytokinesis thus gives rise to a two-cell embryo, with each cell containing a diploid genome.



Early development of the embryo. Division of the fertilized egg results in the formation of daughter cells called blastomeres, which form a ball of cells called a morula. After six or so divisions, a cavity appears at one end of the morula to form the blastocyst. Until this stage, each of the cells divides evenly to produce two daughter cells, each of which retains the same characteristics, that is, they are totipotent. The inner cell mass of the blastula is the source of ESCs (see below). Invagination of one pole of the blastula leads to formation of the primitive germ cell layers, with formation of the gastrula. During gastrulation and later formation of the fetus, the daughter cells lose potential as they gain specialized function. This process is known as determination.


## Reprogramming of epigenetic marks by dynamic replacement of histone H3 variants



Representative pictures of MII oocytes prior to ( A , F , and K ) and after insemination using ICSI ( Day 1 : B , G , and L ; Day 3 : C , H , and M ; Day 5 : D , I , and N ; Day 7 : E , J , and O ). After insemination, the presence of two pronuclei (PN) and two polar bodies (PB) was checked to confirm fertilization ( B , G , and L ). Some embryos arrested at very early stages (e.g., four cell; D and E ), whereas others continued cleaving until the morula stage ( J and O ), when they arrested.

**EPIGENETIC REPROGRAMMING**  
*Gene activation*


# Reprogramming of epigenetic marks by dynamic replacement of histone H3 variants



**Female Gametogenesis:** In females, the total number of eggs ever to be produced are present in the newborn female initially arrested at the diplotene stage of the meiosis I from fetal life through childhood until puberty, when the lutenizing hormone (LH) surges stimulate the resumption of meiosis. **All eggs are arrested at an early stage (prophase I) of the first meiotic division as a primary oocyte (primordial follicle).** Following puberty, during each menstrual cycle, pituitary gonadotrophin stimulates completion of meiosis 1 the day before ovulation. In meiosis 1, a diploid cell becomes 2 haploid (23 chromosomes) daughter cells, each chromosome has two chromatids. One cell becomes the secondary oocyte the other cell forms the first polar body. **The secondary oocyte then commences meiosis 2 which arrests at metaphase and will not continue without fertilization.** At fertilization meiosis 2 completes, forming a second polar body. Note that the first polar body may also undergo this process forming a third polar body.

**Polar Body:** The breakdown of the germinal vesicle indicates a resumption of meiosis and the extrusion of the first polar body (1 PB) indicates completion of the first meiotic division in human oocytes. The polar body is a small cytoplasmic exclusion body formed to enclose the excess DNA formed during the oocyte (egg) meiosis and following sperm fertilization. There are 2-3 polar bodies derived from the oocyte present in the zygote, the number is dependent upon whether polar body 1 (the first polar body formed during meiosis 1) divides during meiosis 2. This exclusion body contains the excess DNA from the reductive division (the second and third polar bodies are formed from meiosis 2 at fertilization). These polar bodies do not contribute to the future genetic complement of the zygote, embryo or fetus. Recent research in some species suggest that the space formed by the peripheral polar body (between the oocyte and the zona pellucida) can influence the site of spermatozoa fertilization.

## Reprogramming of epigenetic marks by dynamic replacement of histone H3 variants




P: paternal pronucleus  
M: maternal pronucleus

H3.2: canonical H3 inserted into pronuclei + 2 cell embryo  
H3.3 RI histone: inserted into into pronuclei + 2 cell embryo

MII oocytes injected with  
flag-tagged H3.1 or H3.2  
or H3.3 variants →  
In vitro fertilization →  
Allow 1-cell and 2 cell  
development →  
Anti-flag IF

## Reprogramming of epigenetic marks by dynamic replacement of histone H3 variants



2 cell stage → blastocyst


H3.1: incorporation into blastocyst chromatin (eu+heterochromatin)

H3.2: incorporation in 2 cell and blastocyst chromatin (eu+heterochr.)

H3.3: incorporation in 2 cell embryos (hetero + euchromatin)  
**incorporation in blastocyst euchromatin** (DNA not condensed)

→ ACTIVATION OF GENE EXPRESSION PARALLELED BY H3.3 INCORPORATION


## Reprogramming by dynamic, replication dependent replacement of histone H3 variants with H3.3



**Blastocyst:** Loss of CAF function (sip150) --> lack of H3.1 and H3.2 incorporation

H3.3 incorporation not affected (increases)


→ H3.3 is an RI histone (incorporated by HIRA; DAXX or DEK)



MII oocytes fertilized →  
 Develop to 1 cell stage →  
 injected with siRNAs →  
 Injection of Flag tagged variants  
 In 1 blastomere of 2 cell embryo  
 → analysis of morula cells

## 2. H3.3 is enriched at repeat elements and contributes to genomic stability

### CHROMATIN REGULATION OF REPETITIVE DNA



In addition to genes, recent studies have shown that H3.3 is enriched at repetitive regions such as

- **Telomeres**
- **Pericentric repeats**
- **Endogenous retroelements**


Deposition at these regions is mediated by the **ATRX/DAXX complex**.

This complex is mutated in many human cancers, with links to a process known as alternative lengthening of telomeres, or ALT.

## DAXX – ATRX insert H3.3 at repeat elements: at heterochromatic telomeres and centromeric repeats

Daxx: histone chaperon

ATRX: ATP nucleosome remodeling complex



H3.3 also enriched at tandem repeat containing sequences (i.e telomere or centromere, ...)

ATRX binds to H3-tails that are H3K4me0 and H3K9me3

ATRX is enriched at telomeres and centromeric repeats


ATRX brings in DAXX → H3.3 RI assembly

There is transcription at telomeres and centromeres → nucleosome turnover

**Figure 8.** Model for maintenance of histone modifications by the concerted action of multiple chromatin regulators via RI replacement with H3.3. We address the question of how a histone modification can be inherited when a nucleosome is lost and replaced. (A) The Suv39h H3K9 methyltransferase (an ortholog of fly Su(var)3-9) is recruited by HP-1 protein, which binds specifically to methylated H3K9. To perpetuate this mark when the nucleosome turns over, we speculate that the ATRX ATPase is recruited to the site via its ATRX-DNMT3-DNMT3L (ADD) domain, which binds with high specificity to methylated H3K9 on tails that entirely lack H3K4 methylation (because there are no H3K4 methyltransferases in this region of the genome). (B) ATRX provides the energy of ATP and works together with the H3.3-specific DAXX histone chaperone complex to incorporate the new nucleosome (Goldberg et al. 2010), or half-nucleosome in the case of partial eviction (Xu et al. 2010). The high local concentration of Suv39h results in a new nucleosome with the same H3K9 methylation as the nucleosome that was lost.




## A model for DAXX-ATRX mediated suppression genomic instability: telomeres as an example



Loss of ATRX/DAXX function → lack of H3.3 incorporation → abnormal chromatin → breaks → induction of homologous recombination

= ALT (alternative lengthening of telomeres)




## A model for DAXX-ATRX mediated suppression genomic instability: telomeres as an example



The MRN complex (MRX complex in yeast) is a protein complex consisting of Mre11, Rad50 and Nbs1 (also known as Nibrin in humans and as Xrs2 in yeast). In eukaryotes, the MRN/X complex plays an important role in the initial processing of double-strand DNA breaks prior to repair by homologous recombination or non-homologous end joining.

**Figure 6 | Model for ATRX-mediated suppression of the ALT pathway.** ATRX together with DAXX deposits histone H3.3 at telomeres, which in turn may facilitate DNA replication through G-quadruplex sequences. The presence of G-quadruplex structures in an ATRX null tumour cell leads to replication fork stalling and collapse, providing a substrate for MRN-dependent homologous recombination and maintenance of telomere length through ALT. ATRX additionally interacts with the MRN complex, facilitating its distribution away from PML bodies and telomeres, further limiting HR.

# ROLE OF H3.3 IN TUMORFORMATION




## ROLE OF H3.3 IN TUMORFORMATION


PanNETs: human pancreatic neuroendocrine tumors: 40% have loss of function mutations in ATRX or Daxx

Loss of ATRX or Daxx is associated with the induction of recombination at telomeres  
→ telomerase independent telomere elongation


Pediatric glioblastoma; Mutations in H3.3 G34R/V associated with ATRX loss → tumor formation



Alternative lengthening of Telomeres (ALT)




Telomerase




Reverse Transcription from Template

## ROLE OF ATRX/DAXX AND H3.3 IN Pancreatic neuroendocrine tumors (PanNETs)



Pancreatic neuroendocrine tumors (PanNETs, PETs, or PNETs), often referred to as "islet cell tumors", or "pancreatic endocrine tumors" are neuroendocrine neoplasms that arise from cells of the endocrine (hormonal) and nervous system within the pancreas. PanNETs are a type of neuroendocrine tumor, representing about one third of gastroenteropancreatic neuroendocrine tumors (GEP-NETs). Many PanNETs are benign, while some are malignant. Aggressive PanNET tumors have traditionally been termed "islet cell carcinoma". PanNETs are quite distinct from the usual form of pancreatic cancer, the majority of which are adenocarcinomas, which arises in the exocrine pancreas. Only 1 or 2% of clinically significant pancreas neoplasms are PanNETs.

## ROLE OF ATRX/DAXX AND H3.3 IN PanNETs



**Figure 1.**

Representative examples of PanNETs assessed by DAXX and ATRX IHC and telomere-specific FISH. **A**, PanNET with preserved nuclear expression for both DAXX (**B**) and ATRX (**C**) and absence of the ALT phenotype (**D**). **E**, PanNET with DAXX loss (**F**), but preserved expression for ATRX (**G**). The loss of DAXX expression correlated with the presence of large, ultrabright intranuclear foci by telomere-specific FISH, consistent with ALT (**H**). **I**, PanNET with preserved expression for DAXX (**J**), but ATRX loss (**K**) and ALT positive (**L**) by telomere-specific FISH.


## ROLE OF ATRX/DAXX AND H3.3 IN PanNETs

**Table 1.** Clinical and pathologic comparison of ALT and DAXX/ATRX status in PanNETs


Patient or tumor characteristics	ALT-positive, n = 98 (31%)	ALT-negative, n = 223 (69%)	P	DAXX/ATRX-negative, n = 84 (26%)	DAXX/ATRX-positive, n = 237 (74%)	P
Gender						
Female	35 (35%)	108 (51%)	0.011 <sup>a</sup>	29 (35%)	121 (51%)	0.011 <sup>a</sup>
Male	63 (65%)	115 (49%)		55 (65%)	116 (49%)	
Mean age (range), years	60.5 (31–85)	58.6 (29–83)	0.195	61.4 (31–85)	58.4 (29–83)	0.050
Mean tumor size (range), cm	5.0 (1.0–15.0)	2.8 (0.6–18.0)	<0.001 <sup>a</sup>	5.0 (1.0–15.0)	2.8 (0.6–18.0)	<0.001 <sup>a</sup>
Functional						
No	95 (97%)	190 (85%)	0.002 <sup>a</sup>	81 (96%)	204 (86%)	0.008 <sup>a</sup>
Yes	3 (3%)	33 (15%)		3 (4%)	33 (14%)	
Location						
Head and uncinata	37 (38%)	90 (40%)	0.300	29 (35%)	98 (41%)	0.300
Body and tail	61 (62%)	133 (60%)		55 (65%)	139 (59%)	
WHO grade						
Low (G1)	28 (29%)	157 (70%)	<0.001 <sup>a</sup>	25 (30%)	160 (68%)	<0.001 <sup>a</sup>
Intermediate (G2)	66 (67%)	66 (30%)		56 (66%)	76 (32%)	
High (G3)	4 (4%)	0 (0%)		3 (4%)	1	
Lymphovascular invasion						
Absent	22 (22%)	163 (73%)	<0.001 <sup>a</sup>	20 (24%)	165 (70%)	<0.001 <sup>a</sup>
Present	76 (78%)	60 (27%)		64 (76%)	72 (30%)	
Perineural invasion						
Absent	55 (56%)	185 (83%)	<0.001 <sup>a</sup>	35 (42%)	191 (81%)	<0.001 <sup>a</sup>
Present	43 (44%)	38 (17%)		49 (58%)	46 (19%)	
Primary tumor (pT) stage						
T1	6 (6%)	110 (49%)	<0.001 <sup>a</sup>	6 (7%)	110 (46%)	<0.001 <sup>a</sup>
T2	28 (29%)	71 (32%)		26 (31%)	73 (31%)	
T3	64 (65%)	42 (19%)		52 (62%)	54 (23%)	
Regional node (pN) stage	n = 96	n = 172		n = 83	n = 185	
N0	39 (41%)	129 (75%)	<0.001 <sup>a</sup>	32 (39%)	136 (74%)	<0.001 <sup>a</sup>
N1	57 (59%)	43 (25%)		51 (61%)	49 (26%)	
Synchronous metastases						
Absent	61 (62%)	209 (94%)	<0.001 <sup>a</sup>	54 (64%)	216 (91%)	<0.001 <sup>a</sup>
Present	37 (38%)	14 (6%)		30 (36%)	21 (9%)	
Metachronous metastases	n = 61	n = 209		n = 54	n = 216	
Absent	28 (46%)	200 (96%)	<0.001 <sup>a</sup>	25 (46%)	203 (94%)	<0.001 <sup>a</sup>
Present	33 (54%)	9 (4%)		29 (54%)	13 (6%)	
ALT						
Negative				0 (0%)	223 (94%)	<0.001 <sup>a</sup>
Positive				84 (100%)	14 (6%)	

<sup>a</sup>Indicates that the value in question is statistically significantly better than the relevant control, where significance is defined by  $P < 0.05$ .

## ALT and LOSS OF DAXX/ATRX mediates PanNET aggressiveness



No. at risk	0	3	6	9	12	15
Negative: 209	183	79	40	17	4	
Positive: 61	31	12	5	1	0	



No. at risk	0	3	6	9	12	15
Positive: 216	189	80	41	17	4	
Negative: 54	25	11	4	1	0	

### disease-free survival

In cancer, the length of time after primary treatment for a cancer ends that the patient survives without any signs or symptoms of that cancer. In a clinical trial, measuring the disease-free survival is one way to see how well a new treatment works. Also called DFS, relapse-free survival, and RFS.



# Histones

## Replication dependent, canonical histones


- Gene organization
- Gene expression control mechanisms in S-phase
- Histone chaperons

## Replication independent histones – histone variants



- A list of histone variants
- Histone variant at centromere
- Histone H3.3 a variant with multiple function:
- **Histone H2AX and DNA damage**
- Histone variants with chromosome specific accumulation

## ROLE OF H2A.X IN DNA DAMAGE RESPONSE

### HISTONE H2A VARIANTS




## Epigenetics and Histone Variants meet to organize DNA DAMAGE RESPONSE



DNA double-strand breaks (DSBs) are cytotoxic damages that can be repaired either by the homologous recombination (HR) pathway or by the non-homologous end-joining (NHEJ) pathway. NHEJ, although faster than HR, is less accurate. The early divergent step between the two pathways is end resection, and this step is regulated by numerous factors. In particular, BRCA1 and 53BP1 play a role in determining the balance between the two pathways. 53BP1 restricts resection and promotes NHEJ.


The MRN complex binds avidly to double-strand breaks both in vitro and in vivo and may serve to tether broken ends prior to repair by non-homologous end joining or to initiate resection prior to repair by homologous recombination. Recruits ATM. The MRN complex also participates in activating the checkpoint kinase ATM in response to DNA damage.

## Epigenetics and Histone Variants meet to organize DNA DAMAGE RESPONSE



Cells cut with laser →  
DNA breaks across nucleus →  
phosphorylation of H2AX at Ser 139 by ATM,  
ATR and DNA-PK =  $\gamma$ H2AX

### ACTIVATION OF DNA DAMAGE RESPONSE




$\gamma$ -H2AX

*H2AX is distributed throughout the genome.  
DNA damage -> phosphorylation of H2AX  
by ATM/ATR/DNA PK that spreads max 1 Mb from DNA damage site*


**=DNA DAMAGE FOCI**

## Epigenetics and Histone Variants meet to organize DNA DAMAGE RESPONSE

DNA Damage → Exposure of DNA breaks → activation of PI3K family kinases (ATR/ATM/DNA PK) → phosphorylation cascade



DNA Damage Foci in genomic DNA of cell nucleus





$\gamma$ H2AX

53BP1  
MRE11  
P-ATM

Histone H2AX is a substrate of several phosphoinositide 3-kinase-related protein kinases (PIKKs), such as ATM (ataxia teleangiectasia mutated), ATR (ATM and Rad3-related), or DNA-dependent protein kinase (DNA-PK). Serves as a platform to recruit DNA damage response factors

## ROLE OF H2A.X IN DNA DAMAGE RESPONSE





### Spatial organization of DDR protein accumulation at DNA DSBs.

(A) DDR signal spreading. DDR proteins initially accumulate at DSB sites and then spread at distance via a positive feedback loop involving MDC1, which binds  $\gamma$ H2AX, the MRN complex, and ATM kinase, which phosphorylates additional H2AX molecules further away from the break site.

(B) Regional distribution of DDR proteins around DSBs. Factors involved in ATR signaling accumulate proximal to the break site on ssDNA generated by DNA end resection, while ATM signaling factors localize on flanking chromatin regions.


## ROLE OF H2A.X IN DNA DAMAGE RESPONSE



$\gamma$ H2A.X  
53BP1  
MRE11  
P-ATM

## Epigenetics and Histone Variants meet to organize DNA DAMAGE RESPONSE

Several lines of evidence suggest the critical role of H2AX phosphorylation at DSB sites for nuclear foci formation and induction of DSB repair.



- (i) H2AX-knockout cells manifested impaired recruitment of NBS1, 53BP1, and BRCA1 to irradiation-induced foci
- (ii) H2AX loss of function mouse thymocytes show an increase in chromosomal aberrations. Mouse embryonic stem (ES) cells deficient in H2AX phosphorylation have alterations in efficiency of DNA repair by NHEJ or HR. As a result of these defects in DNA damage repair, cells show increased sensitivity to DNA damage.
- (iii) H2AX knock-out mice show male-specific infertility and reduced levels of secondary immunoglobulin isotypes, suggesting defects in class switch recombination (CSR). It was shown that efficient resolution of DSBs induced during CSR in lymphocytes requires H2AX, and its absence is associated with chromosome abnormalities involving the immunoglobulin locus.

These facts suggest that H2AX might serve as a docking site for DNA damage/repair proteins and functions to promote DSB repair and genome stability.



# Histones


## Replication dependent, canonical histones

- Gene organization
- Gene expression control mechanisms in S-phase
- Histone chaperons

## Replication independent histones – histone variants


- A list of histone variants
- Histone variant at centromere
- Histone H3.3 a variant with multiple function:
- Histone H2AX and DNA damage
- **Histone variants with chromosome specific accumulation**

## ROLE OF macroH2A AND H2A.B ON THE X-CHROMOSOME




# MacroH2A and H2A.B and X inactivation

X inactivation is associated with the exchange of H2A variants




macroH2A is enriched at the Xi



**Figure 11.** H2A variants and the inactive X chromosome of human females. (A) macroH2A (red) stains discrete regions of the inactive X chromosome that alternate with a marker for heterochromatin (histone H3K9me3). (B) H2A.B (green) is excluded from the inactive X chromosome (red dot with arrowhead pointing to it). (C) Same nucleus as in B, but stained with DAPI to show chromatin. (A, Reprinted, with permission, from [Chadwick and Willard 2004](#), © National Academy of Sciences; B,C, reprinted, with permission, from [Chadwick and Willard 2001](#), © 2001 The Rockefeller University Press. Originally published in *Journal of Cell Biology* **152**: 375–384. doi: 10.1083/jcb.152.2.375.)

## ROLE OF H2A.Z IN CHROMATIN DYNAMICS



### Different heterodimers possible:

1. H2A.Z/H2B + H2A.Z/H2B = ZZ
2. H2A.Z/H2B + H2A/H2B
3. H2A/H2B + H2A/H2B

### Inserted of RI H2AZ into chromatin by:

- INO80 and SWR-C complexes (chromatin remodelers with ATPase domain);
- **histone chaperon**: human YL1; yeast: Chz1

H2A.Z shows less adherence to DNA and is frequently localized **around** the transcriptional start site:



Yeast: ZZ: 32%

ZA: 24%

AA: 44%


→ >50% of genes have a H2A.Z at transcription start sites! WHY?

## NUCLEOSOME DEPLETED REGIONS (NDRs)



**Figure 2** The different physiological outcomes of nucleosome sliding. (a) Nucleosome-remodeling enzymes can introduce and maintain regular spacing of nucleosomes. (b) The phasing of nucleosomal arrays with respect to a nucleosome-depleted region (NDR), a prominent feature of promoters, depends on nucleosome-remodeling enzymes. (c) Nucleosome-sliding activity regulates the accessibility of DNA sequences by positioning individual nucleosomes.

## NUCLEOSOME DEPLETED REGIONS (NDRs)





### **Nucleosome-depleted regions at the 5' and 3' ends of transcripts.**

As observed in eukaryotes, Both 5' and 3' end profiles are overlaid in this figure for comparison.

The 5' NDR is, on average, more depleted and longer. DOI:<http://dx.doi.org/10.7554/eLife.00078.005>

## ROLE OF H2A.Z IN CHROMATIN DYNAMICS



**Figure 9.3** DNA strands separate to form a transcription bubble. RNA is synthesized by complementary base pairing with one of the DNA strands.

At promoter and +1 position high turn-over of nucleosomes

The initiation site of transcription is nucleosome-free

Nucleosomes containing H2A.Z (or H3.3) are **more instable**; leaving the promoter and +1 position more flexible

During G1, S, G2, M phase nucleosome at +1 position (transcriptional start site) are **not present**.


Around these position H2A.Z is abundant!  
In this way the TSS acquires an open configuration  
- Also when cell pass trough M-Phase

**=TRANSCRIPTIONAL MEMORY TO MARK START SITE IN ACTIVE GENES!!!**

“normal” histone modifications do not precisely mark the start site of transcription

MECHANISMS THAT DECIDES HOW H2A.Z IS INCORPORATED (RI assembly) IS UNKNOWN

## ROLE OF H2A.Z IN CHROMATIN AND EPIGENETIC INHERITANCE IN STEM CELLS



EMBRYONIC STEM CELLS:  
H2A.Z at bivalent promoters  
(colocalization with PcG/Trx)

### Depletion of H2A.Z:

- More stable association of nucleosomes with promoters and enhance
- Reduced activation of expression
- differentiation defects

(b)


Genes involved in	ES	MEF	NPC
Haematopoiesis	Bivalent	K27 or neither	K27 or neither
Mesenchyme	Bivalent	K4	K27 or neither
Adipogenesis	Bivalent	Bivalent	K27 or neither
Neurogenesis	Bivalent	K27 or neither	Bivalent



## H2A.Z IS CELL CYCLE REGULATED AT THE TSS

-1, 0 NDR: transcriptional start site

Brown color: highly expressed gene



Position -2: high peak of H2A.Z

Position -1,0: Nucleosome depleted region

Position +1: peak of H2A.Z

Position -2: low high peak of H2A.Z

Position -1,0: Nucleosome depleted region

Position +1: peak of H2A.Z

Position -2: low high peak of H2A.Z

Position -1,0: Nucleosome depleted region

Position +1: peak of H2A.Z

**EXPERIMENTS: Trophoblasts in G1, S, M Phase**

→ gene expression array +

→ ChIP H2A.Z and ANALYSIS OF OCCUPANCY OF GENES THAT ARE ACTIVE IN G1, S, M PHASE

**H2A.Z around transcriptional start site is cell cycle regulated**