



# 993SM - Laboratory of Computational Physics lecture 8 - part 1 May 5, 2021

**Maria Peressi**

Università degli Studi di Trieste - Dipartimento di Fisica  
Sede di Miramare (Strada Costiera 11, Trieste)

e-mail: [peressi@units.it](mailto:peressi@units.it)

tel.: +39 040 2240242

# The Ising model in the canonical ensemble

- Introduction to the Ising model
- The Ising model in the canonical ensemble: application of Metropolis Monte Carlo Method
- Implementation in a code

M. Peressi - UniTS - Laurea Magistrale in Physics  
Laboratory of Computational Physics - Unit VIII

(from previous lecture)

# Metropolis algorithm in the canonical ensemble

$$T(i \rightarrow j) = \min(1, e^{-\beta\Delta E}) \quad (\text{Metropolis algorithm}),$$

where  $\Delta E = E_j - E_i$ .

1) ERGODICITY implicitly assumed!

2) TEMPERATURE:

If  $E_B > E_A$ , accept the new (higher energy) configuration with probability  $p = e^{-(E_B - E_A)/k_B T}$ . This means that when the temperature is high, we don't mind taking steps in the "wrong" direction, but as the temperature is lowered, we are forced to settle into the lowest configuration we can find in our neighborhood.

# stochastic simulations of an interacting spin ensemble

We need a model for:

- **energy**

- **evolution**

(to build the trajectory / the Markov chain)

# stochastic simulations of an interacting spin ensemble

We need a model for:

- **energy** => Ising model

- **evolution**


(to build the trajectory / the Markov chain)

# Ising model

the simplest model of interacting spin on a lattice

$$\mathcal{H}^{spin} = -J \sum_{i,j=1}^N s_i s_j$$

$$s_i = \pm 1$$



But where does it come from ?

# 2 interacting spins

Consider two fermions,  $\vec{s}_1, \vec{s}_2$  (e.g. electrons in He) described by a hamiltonian not depending explicitly on spin:

$$\mathcal{H} = h_1 + h_2 + V_{12}$$

=> the eigenstates can be in a factorized form (symm/antisymm under particle exchange due to the Pauli exclusion principle):

$$\Psi_{+,-}(1,2) = \Psi_{+,-}^{orb}(1,2) \chi(1,2)$$

In the Hilbert **spin subspace** we can choose a **coupled representation**, referring to the spin sum  $\vec{S}$ , with eigenstates labeled by  $|S, S_z\rangle$  :

$$\Psi_{+,-}(1,2) = \Psi_{+,-}^{orb}(1,2) \chi_{S,S_z}(1,2) = \Psi_{+,-}^{orb}(1,2) |S, S_z\rangle$$

# 2 interacting spins

## Eigenstates

$$\Psi_{+,-}^{orb}(1, 2)|S, S_z\rangle = (\phi(1)\psi(2) + (-)^S \phi(2)\psi(1))/\sqrt{2} |S, S_z\rangle$$

where  $|S, S_z\rangle = \chi_{S, S_z}(\sigma_1, \sigma_2)$  have the explicit form:

$$\begin{aligned} \chi_{1,1} &= v_+(\sigma_1)v_+(\sigma_2) \\ \chi_{1,0} &= \frac{1}{\sqrt{2}} [v_+(\sigma_1)v_-(\sigma_2) + v_-(\sigma_1)v_+(\sigma_2)] \\ \chi_{1,-1} &= v_-(\sigma_1)v_-(\sigma_2) \end{aligned} \left. \vphantom{\begin{aligned} \chi_{1,1} \\ \chi_{1,0} \\ \chi_{1,-1} \end{aligned}} \right) \text{Spin triplet: symmetric}$$
$$\chi_{0,0} = \frac{1}{\sqrt{2}} [v_+(\sigma_1)v_-(\sigma_2) - v_-(\sigma_1)v_+(\sigma_2)] \quad \text{Spin singlet: antisymmetric}$$



# 2 interacting spins

## Energy

$$\begin{aligned}\langle \Psi_{+,-} | \mathcal{H} | \Psi_{+,-} \rangle &= \langle \Psi_{+,-}^{orb} | \mathcal{H} | \Psi_{+,-}^{orb} \rangle = \langle \phi | h | \phi \rangle + \langle \psi | h | \psi \rangle + J_{12} + (-1)^S K_{12} \\ &= E_0 + J_{12} + (-1)^S K_{12}\end{aligned}$$

with the Hartree and the exchange terms:

$$\begin{aligned}J_{12} &= \langle \phi(1)\psi(2) | V_{12} | \phi(1)\psi(2) \rangle \\ K_{12} &= \langle \phi(1)\psi(2) | V_{12} | \phi(2)\psi(1) \rangle\end{aligned}$$

The Pauli principle  $\Rightarrow$  the energy is affected by spin even if  $\mathcal{H}$  does not depend explicitly on it:

$$\begin{aligned}\langle S = 0 | \mathcal{H} | S = 0 \rangle &= E_s && \text{singlet} \\ \langle S = 1 | \mathcal{H} | S = 1 \rangle &= E_t && \text{triplet}\end{aligned}$$

Hence:

$$\begin{aligned}\langle S = 0 | \mathcal{H} | S = 0 \rangle - \langle S = 1 | \mathcal{H} | S = 1 \rangle &= E_s - E_t \\ &= 2K_{12}\end{aligned}$$

# 2 interacting spins

## Energy

The relative order of  $E_s$ ,  $E_t$  depends on  $V_{12}$  ( $K_{12}$ )

In He, with  $V_{12}$ = repulsive Coulomb interaction,

$$E_s > E_t \quad (K > 0)$$

(a part from the ground state, when only singlet is allowed)

But present discussion is independent on the sign of  $E_s - E_t$

# 2 interacting spins

**Idea: write a model hamiltonian explicitly dependent on spin (and only dependent on spin) which gives the same difference between the energy eigenvalues of its eigenstates as the original one.**

Consider the operator:

$$\Sigma_{12} = \vec{s}_1 \cdot \vec{s}_2 = \frac{1}{2} S^2 - \frac{3}{4}$$

which is diagonal on the coupled basis, with eigenvalues:

$$\langle S = 0 | \Sigma_{12} | S = 0 \rangle = -\frac{3}{4}, \quad \langle S = 1 | \Sigma_{12} | S = 1 \rangle = \frac{1}{4}$$

Consider then:  $\mathcal{H}^{spin} = -(E_s - E_t) \Sigma_{12}$

We have:

$$\langle S = 0 | \mathcal{H}^{spin} | S = 0 \rangle - \langle S = 1 | \mathcal{H}^{spin} | S = 1 \rangle = E_s - E_t$$

# Heisenberg hamiltonian

$$\mathcal{H}^{spin} = -(E_s - E_t)\Sigma_{12}$$

is therefore OK! Defining:  $J \equiv E_s - E_t$  , we have  
(going back to the individual basis representation):

$$\mathcal{H}^{spin} = -J\vec{s}_1 \cdot \vec{s}_2$$

$J > 0$  ( $E_s > E_t$ )  $\uparrow\uparrow$  spins favored  $\Rightarrow$  ferromagnetic case

$J < 0$  ( $E_s < E_t$ )  $\uparrow\downarrow$  spins favored  $\Rightarrow$  antiferromagnetic case

# Heisenberg hamiltonian

Extension to the case of several spins:

$$\mathcal{H}^{spin} = - \sum_{\substack{i,j=1 \\ i \neq j}}^N J_{ij} \vec{s}_i \cdot \vec{s}_j$$

(see also: Ashcroft & Mermin, chap. 32,  
“Spin hamiltonian and Heisenberg model”)


# Ising model

Consider only the possibility :  $s_i = \pm 1$  and **nearest neighbor interaction only**, with the same interaction constant  $J$



$$\mathcal{H}^{spin} = -J \sum_{i,j=1}^N s_i s_j$$

$J > 0$  ( $E_s > E_t$ )  $\uparrow\uparrow$  spins favored  $\Rightarrow$  **ferromagnetic case**


$J < 0$  ( $E_s < E_t$ )  $\uparrow\downarrow$  spins favored  $\Rightarrow$  **antiferromagnetic case**



$$E = -J$$



$$E = +J$$




(choosing the kind of interaction, we specify the **energy model**)

# Ising model on a lattice

lattice containing  $N$  sites

lattice site  $i$  has associated with it a number  $s_i$ , where  $s_i = +1$  for an “up” ( $\uparrow$ ) spin and  $s_i = -1$  for a “down” ( $\downarrow$ ) spin. A particular configuration or **microstate** of the lattice is specified by the set of variables  $\{s_1, s_2, \dots, s_N\}$  for all lattice sites.



Lowest energy state of the 2D Ising model on a square lattice with ferromagnetic ( $J > 0$ ) and antiferromagnetic ( $J < 0$ ) interactions. Solid and open circles correspond to  $+1$  and  $-1$  spins, respectively.

# Ising model: interesting quantities



# Ising model: energy

lattice containing  $N$  sites  
No external magnetic field:

$$E = -J \sum_{i,j=\text{nn}(i)}^N s_i s_j$$

(nn=nearest neighbor)

Energy in presence of an external magnetic field:

$$E = -J \sum_{i,j=\text{nn}(i)}^N s_i s_j - H \sum_{i=1}^N s_i,$$

or, better, define an **average energy per spin**:  $E/N$

# Ising model: magnetization

$$M = \sum_{i=1}^N s_i$$

**“Order parameter”**: total magnetization, or -better- average magnetization per spin:

$$m = \frac{M}{N} = \frac{1}{N} \sum_{i=1}^N s_i$$

$$-1 \leq m \leq +1$$

# Ising model: configurations and energy

$2^n$  different configurations for  $n$  spins.  
(microstates)

e.g.  $2^4 = 16$  spin configurations for  $2 \times 2$  lattice

+	+	-	-		+	-	-	+
+	+	-	-		-	+	+	-

-	+	+	-	+	+	+	+	+	-	+	-	-	-	-	-	-	+	-	+	+	-	+	
+	+	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	+

---

For  $J > 0$  the state of lowest energy is when all the spins are aligned.  
The state has macroscopic magnetization (**ferromagnetic**).

The **ground state energy** per spin

(ferromagnetic case, thermodynamic limit ( $N$  large), no ext. field) is:


$$E_0/N = - 2J$$

# Ising model: dynamics?

Beside an **energy model**, we must define a **dynamics** in order to simulate the evolution of the system (i.e. to generate the trajectory in the phase space, to generate the configurations of the Markov chain)

# Ising model: spin flip dynamics


Consider  $nn$  interactions, choose a random spin and flip it: it's **a new configuration** (a microstate)



Apply **Metropolis Monte Carlo** method for evolution in the **canonical ensemble** (fix  $T$ ). Evolution is driven by the **energy change** between the old and the new configuration,  $\Delta E$ .

**Remark:** Is it sufficient to calculate only  $\Delta E$ , not  $E$  at each new configuration!

# Ising model: spin flip dynamics



The five possible transitions of the Ising model on the square lattice with spin flip

# Ising model: boundary conditions

Of course we cannot simulate an infinite system (the thermodynamic limit).

We have two choices for the simulation cell:

- free (open) boundary conditions
- periodic boundary conditions (PBC)

# Ising model: free boundary conditions

in a  $N=L \times L$  spin lattice there are  $2L(L-1)$  nn interactions;  
for the ferromagnetic g.s. configuration, for instance, the energy is:


			+	+	+	+
		+	+	+	+	+
+	+	+	+	+	+	+
+	+	+	+	+	+	+

Let's count the interactions...



# Ising model: free boundary conditions

in a  $N=L \times L$  spin lattice there are  $2L(L-1)$  nn interactions;  
for the ferromagnetic g.s. configuration, for instance, the energy is:



$E_0/N = -J$      
  $E_0/N = -(12/9)J$      
  $E_0/N = -(24/16)J$     ...   
  $E_0/N = -J \times 2L(L-1)/L^2$   
 $= -2J \times (1 - 1/L)$


(volume term)
(surface term)

Energy per spin in the ground state converges to the value

$E_0/N = -2J$  in the thermodynamic limit

(with deviations  $\sim 1/L$ )

# Ising model: PBC



The energy is a  $2N$ -term sum:  
each spin interacts with its NN  
within the simulation cell or  
with the NN images


One of the  $2^N$  possible configurations of a system of  $N = 16$  Ising spins on a square lattice.  
with periodic boundary conditions.

# Ising model: PBC

We have always:

$2^4 = 16$  spin configurations for  $2 \times 2$  lattice

but the energy for each configuration in case of free boundary conditions and PBC is different:



Two different configurations with 2 spins up

# of spins UP	Degeneracy	Energy	Magnetisation
4	1	$-8J$	4
3	4	0	2
2	4	0	0
2	2	$8J$	0
1	4	0	-2
0	1	$-8J$	-4

Energy and magnetization of 16 configurations of the  $2 \times 2$  Ising model with PBC

Energy per spin in the ground state is always equal to the value  $E_0/N = -2J$  in the thermodynamic limit


# Ising model: phase transition

**Low T: spin configuration minimizes energy**

(if  $J > 0$ : spins tend to align => **high (absolute) magnetization**)

**High T: spin configuration maximizes entropy**

(=**disorder**) (spins tend to disalign => **low magnetization**)




**existence of a  
Critical temperature  $T_c$**

in 2D the model has an  
analytical solution:


$$T_c = 2.269 J/k_B$$

# Apply **Metropolis Monte Carlo** method for evolution in the **canonical ensemble** (fix $T$ ):

Average magnetization per spin as a function of time for a lattice  $10 \times 10$  and two different  $T$




Fluctuations! do, as usual, **temporal averages**:  $\langle M \rangle / N$ ,  $\langle E \rangle / N$



Magnetisation as a function of the temperature for the 2D Ising model.  
 (data averaged on time; red lines indicate the  $T$  values in the previous plot)


Magnetization distribution for  $T < T_c$  (solid) and  $T > T_c$  (dashed curve)



(data collected during time evolution, at equilibrium)

# Ising model: phase transition

$T_c$  also for energy, not only for magnetization:



and also the energy fluctuates during time evolution...



# Ising model: fluctuations

Fluctuations are intrinsic to the system evolution  
and are important!

Linear response functions are related to  
equilibrium fluctuations:

(already proved):  $C = \frac{\partial \langle E \rangle}{\partial T}$  ,  $C = \frac{1}{kT^2} (\langle E^2 \rangle - \langle E \rangle^2)$

but also:  $\chi = \lim_{H \rightarrow 0} \frac{\partial \langle M \rangle}{\partial H}$  ,  $\chi = \frac{1}{kT} (\langle M^2 \rangle - \langle M \rangle^2)$

where  $\langle M \rangle$  and  $\langle M^2 \rangle$  are evaluated in zero magnetic fields.

# Ising model:

## fluctuations and phase transition

Rapid change in  $\langle E \rangle$  and  $\langle M \rangle \Rightarrow$  singularities in  $C$  and  $\chi$


(Large fluctuations near the phase transition:  
Second Order phase transition)

specific heat:

$$C = \frac{\partial \langle E \rangle}{\partial T}$$

magnetic susceptibility:

$$\chi = \lim_{H \rightarrow 0} \frac{\partial \langle M \rangle}{\partial H}$$



# Implementing the Ising model in the code

# Implementing the Ising model

on a 2D square lattice in the canonical ensemble  
zero-field, nearest neighbor interactions only

$$\mathcal{H}^{spin} = -J \sum_{i,j=1}^N s_i s_j \quad s_i = \pm 1$$

Input parameters are:

- $L$  (linear lattice dimension, which gives the number of spins:  $N=L*L$ )
- $nmcs$  (number of total MC steps per spin)
- $nequil$  (number of equilibration MC steps per spin)
- $T$  (temperature of the thermal bath).

# Implementing the Ising model

```
program ising
  ! metropolis algorithm for the ising model on a square lattice
  use common
  integer :: imcs,ispin,jspin
  real (kind = double), dimension(5) :: cum
  call initial(nequil,cum)
  ! equilibrate system
  do imcs = 1,nequil
    call metropolis()
  end do
  ! accumulate data while updating spins
  do imcs = 1,nmcs
    call metropolis()
    call data(cum)
  end do
  call output(cum)
  .....
  .....
end program ising
```

# Ising model on a lattice

L : linear lattice dimension

N = LxL : number of spins

**a configuration** (a microstate) is the whole sequence of spins, i.e. the LxL array spin(x,y)

```
module common
```

```
.....
```

```
integer, public, dimension(:, :), allocatable :: spin
```

```
subroutine initial(nequil, cum)
```

```
.....
```

```
allocate(spin(L,L))
```

```
.....
```

```
spin(x,y) = 1
```

```
else
```

```
spin(x,y) = -1
```

```
.....
```

$$S_i = \pm 1$$

# Ising model: magnetization

Total magnetization, or define an average magnetization per spin:

$$M = \sum_{i=1}^N s_i$$

$$m = \frac{M}{N} = \frac{1}{N} \sum_{i=1}^N s_i$$

$$-1 \leq m \leq +1$$

```
! compute initial magnetization
M = 0.0_double
do y = 1,L
  do x = 1,L
    .....
    .....
    M = M + spin(x,y)
  end do
end do
```

(Instead of the loop over x,y, write: `M=sum( spin )` )

# Ising model: energy

$$E = -J \sum_{i,j=\text{nn}(i)}^N s_i s_j$$

```
! compute initial energy
```

```
E = 0.0_double
```

```
do y = 1,L
```

```
.....
```

```
  do x = 1,L
```

```
    .....
```

```
    sums = spin(x,up) + spin(right,y)
```

```
! calculate the initial energy summing all over pairs
```

```
! (for a given spin, consider only the up NN and the right NN
```

```
! - NOT the down and the left NN - : each interaction is counted once
```

```
  E = E - spin(x,y)*sums
```


```
  end do
```

```
end do
```



# Ising model: energy with PBC

```
do y = 1,L
  ! periodic boundary conditions
  if (y == L) then
    up = 1
  else
    up = y + 1
  end if
  do x = 1,L
    if (x == L) then
      right = 1
    else
      right = x + 1
    end if
    sums = spin(x,up) + spin(right,y)
  ! calculate the initial energy summing all over pairs
  ! (for a given spin, consider only the up NN and the right NN
  ! - NOT the down and the left NN - : each interaction is counted once
    E = E - spin(x,y)*sums
  end do
end do
```



# Ising model: spin flip dynamics

Choose a random spin and flip it:  
it's **a new configuration** (a microstate)

```
do ispin = 1,N
  !      random x and y coordinates for trial spin
  call random_number(rnd)
  x = int(L*rnd) + 1      ←  $1 \leq x \leq L$ 
  call random_number(rnd)
  y = int(L*rnd) + 1      ←  $1 \leq y \leq L$ 
  .....
```

Flip is:  $\text{spin}(x,y) = -\text{spin}(x,y)$

but do it later, only if you decide to accept the flip (according to Metropolis)

# Ising model: energy variations per spin flip

Evolution is driven by the **energy change** between the old and the new configuration (Metropolis MC)

```
dE = DeltaE(x,y) ← energy variation for spin(x,y) flip
call random_number(rnd)
if (rnd <= w(dE)) then ← w(dE) is  $e^{-\Delta E/k_B T}$ 
    spin(x,y) = -spin(x,y)
    accept = accept + 1
    .....
```


```
function DeltaE(x,y) result (DeltaE_result)
.....
DeltaE_result = 2*spin(x,y)*(left + right + up + down)
.....
```

# Energy variations per spin flip with PBC

```

function DeltaE(x,y) result (DeltaE_result)
  ! periodic boundary conditions
  .....
  if (x == 1) then
    left = spin(L,y)
    right = spin(2,y)
  else if (x == L) then
    left = spin(L-1,y)
    right = spin(1,y)
  else
    left = spin(x-1,y)
    right = spin(x+1,y)
  end if
  if (y == 1) then
    up = spin(x,2)
    down = spin(x,L)
  else if (y == L) then
    up = spin(x,1)
    down = spin(x,L-1)
  else
    up = spin(x,y+1)
    down = spin(x,y-1)
  end if
  DeltaE_result = 2*spin(x,y)*(left + right + up + down)
  .....

```




# Ising model: storage of Boltzmann's coeff.

```
! Choosing the interaction parameter J=1,  
! possible energy variations per spin flip are -8,-4,0,+4,+8:
```

```
do dE = -8,8,4  
  w(dE) = exp(-dE/T)  
end do  
accept = 0  
do i = 1,5  
  cum(i) = 0.0_double  
end do
```

Convenient to store the Boltzmann's coefficient for these discrete values of energy variations



The five possible transitions of the Ising model on the square lattice with spin flip

# Ising model: updating energy and magnetization

```
subroutine metropolis()  
  ! one Monte Carlo step per spin  
  .....  
  do ispin = 1,N  
    .....  
    dE = DeltaE(x,y)  
    call random_number(rnd)  
    if (rnd <= w(dE)) then  
      spin(x,y) = -spin(x,y)  
      accept = accept + 1  
      M = M + 2*spin(x,y) ! factor 2 is to account for the variation:  
      E = E + dE          ! (-(-)+(+))  
    end if  
  end do  
end subroutine metropolis
```

**DO NOT CALCULATE  
EVERYTHING FROM THE  
SCRATCH!!**

$\Delta E$  is already a variation

# Spin flip dynamics: how to choose spin to flip?

## Random ...

```
do ispin = 1,N
  !      random x and y coordinates for trial spin
  call random_number(rnd)
  x = int(L*rnd) + 1      ←  $1 \leq x \leq L$ 
  call random_number(rnd)
  y = int(L*rnd) + 1      ←  $1 \leq y \leq L$ 
  .....
```

## or ordered (sequential) ...

```
do x = 1,L
do y = 1,L
  ...
```

$$\text{spin}(x,y) = -\text{spin}(x,y)$$

# Spin flip dynamics: how to choose spin to flip?

- ORDERED: in some cases, it could go more slowly towards equilibrium (see later: correlation time), but it depends...
- NO appreciable differences in the statistics at equilibrium



# Measuring physical quantities: how to accumulate data?

```
subroutine data(cum)
  ! accumulate data after every Monte Carlo step per spin
  real (kind = double), dimension(5), intent (inout) :: cum
  cum(1) = cum(1) + E
  cum(2) = cum(2) + E*E
  cum(3) = cum(3) + M
  cum(4) = cum(4) + M*M
  cum(5) = cum(5) + abs(M)
end subroutine data
```

After one MC step per spin for all spins:

```
do imcs = 1, nmcs
  call metropolis()           contains the loop over all the spins
  call data(cum)
end do
```

Alternatively, do it after each MC step per individual spin...

# Measuring physical quantities: how to accumulate data?

## Further remarks...

- Use statistically **INDEPENDENT** configurations
- Calculate therefore the **CORRELATION TIME** by considering the **autocorrelation functions**:


$$C_M(t) = \langle M(t)M(0) \rangle - \langle M \rangle^2, \quad C_E(t) = \langle E(t)E(0) \rangle - \langle E \rangle^2$$

$$(C_M(0) \propto \chi, \quad C_E(0) \propto C_V)$$

$C_M(t) \rightarrow 0$  and  $C_E(t) \rightarrow 0$  exponentially for  $t \rightarrow \infty$

with a certain decay time  $\tau$ : consider intervals longer than  $\tau$  for statistical averages

# Autocorrelation functions



(NOTE: "critical slowing down" for  $T \rightarrow T_C$ )

=> configurations change very slowly, and it is difficult to sample enough configurations

# Measuring physical quantities: how to accumulate data?

## Further remarks...

- see also CORRELATION LENGTH between magnetic domains,  $\zeta(T)$
- close to  $T_c$ , also the correlation length increases (spin alignments are more correlated), up to divergence

# Measuring physical quantities: which errors?

- necessary to give the ERROR ESTIMATE corresponding to the measured physical quantity !!!
- (see Tab. I of D.P. Landau, PRB 13, 2997 (1976), “Finite size behavior of the Ising square lattice”)
- do also BLOCKING (called “coarse grained technique” in that paper)


# How to do efficiently simulations as a function of $T$ ?

- Sometimes EQUILIBRATION time is long...
- IDEA: for  $T'$  close to  $T$ , choose as starting point the equilibrated output of  $T$


# How to do efficiently simulations as a function of $T$ ?

*Magnetization (System : 10\*10)*

*Thick :  $T = 0.9 T_c$  Dashed :  $T = 2 T_c$*




*Magnetization distribution*



# Ising model: size problems

We cannot simulate an INFINITE system!



The temperature dependence of the specific heat  $C$  (per spin) of the Ising model



# Ising model: size problems

## INTERFACE EFFECTS:

example of energy for HALF UP/HALF DOWN configurations:

$$L=2 \quad E=0$$

$$L=4 \quad E=-1$$

$$L=8 \quad E=-1.5$$


$$L=16 \quad E=-1.75$$

$$L=20 \quad E=-1.8$$

$$L=32 \quad E=-1.875$$

... for an infinite system:  $E=-2$

We have a (“interface”) term proportional to  $1/L$



“interface”

# Ising model: alternative dynamics

- in the **SPIN FLIP dynamics** the order parameter is not conserved (**M changes** during evolution)
- alternative: **NN spin exchange (Kawasaki dyn.)**  
(exchange two NN spins picked at random;  
**M is conserved**; this is equivalent to **LATTICE GAS MODELS** with fixed number of particles)

# Ising model: Kawasaki dynamics

Fixed magnetization : change of thermodynamical ensemble

No modification of the equilibrium properties  
except phase separation



# Addition of further interactions

- Attractive ( $J>0$ ) nearest-neighbor (NN) interaction only:  
total energy of the system:

$$E = -\frac{J}{2} \sum_{\langle ij \rangle} n_i n_j$$


=> Trend to aggregation (diffusive behavior is limited to a transient)

- Add a repulsive ( $J<0$ ) next-nearest-neighbor (NNN) interaction: total energy of the system:

$$E = -\frac{1}{2} \sum_{\langle ij \rangle} J_{ij} n_i n_j$$

=> The behavior depends on the ratio  $R = J_{NNN}/J_{NN}$

- With finite NN and/or NNN interactions, temperature plays a role




**T=10, starting from random configuration**


# Ising model: other generalizations

- SPINS: XY, Heisenberg, Potts...
- LATTICES: Square, Triangle, Cubic, Honeycomb, Kagome....
- INTERACTIONS: Magn. Field, Antiferrom., Next Nearest Neighbor (NNN)....

Different behaviour according to the geometry and the kind of interactions.  
Example: frustration in the triangular antiferromagnetic Ising model:



FM



AFM




FIG. 15. (Color online) Behavior of the two dimensional Ising model (top) and XY model (bottom) at  $T < T_c$ ,  $T = T_c$  and  $T > T_c$ . For the Ising model, an  $\uparrow$  spin ( $s_i = 1$ ) is represented by red and a  $\downarrow$  spin ( $s_i = -1$ ) is represented by blue. For the XY model, because every spin is determined by  $\theta$ , so  $\mathbf{s}_i = (\cos \theta, \sin \theta)$ , an angle of  $2\pi$  is represented by red and an angle of 0 by blue. In the low-temperature phase, the Ising model exhibits spontaneous magnetization while in the XY model, vortex buddies appear (characterized by points where a continuum from blue to red, or viceversa, circle the point. It is worth to notice that these points are present by pairs with opposite circulation).

# Program:

On moodle2:

**ising.f90**



# Exercise

(a) Choose  $L=30, T=2$ , and initially  $\text{spin}=\pm 1$  randomly. Calculating and plotting the instantaneous values of the energy  $E/N$  and the magnetization  $M/N$  per spin (averaged over the lattice) as a function of Metropolis-MC steps, how much time (i.e. how many *nequil* MC steps) is it necessary to equilibrate the system?


Hint:

- Since initially  $\text{spin}=\pm 1$  randomly,  $E/N$  and  $M/N$  initially will be far from the expected equilibrium average value.

First, set *nequil*=0 and plot instantaneous values of  $E/N$  and  $M/N$ .

Estimate *nequil* from that plot!!! Visualization is important!!!!

*nequil* of course depends on  $T$  and on the initial situation



Then, set *nequil* not zero and calculate the time average  $\langle E \rangle/N$  and  $\langle M \rangle/N$ ; increasing the total *nmcs*, the two quantities should converge...

# Exercise


(a) Choose  $L=30, T=2$ , and initially  $\text{spin}=\pm 1$  randomly. ...

Plot a snapshot of the spin pattern: does the system appear ordered or disordered?

it should appear **ordered**...

**p 'ising-up.dat' ps 3 pt 7, 'ising-down.dat' ps 3 pt 7**

Plotting “ising-up.dat” and “ising-down.dat” which contain the coordinates of spin up and down respectively, one should get something like that:




Calculate also  $c$  and  $\chi$ .

# Exercise

(a) Choose  $L=30, T=2$ , and initially  $\text{spin}=\pm I$  randomly. ...


Calculate also  $c$  and  $\chi$ .

(b) Choose  $T=I$  and repeat (a)...




## Raw data: traces, covariance and autocorrelation time


Trace: magnetization for  $T = 2.27J/k_B \approx T_C$  ( $10^5$  sweeps)




# Magnetization ( $10^5$ sweeps)




# Energy ( $10^5$ sweeps)



# Magnetic susceptibility ( $10^5$ sweeps)




# Magnetic susceptibility near $T_c$ ( $10^6$ sweeps)






## Specific heat ( $10^5$ sweeps)



## Specific heat near $T_c$ ( $10^6$ sweeps)



A few references:

Ising E 1925 Zeitschrift fur Physik 31 253–258

Bethe H 1931 Zeitschrift fur Physik 71 205

Heisenberg W 1928 Zeitschrift fur Physik 49 205

# Universality and critical exponents

Near  $T_c$ , we can characterize the behavior of many physical quantities by power law behavior just as we characterized the percolation threshold (see Table 13.1). For example, we can write  $m$  near  $T_c$  as

$$m(T) \sim (T_c - T)^\beta, \quad (17.22)$$

where  $\beta$  is a critical exponent (not to be confused with the inverse temperature). Various thermodynamic derivatives such as the susceptibility and heat capacity diverge at  $T_c$ . We write

$$\chi \sim |T - T_c|^{-\gamma} \quad (17.23)$$

and

$$C \sim |T - T_c|^{-\alpha}. \quad (17.24)$$

We have assumed that  $\chi$  and  $C$  are characterized by the same critical exponents above and below  $T_c$ .

Another measure of the magnetic fluctuations is the linear dimension  $\xi(T)$  of a typical magnetic domain. We expect the *correlation length*  $\xi(T)$  to be the order of a lattice spacing for  $T \gg T_c$ . Because the alignment of the spins becomes more correlated as  $T$  approaches  $T_c$  from above,  $\xi(T)$  increases as  $T$  approaches  $T_c$ . We can characterize the divergent behavior of  $\xi(T)$  near  $T_c$  by the critical exponent  $\nu$ :

$$\xi(T) \sim |T - T_c|^{-\nu}. \quad (17.25)$$

# Universality and critical exponents

More precisely, the magnetisation follow a power law close to the transition only approaching from smaller T:

$$M = 0 \quad T \geq T_c$$

$$M \sim |1 - T/T_c|^\beta \quad T < T_c$$

If we use the Reduced temperature :  $\Delta T = (T - T_c)/T_c$

$$C \sim |\Delta T|^{-\alpha}$$

$$M \sim |\Delta T|^\beta \quad \text{for } \Delta T < 0$$

$$\chi \sim |\Delta T|^{-\gamma}$$

$$\xi \sim |\Delta T|^{-\nu}$$

# Universality and critical exponents

the critical exponents are not independent from each other, because of the following scaling laws (●):

$$\begin{aligned}\gamma &= \nu(2 - \eta), \\ 2 &= \alpha + 2\beta + \gamma, \\ \nu d &= 2 - \alpha, \\ \gamma &= \beta(\delta - 1),\end{aligned}$$

so it is only necessary to know two of them to determine the others.

For the 2D Ising model:

$\alpha$		0
$\beta$		0.125
$\gamma$		1.750
$\nu$		1

- Kerson Huang, Introduction to Statistical Physics (CRC Press)

# Universality and critical exponents

If the heat capacity goes like:  $C(T) \propto |T - T_c|^{-\alpha}$

we could plot  $\frac{1}{C(T)}$  as a function of  $T$  and make a linear fit:

$$\frac{1}{C(T)} = a + bT$$

and similarly for the other quantities.

**BUT:** Because we can simulate only finite lattices, a direct fit of the measured quantities does not yield good estimates for the corresponding exponents  $\alpha$ ,  $\nu$ ,  $\beta$ , and  $\gamma \Rightarrow$  we have to take into account the finite size of the system

**$\Rightarrow$  finite size scaling**

# Universality and critical exponents

The shift in the peak position of  $C$  and  $\chi$  with respect to the critical temperature corresponding to the thermodynamic limit is described by:  $T_c(L) - T_c(\infty) \propto L^{-\lambda}$




FIG. 1. Typical behaviour of a physical quantity  $A$  vs temperature close to the critical point for various system sizes. Figure taken from Thijsen<sup>8</sup>.

[Thijsen, Computational Physics (Cambridge University Press)]




FIG. 9. (Color online) Inverse of the critical temperature  $T_c^{-1}$  vs inverse of lattice size  $L^{-1}$  for the 2D Ising model. The curve was fitted with a power law  $T_c^{-1} = T_c^{-1} - bL^{-1/\nu}$ , and the critical exponent  $\nu$  was determined.

- From: E. Ibarra-Garcia-Padilla et al., European Journal of Physics 37(6):065103 DOI: [10.1088/0143-0807/37/6/065103](https://doi.org/10.1088/0143-0807/37/6/065103)



# Universality and critical exponents

(This is referred to percolation)

Because we can simulate only finite lattices, it is difficult to obtain estimates for the critical exponents  $\alpha$ ,  $\beta$ , and  $\gamma$  by using the definitions (17.22)–(17.24) directly. We learned in Section 13.4, we can do a *finite size scaling analysis* to extrapolate finite  $L$  results to  $L \rightarrow \infty$ . For example, from Fig. 17.2 we see that the temperature at which  $C$  exhibits a maximum becomes better defined for larger lattices. This behavior provides a simple definition of the transition temperature  $T_c(L)$  for a finite system. According to finite size scaling theory,  $T_c(L)$  scales as

$$T_c(L) - T_c(L = \infty) \sim aL^{-1/\nu}, \quad (17.27)$$

where  $a$  is a constant and  $\nu$  is defined in (17.25). The finite size of the lattice is important when the correlation length

$$\xi(T) \sim L \sim |T - T_c|^{-\nu}. \quad (17.28)$$

As in Section 13.4, we can set  $T = T_c$  and consider the  $L$ -dependence of  $M$ ,  $C$ , and  $\chi$ :

$$m(T) \sim (T_c - T)^\beta \rightarrow L^{-\beta/\nu} \quad (17.29)$$

$$C(T) \sim |T - T_c|^{-\alpha} \rightarrow L^{\alpha/\nu} \quad (17.30)$$

$$\chi(T) \sim |T - T_c|^{-\gamma} \rightarrow L^{\gamma/\nu}. \quad (17.31)$$

## Scaled magnetization vs unscaled $T$ ( $10^5$ sweeps)

