

**Università di Trieste**  
**LAUREA MAGISTRALE IN GEOSCIENZE SM62**  
**Percorso Esplorazione Geologica**

**Anno accademico 2021 - 2022**

# **Geologia Marina 953SM**


## Parte I

**Modulo 1.2**     Introduzione ai fondali oceanici oceanici: Acqua oceanica, sedimento, e fluidi interstiziali

Docente  
**Angelo Camerlenghi**


# Oceans and Seas:

~ 70% of Earth Surface




# Oceans and Seas:

## Global sinks of sediments eroded on continents





## Total Sediment Thickness of the World's Oceans & Marginal Seas




A digital total sediment thickness database for the world's oceans and marginal seas is being compiled by the National Geophysical Data Center (NGDC) Marine Geology & Geophysics Division. The data are gridded with a spacing of 5 arc-minutes by 5 arc-minutes. Sediment thickness data were compiled from three principle sources: previously published isopach maps; ocean drilling results, both ODP and DSDP; and seismic reflection profiles archived at NGDC as well as seismic data and isopach maps available as part of the IOC's Geological/Geophysical Atlas of the Pacific (GAPAP) project.

The distribution of sediments in the oceans is controlled by five primary factors:

- 1) Age of the underlying crust
- 2) Tectonic history of the ocean crust
- 3) Structural trends in basement
- 4) Nature and location of sediment sources, and
- 5) The nature of the sedimentary processes delivering sediments to depocenters

The data values are in meters and represent the depth to acoustic basement. It should be noted that acoustic basement may not actually represent the base of the sediments. These data are intended to provide a minimum value for the thickness of the sediment in a particular geographic region.


<http://www.ngdc.noaa.gov/mgg/sedthick/sedthick.html>





## Terrigenous Sediments (also: *lithogenous*)

- Eroded rock fragments from land
- Transported from land by
  - Water (e.g., river-transported sediment)
  - Wind (e.g., windblown dust) - *aeolian transport*
  - Ice (e.g., ice-rafted rocks)
  - Gravity (e.g., turbidity currents)



## Terrigenous Sediments (also: *lithogenous*)

### CLAYS (hydrates aluminosilicate mineral)

- Kaolinite
- Chlorite
- Illite
- Montmorillonite
- Particles are generally  $< 2 \mu\text{m}$

Also generated by dissolution of calcareous plankton and benthos (red clays, or pelagic clays) below the **Carbonate Compensation Depth (CCD)** in the open ocean.

## Terrigenous Sediments (also: *lithogenous*)

SILTS and SANDS

Silts (2 - 63  $\mu\text{m}$ )

Sands (>63  $\mu\text{m}$ )

Mainly transported by turbidity currents, debris flows, icebergs

Reflect composition of surrounding land masses



## Biogenic Sediments (also: *biogenous*)

Hard remains of dead organisms


- Macroscopic (large remains)
  - Shells, bones, teeth
- Microscopic (small remains)
  - Tiny shells or tests settle through water column
  - Biogenic ooze (30% or more tests)
  - Mainly algae and protozoans
- Calcium carbonate ( $\text{CaCO}_3$ )
- Silica ( $\text{SiO}_2$  or  $\text{SiO}_2 \cdot n\text{H}_2\text{O}$ )

# Biogenic Sediments (also: *biogenous*) continued

## Calcium carbonate in biogenous sediments

### Coccolithophores (algae)

- Photosynthetic
- Coccoliths (nano-plankton)



(a)

Copyright © 2008 Pearson Prentice Hall, Inc.

# Biogenic Sediments (also: *biogenous*) continued


## Calcium carbonate in biogenous sediments

### Foraminifera

(Benthic and Planktonic  
Protozoans)

Calcite

30 $\mu$ m - 1 mm



(c)

Copyright © 2008 Pearson Prentice Hall, Inc.



## Biogenic Sediments (also: *biogenous*) continued

### Pteropods

(planktonic gastropods molluscs)

### Aragonite

(a variety of calcite, more soluble)



# Biogenic Sediments (also: *biogenous*) continued

## Silica in biogenic sediments


Diatoms (algae)  $< 200 \mu\text{m}$

Radiolarians (protozoans)  $50\text{-}300 \mu\text{m}$



(b)

Copyright © 2008 Pearson Prentice Hall, Inc.



(a)

Copyright © 2008 Pearson Prentice Hall, Inc.

## Authigenic Sediments (also: *Hydrogenous*)

- Minerals precipitate directly from seawater
  - Manganese nodules
  - Phosphates (beneath areas in surface ocean high biological productivity)
  - Carbonates (Aragonite and calcite)
  - Metal sulfides (Associated with hydrothermal)
  - Evaporites (Minerals that form when seawater evaporates)
- Small proportion of marine sediments




(a)

Copyright © 2008 Pearson Prentice Hall, Inc.



## Volcanogenic Sediments (also: *Hydrogenous*)

- Ash layers
- Lava basalts
- Tephra layers



Glass shards

## Cosmogenous Sediments

- Macroscopic meteor debris
- Microscopic iron-nickel and silicate spherules
- Tektites
- Space dust
- Overall, insignificant proportion of marine sediments

# Microplastics


- “New and emerging issue” — GESAMP (Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection) is an advisory body, established in 1969, that advises the United Nations (UN) system on the scientific aspects of marine environmental protection.
- Plastic particles < 5mm
- Primary microplastics are deliberately manufactured
- Secondary microplastics are break-down products of larger debris
- Most plastic derives from land-based sources
  - Household and industrial waste + wastewater
  - Fishing, aquaculture, shipping, tourism, etc.





Front. Earth Sci., 30 April 2019 | <https://doi.org/10.3389/feart.2019.00080>

# Dispersion, Accumulation, and the Ultimate Fate of Microplastics in Deep-Marine Environments: A Review and Future Directions



# MARINE SEDIMENTS


## Distribution

### Neritic

- Shallow water deposits
- Close to land
- Dominantly lithogenous
- Typically deposited quickly

### Pelagic (Also Oceanic)

- Deeper water deposits
- Finer-grained sediments
- Deposited slowly



**Neritic**

**Oceanic**

**Abyssal clay**

**Calcareous Ooze**

**Siliceous Ooze**



## Distribution of pelagic sediment

Dominant component	Composition	Atlantic	Pacific	Indian	Total %
Foraminiferal and nannofossil ooze	Calcium carbonate	65	36	54	47
Pteropod ooze	Calcium carbonate	2	0.1	-	0.5
Diatom ooze	Silica (opal)	7	10	20	12
Radiolarian ooze	Silica (opal)	-	5	0.5	3
Red (actually brown) clay K, Fe	Al silicate	26	49	25	38

*Source P.Pinet Invitation to Oceanography, 2000 2nd Edition, Jones and Barlett Publishers, Massachusetts*

## Take-home messages

Marine sediments are very heterogeneous material, whose solid component is made by a mixture of particles with different composition, shape and size

All of the inorganic solid components of marine sediments undergo some kind of chemical alteration through time essentially in consequence of the exposure to salt water






# SEAWATER

Water has unique and unusual properties both in pure form and as a solvent.

These properties influence many of the chemical reactions taking place in the oceans.

“Distorted” tetrahedral arrangement, with 2 electrons in each lobe. Two lobes are used for O-H bonds and two lobes have free lone pairs of electrons.

The H-O-H tetrahedral angle of  $105^\circ$ , results in dipole moments, which means that this is a polar molecule.



Propensity to form hydrogen bonds. “Cooperative Bonding” in which the water molecules link together to form regions with structure.

## SEAWATER

- SALINITY: dissolved ion content by weight.
- Average 3.5 % or 35 parts per thousands
- Varying between 3.1% and 3.8%
- Not uniform distribution (horizontal and vertical)

## Seawater composition (by mass)

Element	%	
<u>Oxygen</u>	85.84	
<u>Hydrogen</u>	10.82	
<u>Chlorine</u>	<u>1.94</u>	because salinity is directly proportional to the amount of chlorine in sea water, and because chlorine can be measured accurately by a simple chemical analysis, <b>salinity S is defined using chlorinity</b>
<u>Sodium</u>	1.08	
<u>Magnesium</u>	0.1292	
<u>Sulfur</u>	0.091	
<u>Calcium</u>	0.04	
<u>Potassium</u>	0.04	<b><math>S (‰) = 1.80655 \times Cl (‰)</math></b>
<u>Bromine</u>	0.0067	
<u>Carbon</u>	0.0028	

**Now Salinity is measured by Electroconductivity in Practical Salinity Units (PSU)**

## Seawater composition

### Marcet's Principle (constancy of composition):


$$X/Cl = \text{constant}$$

- True for **conservative** elements, which include most of the major ions in seawater ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{SO}_4^{-2}$ , and  $\text{Cl}^-$ ).
- The concentration of these elements, normalized to salinity, is constant with depth and in the different oceans.
- The ratio of one conservative element to another will also be constant.



# Seawater composition

→ Conservative elements



# What does and what does not affect the constancy of composition of salt in seawater?

## Does Not:

- Precipitation – evaporation
- Freezing – thawing
- Turbulent mixing between water masses


## Does:

- Marginal seas receiving significant river runoff, e.g. Baltic Sea
- Anoxic basins where sulfate reduction occurs, e.g. Black Sea
- Shallow water environments where significant inorganic aragonite (a form of  $\text{CaCO}_3$ ) precipitation occurs – oolites
- Hydrothermal vents where seawater influx through hydrogeological processes causes low and high temperature alteration of basalts
- Evaporite basins, e.g. Dead Sea
- Interstitial waters of sediments

# Origin of dissolved ions

rain water does not look like river water

seawater doesn't look like river water



## Origin of dissolved ions

IGNEOUS ROCK + RAIN WATER >>>> SEDIMENTARY ROCK + RIVER WATER

Order of loss of cations during weathering of igneous rocks:



$\text{Ca}^{++} > \text{Na}^{+} > \text{Mg}^{++}$  are the most abundant cations in river water

$\text{K}^{+}$  is retained in clay minerals and is not abundant in river water





## Origin of dissolved ions

Product of alteration of igneous rocks:

LIMESTONES                      5 - 15%


SANDSTONES                    11 - 30%

**SHALES                            65 - 80%**

MINOR COMPONENTS (evaporites, cements...)

# Other input of ions to the oceans

## Groundwaters



Around the world groundwater discharge was observed at several coastal regions

Considerable flow rates of more than 1000 liters per minute were measured for submarine springs in the northeastern Gulf of Mexico.

Using the enrichment of  $^{226}\text{Ra}$ , Moore (1995) estimated that the groundwater flux to the coastal waters of South Atlantic Bay must be **~40% of the river-water flux (LOCAL)**.



## Hydrothermal Systems

- Seawater is entrained at spreading centers and contributes to hydrothermal circulation.
- Rate of hydrothermal circulation 0.3 to 3% of river input.
- Reactions between seawater and basalt modify composition of the circulating fluids.





FIGURE 19.2. Hydrothermal convection cells.

By **HIGH** and **LOW**  
temperature  
geochemistry


## Salt sinks

### Biological factors



Centric diatoms – an algae

Make a skeleton  
based on the element  
Si – ‘**biogenic silica**’  
or  $\text{SiO}_2$



*Emiliana huxleyi*, a coccolithorophorid

Make skeletal  
material from  
**calcium carbonate**  
 $\text{CaCO}_3$



## Salt sinks

Microbiology: Sulfate reduction in anoxic basins



Organic matter


Increase in alkalinity



Pyrite






Calcite



# Salt sinks

## Evaporation




The minerals precipitate out of solution in the reverse order of their solubilities, such that the order of precipitation from sea water is:

- 1 Calcite ( $\text{CaCO}_3$ ) and dolomite ( $\text{CaMg}(\text{CO}_3)_2$ )
- 2 Gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) and anhydrite ( $\text{CaSO}_4$ )
- 3 Halite (i.e. common salt,  $\text{NaCl}$ )
- 4 Potassium and magnesium salts


Limestone (calcite) and dolomite are more common than gypsum, which is more common than halite, which is more common than potassium and magnesium salts.

## *Salt sinks*

Extra-cellular polysaccharides may play a role in the formation of aggregates and transport of trace metals and pollutants in aquatic systems



[http://loer.tamug.tamu.edu/People/Santschi/santschi\\_research.htm](http://loer.tamug.tamu.edu/People/Santschi/santschi_research.htm)



= cation exchanging IN TO clay  
 = cation exchanging OUT OF clay  
 = clay minerals

## Salt sinks

### Ion Exchange on Sedimentary Particles

Ca introduced in rivers is removed from clays at the expense of Na and K uptake

**TABLE 12-4** CHANGE IN EXCHANGEABLE CATIONS WHEN RIVER-BORNE CLAYS ENTER SEAWATER<sup>a</sup> (after Sayles and Mangelsdorf, 1977).

	Average equivalent fraction <sup>b</sup>		Change in equiv. fraction	Net removal from ocean (g/yr) <sup>c</sup>	Percent of river input
	in river water	in seawater			
Na <sup>+</sup>	0.04	0.47	+0.43	$0.45 \times 10^{14}$	20.5 (30) <sup>d</sup>
K <sup>+</sup>	0.01	0.06	+0.05	$0.09 \times 10^{14}$	13
Ca <sup>2+</sup>	0.06	0.16	-0.44	$-0.4 \times 10^{14}$	-8
Mg <sup>2+</sup>	0.25	0.32	+0.07	$0.04 \times 10^{14}$	3
H <sup>+</sup>	0.10	0	-0.10		

<sup>a</sup> A + sign indicates uptake by the clay.

<sup>b</sup> Equivalent fraction is the fraction of the total exchange sites occupied by that cation.

<sup>c</sup> Assuming a suspended sediment input of  $183 \times 10^{14}$  g/yr and a CEC of 25 meq/100 g.

<sup>d</sup> Value in parentheses is corrected for cyclic salts.



## ***Salt sinks***


Hydrothermal systems

**Basalt alteration at 200 °C**

**Source for Ca, sink for Mg, SO<sub>4</sub>**



## Mass balance: Simple box models



From Berner and Berner. Box model appropriate for conservative elements in seawater.

## Physical properties of sea water

In oceanography, the calculation of water movement requires measurements of density with an accuracy of a few parts per million. This is not easy.

Absolute Density of water can only be measured in special laboratories, and only with difficulty. The best accuracy is  $1 : 2.5 \cdot 10^5$  = 4 parts per million.

To avoid the difficulty of working with absolute density, oceanographers use **density relative to density of pure water**. Density  $r$  (S, t, p) is now defined using Standard Mean Ocean Water of known isotopic composition, assuming saturation of dissolved atmospheric gasses.

S, t, p refer to salinity, temperature, and pressure

## Physical properties of sea water

**Density** of water at the sea surface is typically  $1027 \text{ kg m}^{-3}$ .


For simplification, physical oceanographers often quote only the last 2 digits of the density, a quantity they call density anomaly or Sigma ( $S, t, p$ ):

$$\sigma(S, t, p) = \rho(S, t, p) - 1000 \text{ kg m}^{-3}$$

T°C	Salinity (g kg <sup>-1</sup> )				
	20	25	30	35	40
0	16.04	20.06	24.08	28.10	32.14
5	15.84	19.78	23.73	27.68	31.64
10	15.31	19.18	23.07	26.96	30.86
15	14.48	18.30	22.13	25.97	29.82
20	13.39	17.17	20.96	24.75	28.56
25	12.07	15.82	19.57	23.34	27.12



## Vertical variability of seawater temperature and salinity



Typical temperature and salinity profiles in the open ocean. **AAC**: At 62.0. S, 170.0. E in the Antarctic Circumpolar Current on 16 January 1969 as measured by the R/V Hakuho Maru. **Warm Pool**: At 9.5. N 176.3. E in the tropical west Pacific **warm pool** on 12 March 1989 as measured by Bryden and Hall on the R/V Moana Wave. **BATS**: At 31.8. N 64.1. W near Bermuda on 17 April and 10 September 1990 as measured by the Bermuda Biological Station for Research, Inc.

## Physical properties of sea water

### Potential Density


Potential density  $s_t$  is the density a parcel of water would have if it were raised adiabatically to the surface without change in salinity. Written as sigma,  $s_t = s(S, t, 0)$  Potential Density is especially useful because it is a conservative thermodynamic property.

### Potential Temperature


Potential temperature is defined as the temperature of a parcel of water at the sea surface after it has been raised adiabatically from some depth in the ocean. Raising the parcel adiabatically means that it is raised in an insulated container so it does not exchange heat with its surroundings.

# Variability of seawater temperature


*Average Sea-Surface Temperature for July*



*Average Sea-Surface Temperature for January*



# Variability of seawater salinity



## Variability of seawater density






Figure 6.10 Vertical sections of density in the western Atlantic. Note that the depth scale changes at 1000 m depth. **Upper:**  $\sigma_\theta$ , showing an apparent density inversion below 3,000 m. **Lower:**  $\sigma_4$  showing continuous increase in density with depth. After Lynn and Reid (1968).





<https://www.youtube.com/watch?v=p4pWafuvdrY>

# Physical properties of sea water

## (function of temperature, salinity and pressure)

**Mechanical and thermal properties of sea water at salinity 35 g kg<sup>-1</sup> and atmospheric pressure (unless otherwise stated)**

Property	0 °C	20 °C
Dynamic viscosity . . . . .	$1.88 \times 10^{-3} \text{ Pa s}$	$1.08 \times 10^{-3} \text{ Pa s}$
Kinematic viscosity, $\nu$ . . . . .	$1.83 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$	$1.05 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$
Thermal conductivity . . . . .	$0.563 \text{ W m}^{-1} \text{ K}^{-1}$	$0.596 \text{ W m}^{-1} \text{ K}^{-1}$
Thermal diffusivity, $\kappa$ . . . . .	$1.37 \times 10^{-7} \text{ m}^2 \text{ s}^{-1}$	$1.46 \times 10^{-7} \text{ m}^2 \text{ s}^{-1}$
Prandtl number, $\nu/\kappa$ . . . . .	13.4	7.2
Specific heat capacity, $C_p$ . . . . .	$3985 \text{ J kg}^{-1} \text{ K}^{-1}$	$3993 \text{ J kg}^{-1} \text{ K}^{-1}$
Thermal expansion coefficient . . . . .		
Pressure = $0.1 \text{ MN m}^{-2}$ . . . . .	$52 \times 10^{-6} \text{ K}^{-1}$	$250 \times 10^{-6} \text{ K}^{-1}$
Pressure = $100 \text{ MN m}^{-2}$ . . . . .	$244 \times 10^{-6} \text{ K}^{-1}$	$325 \times 10^{-6} \text{ K}^{-1}$
Ratio of specific heat capacities, $C_p / C_v$ . . . . .	1.000 4	1.010 6
Velocity of sound* . . . . .	$1449 \text{ m s}^{-1}$	$1522 \text{ m s}^{-1}$
Compressibility . . . . .	$4.65 \times 10^{-10} \text{ Pa}^{-1}$	$4.28 \times 10^{-10} \text{ Pa}^{-1}$
Freezing point . . . . .		- 1.910 °C
Boiling point . . . . .		100.56 °C

	Seawater (35‰)	Pure Water
Temperature of maximum density:	3.25°C	3.98°C
Freezing point	-1.91°C	0.00°C

## Pore water in marine sediments

Chemically unstable sediment particles:

Igneous rock fragments

Amorphous material (e.g. opal)


Organic matter

Remnants of Organisms (bones, teeth)


Clay minerals

Aragonite and Mg Calcite


# Pore water in marine sediments



## Pore water in marine sediments




depth under sediment surface




concentration




## Pore water Sampling in in marine sediments



## Pore water Sampling in in marine sediments



## Pore water Sampling in in marine sediments





## Pore water Sampling in in marine sediments



## References

- Worthington L.V. 1981. The water masses of the World Ocean: Some results of a fine-scale census. In: Evolution of Physical Oceanography: Scientific surveys in honor of Henry Stommel. Edited by B. A. Warren and C. Wunsch. 42–69. Cambridge: Massachusetts Institute of Technology.
- Millero F.J., C.-T. Chen, A. Bradshaw, and K. Schleicher. 1980. A new high pressure equation of state for seawater. Deep-Sea Research 27A: 255–264.
- Millero F.J., and A. Poisson. 1981. International one-atmosphere equation of state of seawater. Deep-Sea Research 28A (6): 625–629.
- JPOTS Joint Panel on Oceanographic Tables and Standards. 1981. The practical salinity scale 1978 and the international equation of state of seawater 1980. Paris: unesco Technical Papers in Marine Science 36: 25.
- Stewart, R.D., 2007. Introduction To Physical Oceanography. Texas A & M University, Department of Oceanography, 345 pp.  
[http://oceanworld.tamu.edu/resources/ocng\\_textbook/PDF\\_files/book\\_pdf\\_files.html](http://oceanworld.tamu.edu/resources/ocng_textbook/PDF_files/book_pdf_files.html)