

RESEARCH ARTICLE

Does full protection count for the maintenance of β -diversity patterns in marine communities? Evidence from Mediterranean fish assemblages

Luca Appolloni¹ | Stanislao Bevilacqua² | Luisa Sbrescia¹ | Roberto Sandulli¹ | Antonio Terlizzi^{3,4} | Giovanni Fulvio Russo¹

¹Department of Science and Technology, CoNISMa URL University "Parthenope" of Naples, Naples, Italy

²Department of Biological and Environmental Sciences and Technologies, University of Salento, Lecce, Italy

³Department of Life Sciences, University of Trieste, CoNISMa, Trieste, Italy

⁴Stazione Zoologica Anton Dohrn, Naples, Italy

Correspondence

Luca Appolloni, Centro direzionale isola C4 80143, Naples, Italy.

Email: luca.appolloni@uniparthenope.it

Abstract

1. Although it is widely recognized that protection may enhance size, abundance, and diversity of fish, its effect on spatial heterogeneity of fish assemblages and species turnover is still poorly understood.
2. Here the effect of full protection within a Mediterranean marine protected area on β -diversity patterns of fish assemblages along a depth gradient comparing a no-take zone with multiple unprotected areas is explored. The no-take zone showed significantly higher synecological parameters, higher β -diversity among depths, and lower small-scale heterogeneity of fish assemblages relative to unprotected areas.
3. Such patterns might likely depend on the high level of fishing pressure outside the no-take zone, as also abundance-biomass curves seemed to indicate. Results suggested that full protection could play a role in maintaining high β -diversity, thus reducing the fragility of marine communities and ecosystems, and spatial heterogeneity may represent a reliable predictor of how management actions could provide insurance against undesirable phase shifts.

KEYWORDS

coastal, conservation evaluation, ecological status, fish, fishing, marine protected area

1 | INTRODUCTION

Over recent decades, marine protected areas (MPAs) have been recognized as useful tools in mitigating human-driven changes to marine ecosystems (Bevilacqua, Terlizzi, Frascchetti, Russo, & Boero, 2006; Edgar et al., 2014; Halpern, 2003; Micheli & Halpern, 2005). They play an important role in contrasting biodiversity loss (Allison, Lubchenco & Carr, 1998), preventing the overexploitation of natural resources (Roberts, Bohnsack, Gell, Hawkins, & Goodridge, 2001; Rogers-Bennett, Haaker, Karpov, & Kushner, 2002) and enhancing fishery yields within and outside their boundaries (Alcala, Russ, Maypa, & Calumpong, 2005; Di Franco et al., 2012; Goñi et al., 2008).

Most evidence on the effectiveness of MPAs stems from quantitative assessments on fish populations (Claudet et al., 2008; Guidetti et al., 2014; McClanahan, Graham, Calnan, & MacNeil, 2007), owing to their rapid and direct response to protection, especially in no-take areas, leading to a general increase in size and abundance of fish (Claudet,

Garcia-Charton, & Lenfant, 2011; Dugan & Davis, 1993; Guidetti, 2006; Halpern & Warner, 2002). The commercial or ecological relevance of many fish species, moreover, prioritizes the re-establishment of natural stocks as a conservation target, which, in turn, makes fish privileged descriptors of the effectiveness of MPAs (Guidetti, 2006).

The effects of protection on fish assemblages have often been assessed taking into account single response variables, such as density, biomass, or diversity (Claudet & Frascchetti, 2010; Claudet et al., 2008; Guidetti et al., 2008; Lester et al., 2009). Yet, community-wide assessments may provide a more comprehensive picture of the ecological outcomes of MPAs, as changes in the multivariate structure of communities reflect the integrated response of species to human disturbance and protection regimes (Guidetti & Sala, 2007; McClanahan, Davies, & Maina, 2005; Micheli, Halpern, Botsford, & Warner, 2004). Several studies on different ecological compartments, from benthos to fish, demonstrated significant differences in assemblage structure when comparing protected versus unprotected areas (Ceccherelli, Casu, Pala,

Pinna, & Sechi, 2006; Frascchetti, Bevilacqua, Guarnieri, & Terlizzi, 2012; Micheli et al., 2005). Human impacts, in addition, are more likely to alter the spatial heterogeneity of assemblages, increasing or decreasing their spatial variability, rather than modifying exclusively their average structure (Frascchetti et al., 2001; Sandulli et al., 2004; Terlizzi et al., 2005; Warwick & Clarke, 1993). Therefore, it could be expected that protection could also play a role in maintaining natural patterns of spatial heterogeneity of assemblages by reducing the effects of anthropogenic disturbance on their spatial distribution.

The heterogeneity in the distribution of species composing assemblages within a given spatial extent reflects the β -component of species diversity (Anderson et al., 2011; Vellend, 2001). β -diversity was originally defined as the extent of change of community composition (Whittaker, 1960, 1972). Recently, it was also described as either species turnover or as non-directional variations in species composition (Anderson et al., 2011; Baeten et al., 2012). The first notion relates to directional changes in community composition from one position to another along any spatial, temporal or environmental gradient. The second notion, instead, considers variations occurring in community composition among a set of sample units within a given spatial, temporal, or environmental extent, without referring to any particular direction. Therefore, while β -diversity as turnover measures the extent of changes in species composition *between* two communities, β -diversity as variation measures the heterogeneity in species composition *within* communities (Bevilacqua, Plicanti, Sandulli, & Terlizzi, 2012).

β -diversity has a central role in tracking the relationships between local and regional diversity (Thrush et al., 2009; Witman, Etter, & Smith, 2004). It identifies biological and environmental factors structuring biogeographical gradients (Ellingsen, 2002) and, more generally, it highlights ecological processes that drive meta-community assembly (Baiser, Olden, Record, Lockwood, & McKinney, 2012; Thrush, Hewitt, Cummings, Norkko, & Chiantore, 2010). Measuring the spatial heterogeneity of assemblage structure may also provide useful insights into subtle mechanisms underlying ecological hetero-/homogenization related to anthropogenic activities (Balata, Piazza, & Benedetti-Cecchi, 2007; Bevilacqua et al., 2012). Human impacts, in fact, could cause spatial homogeneity of assemblages and smooth natural environmental gradients by selecting a few dominant species that tolerate perturbed environmental conditions (Balata et al., 2007; Claudet & Frascchetti, 2010; Magurran, Dornelas, Moyes, Gotelli, & McGill, 2015), or could increase assemblage heterogeneity by creating a mosaic of disturbed patches (Frascchetti et al., 2001; Guarnieri, Terlizzi, Bevilacqua, & Frascchetti, 2012).

Quantifying patterns of spatial heterogeneity in assemblage structure is indispensable for guiding the siting and spacing of marine reserves (Margules & Pressey, 2000) in order to ensure an effective networking through adequate levels of complementarity among protected communities (Hewitt, Thrush, Halliday, & Duffy, 2005; Winberg, Lynch, Murray, Jones, & Davis, 2007).

Communities characterized by high β -diversity should be more resistant to disturbance, whether natural or anthropogenic, than others, acting as refuges from which neighbouring patches can be sustained (Soininen, 2010). In this respect, β -diversity has been proposed as a measure of ecological connectivity (de Juan, Thrush, & Hewitt, 2013; Thrush et al., 2009), thus providing an indication of

the potential resilience of assemblages (Thrush et al., 2010). At the local scale, strong connectivity among patches could promote recovery after disturbance through supplementary inputs from the whole system (Scheffer et al., 2012). Several studies have documented the role of protection in enhancing the recovery trajectories of disturbed assemblages (Bellwood, Hughes, Folke, & Nyström, 2004; Bevilacqua et al., 2006; Micheli et al., 2004; Mumby et al., 2006), or in increasing their temporal stability (Frascchetti et al., 2012). However, despite the recognized importance of evaluating direct and indirect effects of protection in modifying patterns of spatial heterogeneity of marine assemblages (Benedetti-Cecchi et al., 2003; Charton & Ruzafa, 1999), very few attempts have been made in this direction (Frascchetti, Guarnieri, Bevilacqua, Terlizzi, & Boero, 2013; Guarnieri et al., 2012).

Here, the effects of full protection on the structure and spatial heterogeneity of fish assemblages within a Mediterranean MPA are assessed. Specifically, the response of fish assemblages associated with rocky reefs at different depths are investigated in order to test if: (i) full protection had an effect on the overall assemblage structure, total abundance and biomass, and species richness; (ii) patterns of change in assemblages along the investigated depth gradient were comparable between protected and unprotected areas; (iii) protected and unprotected assemblages exhibited different levels of spatial heterogeneity and if such patterns were consistent among depths.

2 | MATERIALS AND METHODS

2.1 | Study area

The MPA of Punta Campanella extends over 1300 ha, from the coastline to 55–60 m depth, around the distal part of the Sorrento peninsula between the Bay of Naples and the Bay of Salerno (Tyrrhenian Sea) (Figure 1). The MPA is embedded within a human-dominated environmental context with high levels of anthropogenic pressure on the whole coastal ecosystem (Bavestrello, Bo, Canese, Sandulli, & Cattaneo-Vietti, 2014; Cianelli et al., 2011).

The no-take zone considered in the present study, covering 21 ha, is located at about 0.7 nm from the coast around the emerging part of a carbonate pinnacle. A partial protected area (hereafter referred to as 'unprotected area'), where most human activities are allowed (fishing and diving included), surrounds the no-take zone (hereafter referred to as 'protected area'). Geo-morphological features of the rocky coast in the protected area are comparable with those characterizing the unprotected area. Both areas comprise submerged cliffs covered with pre-coraligenous assemblages alternating with rocky terraces dominated by macro-algal stands.

2.2 | Sampling design

Sampling was carried out between June and October 2008 on rocky cliffs within the only protected zone (P) in the Bay of Naples and within an unprotected zone (F) relatively close to the no-take zone (Figure 1). The no-take zone (island of Veruce) is a very small-emerged rock where it was not possible to define more than a single visual census sampling site. However, the unprotected zone is located along the more extended coastline of the Sorrento peninsula where six sampling sites




FIGURE 1 Punta Campanella MPA and location of study area with protected (P) and unprotected (F) sites

(F1, F2, F3, F4, F5, F6) were selected to detect possible differences in the fish assemblages of the rocky reefs of the unprotected zone.

At each site, fish assemblages were sampled at three depths (5 m, 10 m, 20 m). For each site and each depth, abundance of fish species and size of individuals were assessed along three strip transects, for a total of 63 sampling units. Fish species were assigned to seven ranks of abundance (1; 2–5; 6–10; 11–30; 31–50; 51–100; >101) according to La Mesa and Vacchi (1999), and three size categories (i.e. small, medium, large) based on the maximum total length attained by each species were used. Biomass of fishes was estimated using the population size-structure and length–weight relationship (Froese & Pauly, 2012).

Visual counts along strip transects (L = 25 m; W = 5 m; H = 5 m) according to standardized procedures (Harmelin-Vivien et al., 1985) were performed when the horizontal visibility was at least 2.5 m.

2.3 | Data analysis

A distance-based permutation multivariate analysis of variance (PERMANOVA; Anderson, 2001a) was carried out on density and biomass, in order to test for differences in assemblage structure imputable to protection. As sites in the unprotected area were spatially segregated, with three sites north and three sites south of the protected area, a preliminary analysis was carried out considering only unprotected sites in order to exclude confounding effects due to the different geographic position. The design for the analysis involved the factor Depth

(De, fixed, three levels), Position (Po, two levels, fixed), and Site (Si(Po), random, three levels, nested in Po) with $n = 3$. No significant differences were detected for factor Position either analysing abundance or biomass of fish assemblages or all the investigated single response variables (see Results). This allowed testing for differences in fish assemblages related to protection by contrasting the single site within the protected area against all six sites from the unprotected area. In this case, the design for the analysis comprised three factors: Depth (De, fixed, three levels), Protection (Pr, fixed, two levels), and Site (Si(Pr), random, nested in Pr). As there was only one site within the protected area the design was asymmetrical (Glasby, 1997; Terlizzi et al., 2005).

Prior to analyses, fish density and biomass data were square root transformed (Guidetti et al., 2014) to reduce the weight of very abundant species. All multivariate analyses were based on Bray–Curtis similarity, and each term in the analysis was tested by 4999 random permutations (Anderson, 2001b; Anderson & Braak, 2003). Post hoc pair-wise comparisons using the PERMANOVA t -statistic and 4999 permutations were conducted if necessary. Multivariate patterns were visualized through nMDS ordination of De \times Pr centroids. Permutation tests of multivariate dispersion (PERMDISP; Anderson, Ellingsen, & McArdle, 2006) based on Bray–Curtis similarity matrices were carried out for the interaction term De \times Pr to investigate possible effects of protection on small-scale (among replicates) spatial heterogeneity of fish assemblages. Multivariate dispersion values were employed as a

measure of variation in β -diversity (Anderson et al., 2006) between protected and unprotected areas.

PERMANOVA based on Euclidean distance (Terlizzi, Anderson, Frascetti, & Benedetti-Cecchi, 2007) was conducted on species richness, total abundance and total biomass of the whole assemblage, and abundance and biomass of target species (classified into two categories, namely commercial species and species with low commercial value), and non-target species. Species were assigned to different categories according to Froese and Pauly (2012). The design for analyses was the same as for multivariate analyses.

Finally, ABC curves (Warwick & Clarke, 1994) were employed to compare abundance-biomass relationships between protected and unprotected areas.

3 | RESULTS

Thirty-five fish taxa from 14 families were identified. Some taxa were recorded only in the protected area (i.e. *Sphyaena viridensis*, *Diplodus puntazzo*, *Muraena helena*).

PERMANOVA on data (abundance and biomass) from the unprotected area showed no significant effect of geographic position of sites. Also, no significant differences due to position were detected analysing all the investigated single response variables (Table 1). This allows us to test for differences in the contrast between the protected site versus all the unprotected sites.

PERMANOVA on fish abundance and biomass based on the full dataset detected a significant De \times Pr interaction, indicating that differences in assemblage structure among depths varied between protected versus unprotected areas (Table 2).

Post hoc pair-wise comparisons from abundance data showed that, in the unprotected area, fish assemblages did not differ among depths. In contrast, in the protected area, the depth gradient was clear with assemblages differing at each depth. Patterns of difference in biomass among depths were consistent with those for abundance data for the protected area, whereas in the unprotected area, assemblages at 20 m differed from those characterizing the shallower depths (Table 2).

Such patterns were clearly depicted in the nMDS ordination plots of De \times Pr centroids that also showed a clear separation of fish assemblages between protected and unprotected areas (Figure 2).

TABLE 1 Summary of results of PERMANOVA tests for differences related to geographic position in fish assemblages and single response variables. Only tests relevant to the hypothesis were reported; De \times Po = interaction between Depth (De) and Position (Po) (see Method section for further details)

	De \times Po		Po	
	Pseudo-F	P	Pseudo-F	P
Fish assemblages (abundance data)	0.6	0.791	1.1	0.402
Fish assemblages (biomass data)	0.7	0.716	1.1	0.379
Species richness	4.2	0.060	1.3	0.317
Total abundance	4.3	0.060	4.1	0.111
Total biomass	0.8	0.549	3.2	0.135
Total abundance target species	1.9	0.220	1.3	0.305
Total biomass target species	0.4	0.677	0.8	0.407
Total abundance non-target species	1.2	0.346	0.1	0.819
Total biomass non-target species	0.8	0.509	2.0	0.221

TABLE 2 PERMANOVA results for difference in abundance and biomass of fish assemblages between protected and unprotected areas. Pair-wise comparisons for differences in fish assemblages among depths in protected (Pr) and unprotected (F) areas are also reported. Significant tests relevant to the hypothesis are given in bold

Abundance					
Source	df	SS	MS	Pseudo-F	P(perm)
De	2	5812.5	2906.3	41.559	
Pr	1	10571.0	10571.0	20.877	
Si(Pr)	5	25317.0	5063.5	12.616	
De \times Pr	2	2676.0	1338.0	19.133	0.023
De \times Si(Pr)	10	6993.1	699.3	17.424	0.003
Res	42	16857.0	401.3		
Total	62	69761.0			

Groups	Within F		Within Pr	
	t	P	t	P
5 m - 10 m	1.436	0.099	2.135	0.048
5 m - 20 m	26.921	0.108	36.511	0.005
10 m - 20 m	19.547	0.252	29.298	0.011

Biomass					
Source	df	SS	MS	Pseudo-F	P(perm)
De	2	10933.0	5466.4	41.889	
Pr	1	21824.0	21824.0	27.247	
Si(Pr)	5	40047.0	8009.5	10.176	
De \times Pr	2	7165.0	3582.5	27.452	0.004
De \times Si(Pr)	10	13050.0	1305.0	1.658	0.001
Res	42	33057.0	787.1		
Total	62	128000.0			

Groups	Within F		Within Pr	
	t	P	T	P
5 m - 10 m	15.401	0.067	27.482	0.015
5 m - 20 m	25.466	0.011	26.376	0.015
10 m - 20 m	19.346	0.054	27.294	0.011

Results of PERMDISP showed a greater small-scale heterogeneity of fish assemblages in unprotected areas with respect to the protected area at each of the investigated depths (Figure 3; Table 3), whereas multivariate dispersion did not differ significantly along the depth gradient in both areas.





FIGURE 2 3-d nMDS plot of DexPr centroids for abundance (a) and biomass (b). White circles = protected area, black circles = unprotected areas. Depths are also indicated (5 m, 10 m, 20 m)

PERMANOVA on species richness, total abundance and total biomass detected a significant De \times Pr interaction for all the investigated variables (Table 4). However, pair-wise tests showed higher species richness, total abundance, and total biomass comparing protected versus unprotected areas at each depth (Figure 4).

PERMANOVA on total abundance of target species detected a significant De \times Pr interaction (Table 4) and subsequent pair-wise comparisons showed a significantly higher abundance of target species in the protected area with respect to unprotected areas at 5 m and 10 m but not at 20 m (Figure 5). Similar results were obtained when analysing total biomass of target species, although significantly higher




FIGURE 3 Mean (\pm SE, $n = 3$) multivariate dispersion of replicates around De \times Pr centroids calculated on abundance (a) and biomass (b) data for unprotected (black bars) and protected (grey bars) areas

TABLE 3 Results of test for differences in multivariate dispersion (PERMDISP) between protected (Pr) and unprotected (F) areas for abundance and biomass data at different depths

Abundance		Biomass	
F	P	F	P
9.5718	0.0002	7.7374	0.0004
Pair-wise comparisons			
	t	P	
5 m Pr - 5 m F	41.724	0.001	5 m Pr - 5 m F
10 m Pr - 10 m F	45.705	0.002	10 m Pr - 10 m F
20 m Pr - 20 m F	29.414	0.019	20 m Pr - 20 m F

biomass of target species characterized the protected area at each of the depths considered (Figure 5). No significant differences between protected versus unprotected areas were recorded for both total abundance and biomass of non-target species (Table 4).

The ABC curves showed that the abundance curve was above the biomass curve ($W = -0.067$) in fished areas, whereas the opposite occurred in the protected area ($W = 0.318$), indicating a typical pattern of stressed condition of fish assemblages in the unprotected area (Figure 6).

4 | DISCUSSION

The structure of fish assemblages differed significantly in abundance or biomass for protected and unprotected areas. These results demonstrate the potential of full protection to significantly increase diversity, number and size of individuals of fish species as shown in several other

TABLE 4 Summary of results of PERMANOVA tests (see text for details) for species richness, total abundance and total biomass of fish assemblages, and for total abundance and biomass of target and non-target species. Significant tests relevant to the hypothesis are given in bold

	De × Pr		Pr	
	Pseudo-F	P	Pseudo-F	P
Species richness	4.7	0.040		
Total abundance	8.6	0.015		
Total biomass	5702.8	0.000		
Total abundance target species	56.4	0.001		
Total biomass target species	83400.0	0.000		
Total abundance non-target species	2.1	0.170	1.5	0.150
Total biomass non-target species	0.2	0.747	0.1	0.283




FIGURE 4 Mean (\pm SE, $n = 6$) values of (a) species richness, (b) total abundance (no. of individuals), and (c) total biomass (log scale) of fish assemblages for each depth in protected (grey bars) and unfished areas (black bars). Results of pair-wise tests are also reported. * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$

studies (Denny & Babcock, 2004; Di Franco, Bussotti, Navone, Panzalis, & Guidetti, 2009; Floeter, Halpern, & Ferreira, 2006; García-Charton et al., 2004; Halpern, 2003; Harmelin-Vivien et al., 2008; Lester et al., 2009; Seytre et al., 2013). It could be argued that the observed differences between the protected and unprotected areas might depend on the insular nature of the protected site with respect to the unprotected sites on the mainland, since isolated spots of hard substrates could serve as attractors for fish (Cresson, Ruitton, & Harmelin-Vivien, 2014; Scott, Smith, Lowry, Taylor, & Suthers, 2015). However, the abundance-biomass curves highlighted highly stressed




FIGURE 5 Mean (\pm SE, $n = 6$) values of (a) total abundance (no. of individuals) and (b) total biomass (log scale) of target species for each depth in protected (grey bars) and unfished areas (black bars). Results of pair-wise tests were also reported. n.s. = not significant; ** = $P < 0.01$; *** = $P < 0.001$

conditions in unprotected assemblages (Yemane, Field, & Leslie, 2005), suggesting that the observed differences between protected and unprotected areas are human-driven (Matić-Skoko et al., 2011; Penczak & Kruk, 1999). Moreover, differences between protected and unprotected areas were mostly driven by changes in abundance and biomass of commercial species, such as large predators (e.g. *Diplodus puntazzo*, *Epinephelus marginatus*, *Sphyrna viridensis*), whereas non-target fish were almost unaffected by protection. The removal of large individuals from target populations is a typical outcome of intense fishing activities (Erzini et al., 2006; Pelletier et al., 2005), indicating that the high fishing pressure outside the no-take area (Bavestrello et al., 2014; Sbrescia, Di Stefano, Russo, & Russo, 2008, 2010) is likely responsible for the observed patterns, over and beyond site-specific natural variations (e.g. isolation, geomorphology) between protected and unprotected areas.




FIGURE 6 Abundance-biomass curves for unprotected (a) and protected (b) areas

In the MPA of Punta Campanella, fishing activity is allowed within 6 nautical miles from the coast only to local fishermen, and by means of <10 GT boats. The most widely used gears are long-lines and gill nets that allow the capture of pelagic and demersal species (Sbrescia et al., 2008, 2010), most of which are large top predators (Lloret, Zaragoza, Caballero, & Riera, 2008; Pauly et al., 2002; Planque et al., 2010). These species are frequently the most susceptible to fishing (Babcock et al., 2010; Pauly et al., 2002; Perry et al., 2010), and consequently respond positively to protection (Babcock et al., 2010; Bianchi & Morri, 2000; Halpern, 2003; Harmelin, Bachet, & Garcia, 1995). Although long-lines and gill nets are generally considered as highly selective gears (Stobart et al., 2009 and references therein), they also catch numerous by-catch species. Therefore, the high intensity of fishing in the area (Sbrescia et al., 2008, 2010) could have also had more subtle direct and indirect effects on spatial patterns of the whole fish assemblage (Farina et al., 2008; Griffin, Pearce, & Handy, 2012; Marriott, Latchford, & McCarthy, 2010; Sala & Lucchetti, 2010), including changes in small-scale heterogeneity.

In the absence of strong sources of human disturbance, habitat type and related environmental features are among the main factors structuring fish assemblages (Brokovich, Baranes, & Goren, 2006; Lara & González, 1998; McGehee, 1994; Wantiez & Chauvet, 2003), determining their species composition and trophic structure (Carpenter, Mclat, Albaladejo, & Corpuz, 1981; Friedlander & Parrish, 1998; Meekan, Steven, & Fortin, 1995; Nanami & Nishihira, 2002). In such context, spatial distribution of fish assemblages largely reflects local environmental gradients (Hewitt et al., 2005), and well-structured associated patterns of distribution of fish species can be considered as a feature of good ecological conditions (Becker, Levin, Fodrie, & McMillan, 2007; Caley et al., 1996; Levin & Talley, 2002; Lundquist et al., 2006). In the no-take zone, fish assemblages were distinct at

each depth, although assemblages at 10 m depth seemed to have an intermediate structure. At this depth the highest number of species was recorded, composed of a combination of those species found at 5 m and 20 m; it represents a mid-point for the increase of Labridae and Serranidae, and for the decrease of Sparidae and Pomacentridae abundances and, thus, is a transition zone between shallow and deeper assemblages due to the mid-domain effect (Colwell & Lees, 2000, Connolly, Bellwood, & Hughes, 2003; Zapata, Gaston, & Chown, 2003, 2005). In contrast, assemblages in unprotected areas showed comparable structure among depths, indicating disruption of the natural depth gradient as a likely result of fishing pressure.

Results also showed the potential of fishing pressure to significantly increase small-scale heterogeneity (patchiness) of assemblages. Several studies have outlined increased heterogeneity of assemblages as a result of human disturbance (Balata et al., 2007; Bevilacqua et al., 2012; Claudet & Fraschetti, 2010; Warwick & Clarke, 1993). This phenomenon could be seen as a symptom of reduced ecological connectivity in the system (Scheffer et al., 2012; Thrush et al., 2010), which in turn might increase the fragility of the whole system (Fernandes et al., 2005; Steneck et al., 2009; Terlizzi et al., 2005; Thrush et al., 2009), leading to reduced resilience potential (Mellin, Bradshaw, Fordham, & Caley, 2013) and increased probability of phase shifts (Andersen, Carstensen, Hernandez-Garcia, & Duarte, 2009; Anderson & Thompson, 2004; Bevilacqua et al., 2012; Underwood, 1991). Moreover, reduced species richness in unprotected areas might further increase the vulnerability of fish assemblages to environmental changes due to reduced functional redundancy (Guillemot, Kulbicki, Chabanet, & Vigliola, 2011; Micheli & Halpern, 2005).

Resilience capacity is always the result of interactions between species (dominance, inhibition or facilitation), extrinsic factors (e.g. colonist supply and environmental setting) (Dayton, 1989; Thrush et al., 2009) and history of environmental exploitation (Jackson et al., 2001; Lotze & Milewski, 2004; Thrush, Halliday, Hewitt, & Lohrer, 2008). Full protection, through the exclusion of all direct forms of human disturbance, particularly fishing, may have important consequences for the dynamic properties of fish assemblages, such as their ability to persist and absorb fluctuations, by increasing their diversity and homogeneity in space (Holyoak, Leibold, Mouquet, Holt, & Hoopes, 2005). In unprotected areas, the exclusion of large predators and spawners, instead, may profoundly erode the resilience potential of the whole system (Hughes, Bellwood, Folke, McCook, & Pandolfi, 2007; Nyström, Folke, & Moberg, 2000), as a consequence of reduced diversity and disrupted top-down control (Guidetti, 2006; Levine, 1976; Mumby et al., 2006; Sala et al., 2012; Vandermeer, 1980).

Several studies highlighted the role of MPAs in enhancing the temporal stability and recovery trajectories of protected assemblages (Bevilacqua et al., 2006; Fraschetti et al., 2013; Hughes et al., 2007), and understanding how protection may affect such properties of marine communities represents a critical focus of future research in these systems (Mumby & Harborne, 2010). These results suggest that underlying mechanisms could include the potential for protection to mitigate the effects of human disturbance on species turnover along natural environmental gradients and on spatial heterogeneity of assemblages. In this respect, β -diversity may play an important role as a good predictor of how management actions could translate into

an increased ability of communities and ecosystems to escape regime shifts and reverse undesirable trends.

ACKNOWLEDGEMENTS

This research was supported financially by Punta Campanella MPA. We thank the director and staff of the reserve for their cooperation and, in particular, Nicola Guida for important help during the sampling phases.

REFERENCES

- Alcala, A. C., Russ, G. R., Maypa, A. P., & Calumpong, H. P. (2005). A long-term, spatially replicated experimental test of the effect of marine reserves on local fish yields. *Canadian Journal of Fisheries and Aquatic Sciences*, *62*, 98–108.
- Allison, G. W., Lubchenco, J., & Carr, M. H. (1998). Marine reserves are necessary but not sufficient for marine conservation. *Ecological Applications* 8(Special Issue), 79–92.
- Andersen, T., Carstensen, J., Hernandez-Garcia, E., & Duarte, C. M. (2009). Ecological thresholds and regime shifts: Approaches to identification. *Trends in Ecology & Evolution*, *24*, 49–57.
- Anderson, M. J. (2001a). A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, *26*, 32–46.
- Anderson, M. J. (2001b). Permutation tests for univariate or multivariate analysis of variance and regression. *Canadian Journal of Fisheries and Aquatic Sciences*, *58*, 626–639.
- Anderson, M. J., & Braak, C. T. (2003). Permutation tests for multi-factorial analysis of variance. *Journal of Statistical Computation and Simulation*, *73*, 85–113.
- Anderson, M. J., & Thompson, A. A. (2004). Multivariate control charts for ecological and environmental monitoring. *Ecological Applications*, *14*, 1921–1935.
- Anderson, M. J., Crist, T. O., Chase, J. M., Vellend, M., Inouye, B. D., Free-stone, A. L., & Swenson, N. G. (2011). Navigating the multiple meanings of β diversity: A roadmap for the practicing ecologist. *Ecology Letters*, *14*, 19–28.
- Anderson, M. J., Ellingsen, K. E., & McArdle, B. H. (2006). Multivariate dispersion as a measure of beta diversity. *Ecology Letters*, *9*, 683–693.
- Babcock, R. C., Shears, N. T., Alcala, A. C., Barrett, N. S., Edgar, G. J., Lafferty, K. D., ... Russ, G. R. (2010). Decadal trends in marine reserves reveal differential rates of change in direct and indirect effects. *Proceedings of the National Academy of Sciences of the United States of America*, *107*, 18256–18261.
- Baeten, L., Vangansbeke, P., Hermy, M., Peterken, G., Vanhuysse, K., & Verheyen, K. (2012). Distinguishing between turnover and nestedness in the quantification of biotic homogenization. *Biodiversity and Conservation*, *21*, 1399–1409.
- Baiser, B., Olden, J. D., Record, S., Lockwood, J. L., & McKinney, M. L. (2012). Pattern and process of biotic homogenization in the New Pangaea. *Proceedings of the Royal Society B: Biological Sciences*, *279*, 4772–4777.
- Balata, D., Piazzì, L., & Benedetti-Cecchi, L. (2007). Sediment disturbance and loss of beta diversity on subtidal rocky reefs. *Ecology*, *88*, 2455–2461.
- Bavestrello, G., Bo, M., Canese, S., Sandulli, R., & Cattaneo-Vietti, R. (2014). The red coral populations of the gulfs of Naples and Salerno: Human impact and deep mass mortalities. *Italian Journal of Zoology*, *81*, 552–563.
- Becker, B. J., Levin, L. A., Fodrie, F. J., & McMillan, P. A. (2007). Complex larval connectivity patterns among marine invertebrate populations. *Proceedings of the National Academy of Sciences of the United States of America*, *104*, 3267–3272.
- Bellwood, D. R., Hughes, T. P., Folke, C., & Nyström, M. (2004). Confronting the coral reef crisis. *Nature*, *429*, 827–833.
- Benedetti-Cecchi, L., Bertocci, I., Micheli, F., Maggi, E., Fosella, T., & Vaselli, S. (2003). Implications of spatial heterogeneity for management of marine protected areas (MPAs): Examples from assemblages of rocky coasts in the northwest Mediterranean. *Marine Environmental Research*, *55*, 429–458.
- Bevilacqua, S., Plicanti, A., Sandulli, R., & Terlizzi, A. (2012). Measuring more of β -diversity: Quantifying patterns of variation in assemblage heterogeneity. An insight from marine benthic assemblages. *Ecological Indicators*, *18*, 140–148.
- Bevilacqua, S., Terlizzi, A., Frascchetti, S., Russo, G. F., & Boero, F. (2006). Mitigating human disturbance: Can protection influence trajectories of recovery in benthic assemblages? *Journal of Animal Ecology*, *75*, 908–920.
- Bianchi, C. N., & Morri, C. (2000). Marine biodiversity of the Mediterranean Sea: Situation, problems and prospects for future research. *Marine Pollution Bulletin*, *40*, 367–376.
- Brokovich, E., Baranes, A., & Goren, M. (2006). Habitat structure determines coral reef fish assemblages at the northern tip of the Red Sea. *Ecological Indicators*, *6*, 494–507.
- Caley, M. J., Carr, M. H., Hixon, M. A., Hughes, T. P., Jones, G. P., & Menge, B. A. (1996). Recruitment and the local dynamics of open marine populations. *Annual Review of Ecology and Systematics*, *27*, 477–500.
- Carpenter, K. E., Micalot, R. I., Albaladejo, V. D., & Corpuz, V. T. (1981). The influence of substrate structure on the local abundance and diversity of Philippine reef fishes. In *Proceedings of the 4th International Coral Reef Symposium Penang, Malaysia: Reef Base*, *2*, 497–502.
- Ceccherelli, G., Casu, D., Pala, D., Pinna, S., & Sechi, N. (2006). Evaluating the effects of protection on two benthic habitats at Tavolara-Punta Coda Cavallo MPA (north-east Sardinia, Italy). *Marine Environmental Research*, *61*, 171–185.
- Charton, J. A. G., & Ruzafa, Á. P. (1999). Ecological heterogeneity and the evaluation of the effects of marine reserves. *Fisheries Research*, *42*, 1–20.
- Cianelli, D., Uttieri, M., Buonocore, B., Falco, P., Zambardino, G., & Zambianchi, E. (2011). Dynamics of a very special Mediterranean coastal area: The Gulf of Naples. In G. S. Williams (Ed.), *Mediterranean Ecosystems: Dynamics, Management and Conservation*. New York: Nova Science Publishers.
- Claudet, J., & Frascchetti, S. (2010). Human-driven impacts on marine habitats: A regional meta-analysis in the Mediterranean Sea. *Biological Conservation*, *143*, 2195–2206.
- Claudet, J., Garcia-Charton, J. A., & Lenfant, P. (2011). Combined effects of levels of protection and environmental variables at different spatial resolutions on fish assemblages in a marine protected area. *Conservation Biology*, *25*, 105–114.
- Claudet, J., Osenberg, C. W., Benedetti-Cecchi, L., Domenici, P., García-Charton, J. A., Pérez-Ruzafa, Á., ... Planes, S. (2008). Marine reserves: Size and age do matter. *Ecology Letters*, *11*, 481–489.
- Colwell, R. K., & Lees, D. C. (2000). The mid-domain effect: Geometric constraints on the geography of species richness. *Trends in Ecology & Evolution*, *15*, 70–76.
- Connolly, S. R., Bellwood, D. R., & Hughes, T. P. (2003). Indo-Pacific biodiversity of coral reefs: Deviations from a mid-domain model. *Ecology*, *84*, 2178–2190.
- Cresson, P., Ruitton, S., & Harmelin-Vivien, M. (2014). Artificial reefs do increase secondary biomass production: Mechanisms evidenced by stable isotopes. *Marine Ecology Progress Series*, *509*, 15–26.
- Dayton, P. K. (1989). Interdecadal variation in an Antarctic sponge and its predators from oceanographic climate shifts. *Science (Washington)*, *243*, 1484–1486.
- de Juan, S., Thrush, S. F., & Hewitt, J. E. (2013). Counting on β -diversity to safeguard the resilience of estuaries. *PLoS ONE*, *8*, e65575.
- Denny, C. M., & Babcock, R. C. (2004). Do partial marine reserves protect reef fish assemblages? *Biological Conservation*, *116*, 119–129.
- Di Franco, A., Bussotti, S., Navone, A., Panzalis, P., & Guidetti, P. (2009). Evaluating effects of total and partial restrictions to fishing on

- Mediterranean rocky-reef fish assemblages. *Marine Ecology Progress Series*, 387, 275–285.
- Di Franco, A., Coppini, G., Pujolar, J. M., De Leo, G. A., Gatto, M., Lyubartsev, V., ... Guidetti, P. (2012). Assessing dispersal patterns of fish propagules from an effective Mediterranean marine protected area. *PLoS ONE*, 7, e52108.
- Dugan, J. E., & Davis, G. E. (1993). Applications of marine refugia to coastal fisheries management. *Canadian Journal of Fisheries and Aquatic Sciences*, 50, 2029–2042.
- Edgar, G. J., Stuart-Smith, R. D., Willis, T. J., Kininmonth, S., Baker, S. C., Banks, S., ... Thomson, R. J. (2014). Global conservation outcomes depend on marine protected areas with five key features. *Nature*, 506, 216–220.
- Ellingsen, K. (2002). Spatial patterns of benthic diversity: Is there a latitudinal gradient along the Norwegian continental shelf? *Journal of Animal Ecology*, 71, 373–389.
- Erzini, K., Gonçalves, J. M., Bentes, L., Moutopoulos, D. K., Casal, J. A. H., Soriguer, M. C., ... Stergiou, K. I. (2006). Size selectivity of trammel nets in southern European small-scale fisheries. *Fisheries Research*, 79, 183–201.
- Farina, A. C., Azevedo, M., Landa, J., Duarte, R., Sampedro, P., Costas, G., ... Cañas, L. (2008). *Lophius* in the world: A synthesis on the common features and life strategies. *ICES Journal of Marine Science: Journal du Conseil*, 65, 1272–1280.
- Fernandes, L., Day, J., Lewis, A., Slegers, S., Kerrigan, B., Breen, D., ... Tapleton, K. (2005). Establishing representative no-take areas in the Great Barrier Reef: Large-scale implementation of theory on marine protected areas. *Conservation Biology*, 19, 1733–1744.
- Floeter, S. R., Halpern, B. S., & Ferreira, C. E. L. (2006). Effects of fishing and protection on Brazilian reef fishes. *Biological Conservation*, 128, 391–402.
- Fraschetti, S., Bevilacqua, S., Guarnieri, G., & Terlizzi, A. (2012). Idiosyncratic effects of protection in a remote marine reserve. *Marine Ecology Progress Series*, 466, 21.
- Fraschetti, S., Bianchi, C. N., Terlizzi, A., Fanelli, G., Morri, C., & Boero, F. (2001). Spatial variability and human disturbance in shallow subtidal hard substrate assemblages: A regional approach. *Marine Ecology Progress Series*, 212, 1–12.
- Fraschetti, S., Guarnieri, G., Bevilacqua, S., Terlizzi, A., & Boero, F. (2013). Protection enhances community and habitat stability: Evidence from a Mediterranean marine protected area. *PLoS ONE*, 8, e81838.
- Friedlander, A. M., & Parrish, J. D. (1998). Habitat characteristics affecting fish assemblages on a Hawaiian coral reef. *Journal of Experimental Marine Biology and Ecology*, 224, 1–30.
- Froese, R., & Pauly, D. (2012). Fish Base: World wide web electronic publication, version (09/2010). http://www.fishbase.org/manual/fishbasethe_length_weight_table.htm
- García-Charton, J. A., Pérez-Ruzafa, A., Sánchez-Jerez, P., Bayle-Sempere, J. T., Reñones, O., & Moreno, D. (2004). Multi-scale spatial heterogeneity, habitat structure, and the effect of marine reserves on Western Mediterranean rocky reef fish assemblages. *Marine Biology*, 144, 161–182.
- Glasby, T. M. (1997). Analysing data from post-impact studies using asymmetrical analyses of variance: A case study of epibiota on marinas. *Australian Journal of Ecology*, 22, 448–459.
- Goñi, R., Adlerstein, S., Alvarez-Berastegui, D., Forcada, A., Reñones, O., Criquet, G., ... Planes, S. (2008). Spillover from six western Mediterranean marine protected areas: Evidence from artisanal fisheries. *Marine Ecology Progress Series*, 366, 159–174.
- Griffin, R., Pearce, B., & Handy, R. D. (2012). Dietary preference and feeding selectivity of common dragonet *Callionymus lyra* in UK. *Journal of Fish Biology*, 81, 1019–1031.
- Guarnieri, G., Terlizzi, A., Bevilacqua, S., & Fraschetti, S. (2012). Increasing heterogeneity of sensitive assemblages as a consequence of human impact in submarine caves. *Marine Biology*, 159, 1155–1164.
- Guidetti, P. (2006). Marine reserves re-establish lost predatory interactions and cause community changes in rocky reefs. *Ecological Applications*, 16, 963–976.
- Guidetti, P., & Sala, E. (2007). Community-wide effects of marine reserves in the Mediterranean Sea. *Marine Ecology Progress Series*, 335, 43–56.
- Guidetti, P., Baiata, P., Ballesteros, E., Di Franco, A., Hereu, B., Macpherson, E., ... Sala, E. (2014). Large-scale assessment of Mediterranean Marine Protected Areas effects on fish assemblages. *PLoS ONE*, 9(4), e91841.
- Guidetti, P., Milazzo, M., Bussotti, S., Molinari, A., Murenu, M., Pais, A., ... Tunesi, L. (2008). Italian marine reserve effectiveness: Does enforcement matter? *Biological Conservation*, 141, 699–709.
- Guillemot, N., Kulbicki, M., Chabanet, P., & Vigliola, L. (2011). Functional redundancy patterns reveal non-random assembly rules in a species-rich marine assemblage. *PLoS ONE*, 6, e26735.
- Halpern, B. S. (2003). The impact of marine reserves: Do reserves work and does reserve size matter? *Ecological Applications*, 13, 117–137.
- Halpern, B. S., & Warner, R. R. (2002). Marine reserves have rapid and lasting effects. *Ecology Letters*, 5, 361–366.
- Harmelin, J. G., Bachet, F., & Garcia, F. (1995). Mediterranean marine reserves: Fish indices as tests of protection efficiency. *Marine Ecology*, 16, 233–250.
- Harmelin-Vivien, M. L., Harmelin, J. G., Chauvet, C., Duval, C., Galzin, R., Lejeune, P., ... Lasserre, G. (1985). Evaluation visuelle des peuplementsset populations de poissons méthodes et problèmes. *Revue D Ecologie-La Terre et la Vie*, 40, 467–539.
- Harmelin-Vivien, M., Le Diréach, L., Bayle-Sempere, J., Charbonnel, E., García-Charton, J. A., Ody, D., & Valle, C. (2008). Gradients of abundance and biomass across reserve boundaries in six Mediterranean marine protected areas: Evidence of fish spillover? *Biological Conservation*, 141, 1829–1839.
- Hewitt, J. E., Thrush, S. F., Halliday, J., & Duffy, C. (2005). The importance of small-scale habitat structure for maintaining beta diversity. *Ecology*, 86, 1619–1626.
- Holyoak, M., Leibold, M. A., Mouquet, N., Holt, R. D., & Hoopes, M. (2005). A framework for large scale community ecology. In M. Holyoak, M. A. Leibold, & R. D. Holt (Eds), *Metacommunities: Spatial dynamics and ecological communities* (pp. 1–31). Chicago: The University of Chicago Press.
- Hughes, T. P., Bellwood, D. R., Folke, C. S., McCook, L. J., & Pandolfi, J. M. (2007). No-take areas, herbivory and coral reef resilience. *Trends in Ecology and Evolution*, 22, 1–3.
- Jackson, J. B., Kirby, M. X., Berger, W. H., Bjorndal, K. A., Botsford, L. W., Bourque, B. J., ... Warner, R. R. (2001). Historical overfishing and the recent collapse of coastal ecosystems. *Science*, 293, 629–637.
- La Mesa, G., & Vacchi, M. (1999). An analysis of the coastal fish assemblage of the Ustica Island marine reserve (Mediterranean Sea). *Marine Ecology*, 20, 147–165.
- Lara, E. N., & González, E. A. (1998). The relationship between reef fish community structure and environmental variables in the southern Mexican Caribbean. *Journal of Fish Biology*, 53(sA), 209–221.
- Lester, S. E., Halpern, B. S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B. I., Gaines, S. D., ... Warner, R. R. (2009). Biological effects within no-take marine reserves: A global synthesis. *Marine Ecology Progress Series*, 384(2), 33–46.
- Levin, L. A., & Talley, T. S. (2002). Natural and manipulated sources of heterogeneity controlling early faunal development of a salt marsh. *Ecological Applications*, 12, 1785–1802.
- Levine, S. H. (1976). Competitive interactions in ecosystems. *American Naturalist*, 110, 903–910.
- Lloret, J., Zaragoza, N., Caballero, D., & Riera, V. (2008). Biological and socioeconomic implications of recreational boat fishing for the management of fishery resources in the marine reserve of Cap de Creus (NW Mediterranean). *Fisheries Research*, 91, 252–259.

- Lotze, H. K., & Milewski, I. (2004). Two centuries of multiple human impacts and successive changes in a North Atlantic food web. *Ecological Applications*, 14, 1428–1447.
- Lundquist, C. J., Thrush, S. F., Hewitt, J. E., Halliday, J., MacDonald, I., & Cummings, V. J. (2006). Spatial variability in recolonization potential: Influence of organism behaviour and hydrodynamics on the distribution of macrofaunal colonists. *Marine Ecology Progress Series*, 324, 67–81.
- Magurran, A. E., Dornelas, M., Moyes, F., Gotelli, N. J., & McGill, B. (2015). Rapid biotic homogenization of marine fish assemblages. *Nature Communications*, 6. doi:10.1028/ncomms9405
- Margules, C. R., & Pressey, R. L. (2000). Systematic conservation planning. *Nature*, 405, 243–253.
- Marriott, A. L., Latchford, J. W., & McCarthy, I. D. (2010). Population biology of the red gurnard (*Aspitrigla cuculus* L.; Triglidae) in the inshore waters of Eastern Anglesey and northwest Wales. *Journal of Applied Ichthyology*, 26, 504–512.
- Matić-Skoko, S., Stagličić, N., Pallaoro, A., Kraljević, M., Dulčić, J., Tutman, P., & Dragičević, B. (2011). Effectiveness of conventional management in Mediterranean type artisanal fisheries. *Estuarine, Coastal and Shelf Science*, 91, 314–324.
- McClanahan, T. R., Davies, J., & Maina, J. (2005). Factors influencing resource users and managers' perceptions towards marine protected area management in Kenya. *Environmental Conservation*, 32(01), 42–49.
- McClanahan, T. R., Graham, N. A., Calnan, J. M., & MacNeil, M. A. (2007). Toward pristine biomass: Reef fish recovery in coral reef marine protected areas in Kenya. *Ecological Applications*, 17, 1055–1067.
- McGehee, M. A. (1994). Correspondence between assemblages of coral reef fishes and gradients of water motion, depth, and substrate size off Puerto Rico. *Marine Ecology Progress Series*, 105, 243–255.
- Meekan, M. G., Steven, A. D. L., & Fortin, M. J. (1995). Spatial patterns in the distribution of damselfishes on a fringing coral reef. *Coral Reefs*, 14, 151–161.
- Mellin, C., Bradshaw, C. J., Fordham, D., & Caley, M. J. (2013). Strong but opposing β -diversity-stability relationships in coral reef fish communities. *Proceedings of The Royal Society B: Biological Sciences*, 281, 20131993.
- Micheli, F., & Halpern, B. S. (2005). Low functional redundancy in coastal marine assemblages. *Ecology Letters*, 8, 391–400.
- Micheli, F., Halpern, B. S., Botsford, L. W., & Warner, R. R. (2004). Trajectories and correlates of community change in no-take marine reserves. *Ecological Applications*, 14, 1709–1723.
- Micheli, F. L., Benedetti-Cecchi, S., Gambaccini, I., Bertocci, C., Borsini, G. C., & Romano, F. (2005). Alternate states, marine protected areas, and the structure of Mediterranean rocky-reef assemblages. *Ecological Monographs*, 75, 81–102.
- Mumby, P. J., & Harborne, A. R. (2010). Marine reserves enhance the recovery of corals on Caribbean reefs. *PLoS ONE*, 5(1), 1–7.
- Mumby, P. J., Dahlgren, C. P., Harborne, A. R., Kappel, C. V., Micheli, F., Brumbaugh, D. R., ... Gill, A. B. (2006). Fishing, trophic cascades, and the process of grazing on coral reefs. *Science*, 311, 98–101.
- Nanami, A., & Nishihira, M. (2002). The structures and dynamics of fish communities in an Okinawan coral reef: Effects of coral-based habitat structures at sites with rocky and sandy sea bottoms. *Environmental Biology of Fishes*, 63, 353–372.
- Nyström, M., Folke, C., & Moberg, F. (2000). Coral reef disturbance and resilience in a human-dominated environment. *Trends in Ecology & Evolution*, 15, 413–417.
- Pauly, D., Christensen, V., Guénette, S., Pitcher, T. J., Sumaila, U. R., Walters, C. J., ... Zeller, D. (2002). Towards sustainability in world fisheries. *Nature*, 418, 689–695.
- Pelletier, D., García-Charton, J. A., Ferraris, J., David, G., Thébaud, O., Letourneur, Y., ... Galzin, R. (2005). Designing indicators for assessing the effects of marine protected areas on coral reef ecosystems: A multidisciplinary standpoint. *Aquatic Living Resources*, 18, 15–33.
- Penczak, T., & Kruk, A. (1999). Applicability of the abundance/biomass comparison method for detecting human impacts on fish populations in the Pilica River, Poland. *Fisheries Research*, 39, 229–240.
- Perry, R. I., Cury, P., Brander, K., Jennings, S., Möllmann, C., & Planque, B. (2010). Sensitivity of marine systems to climate and fishing: Concepts, issues and management responses. *Journal of Marine Systems*, 79, 427–435.
- Planque, B., Fromentin, J. M., Cury, P., Drinkwater, K. F., Jennings, S., Perry, R. I., & Kifani, S. (2010). How does fishing alter marine populations and ecosystems sensitivity to climate? *Journal of Marine Systems*, 79, 403–417.
- Roberts, C. M., Bohnsack, J. A., Gell, F., Hawkins, J. P., & Goodridge, R. (2001). Effects of marine reserves on adjacent fisheries. *Science*, 294, 1920–1923.
- Rogers-Bennett, L., Haaker, P. L., Karpov, K. A., & Kushner, D. J. (2002). Using spatially explicit data to evaluate marine protected areas for abalone in southern California. *Conservation Biology*, 16, 1308–1317.
- Sala, A., & Lucchetti, A. (2010). The effect of mesh configuration and coded circumference on selectivity in the Mediterranean trawl Nephrops fishery. *Fisheries Research*, 103, 63–72.
- Sala, E., Ballesteros, E., Dendrinis, P., Di Franco, A., Ferretti, F., Foley, D., ... Zabala, M. (2012). The structure of Mediterranean rocky reef ecosystems across environmental and human gradients, and conservation implications. *PLoS ONE*, 7, e32742.
- Sandulli, R., Carriglio, D., Deastis, S., Marzano, A., Gerardi, D., Gallo D'Addabbo, M., & de Zio Grimaldi, S. (2004). Meiobenthic biodiversity in areas of the Gulf of Taranto (Italy) exposed to high environmental impact. *Chemistry and Ecology*, 20(1), 379–386.
- Sbrescia, L., Di Stefano, F., Russo, M., & Russo, G. F. (2008). Influence of sport fishing on gorgonians in MPA of Punta Campanella. *Biologia Marina Mediterranea*, 15, 172–173.
- Sbrescia, L., Di Stefano, F., Russo, M., & Russo, G. F. (2010). La pesca artigianale e ricreativa nell'AMP di Punta Campanella. *Workshop - Porto Cesareo 'Pesca e gestione nelle Aree Marine Protette'* (pp. 129–135).
- Scheffer, M., Carpenter, S. R., Lenton, T. M., Bascompte, J., Brock, W., Dakos, V., ... Vandermeer, J. (2012). Anticipating critical transitions. *Science*, 338, 344–348.
- Scott, M. E., Smith, J. A., Lowry, M. B., Taylor, M. D., & Suthers, I. M. (2015). The influence of an offshore artificial reef on the abundance of fish in the surrounding pelagic environment. *Marine and Freshwater Research*, 66, 429–437.
- Seytre, C., Vanderklift, M. A., Bodilis, P., Cottalorda, J. M., Gratiot, J., & Francour, P. (2013). Assessment of commercial and recreational fishing effects on trophic interactions in the Cap Roux area (north-western Mediterranean). *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23, 189–201.
- Soininen, J. (2010). Species turnover along abiotic and biotic gradients: Patterns in space equal patterns in time? *Bioscience*, 60, 433–439.
- Steneck, R. S., Paris, C. B., Arnold, S. N., Ablan-Lagman, M. C., Alcalá, A. C., Butler, M. J., ... Sale, P. F. (2009). Thinking and managing outside the box: Coalescing connectivity networks to build region-wide resilience in coral reef ecosystems. *Coral Reefs*, 28, 367–378.
- Stobart, B., Warwick, R., González, C., Mallol, S., Díaz, D., Reñones, O., & Goñi, R. (2009). Long-term and spillover effects of a marine protected area on an exploited fish community. *Marine Ecology Progress Series*, 384, 47–60.
- Terlizzi, A., Anderson, M. J., Fraschetti, S., & Benedetti-Cecchi, L. (2007). Scales of spatial variation in Mediterranean subtidal sessile assemblages at different depths. *Marine Ecology Progress Series*, 332, 25–39.
- Terlizzi, A., Benedetti-Cecchi, L., Bevilacqua, S., Fraschetti, S., Guidetti, P., & Anderson, M. J. (2005). Multivariate and univariate asymmetrical analyses in environmental impact assessment: A case study of Mediterranean subtidal sessile assemblages. *Marine Ecology Progress Series*, 289, 27–42.
- Thrush, S. F., Halliday, J., Hewitt, J. E., & Lohrer, A. M. (2008). The effects of habitat loss, fragmentation, and community homogenization on resilience in estuaries. *Ecological Applications*, 18, 12–21.
- Thrush, S. F., Hewitt, J. E., Dayton, P. K., Coco, G., Lohrer, A. M., Norkko, A., ... Chiantore, M. (2009). Forecasting the limits of resilience: Integrating

- empirical research with theory. *Proceedings of the Royal Society of London B: Biological Sciences*, 276, 3209–3217.
- Thrush, S. F., Hewitt, J. E., Cummings, V. J., Norkko, A., & Chiantore, M. (2010). β -diversity and species accumulation in Antarctic coastal benthos: Influence of habitat, distance and productivity on ecological connectivity. *PLoS ONE*, 5, e11899.
- Underwood, A. J. (1991). Beyond BACI: Experimental designs for detecting human environmental impacts on temporal variations in natural populations. *Marine and Freshwater Research*, 42, 569–587.
- Vandermeer, J. (1980). Indirect mutualism: Variations on a theme by Stephen Levine. *American Naturalist*, 116, 441–448.
- Vellend, M. (2001). Do commonly used indices of β -diversity measure species turnover? *Journal of Vegetation Science*, 12, 545–552.
- Wantiez, L., & Chauvet, C. (2003). First data on community structure and trophic networks of *Uvea* coral reef fish assemblages (Wallis and Futuna, South Pacific Ocean). *Cybium*, 27, 83–100.
- Warwick, R. M., & Clarke, K. R. (1993). Increased variability as a symptom of stress in marine communities. *Journal of Experimental Marine Biology and Ecology*, 172(1), 215–226.
- Warwick, R. M., & Clarke, K. R. (1994). Relearning the ABC: Taxonomic changes and abundance/biomass relationships in disturbed benthic communities. *Marine Biology*, 118, 739–744.
- Whittaker, R. H. (1960). Vegetation of the Siskiyou mountains, Oregon and California. *Ecological Monographs*, 30, 279–338.
- Whittaker, R. H. (1972). Evolution and measurement of species diversity. *Taxon*, 21, 213–251.
- Winberg, P. C., Lynch, T. P., Murray, A., Jones, A. R., & Davis, A. R. (2007). The importance of spatial scale for the conservation of tidal flat macrobenthos: An example from New South Wales, Australia. *Biological Conservation*, 134, 310–320.
- Witman, J. D., Etter, R. J., & Smith, F. (2004). The relationship between regional and local species diversity in marine benthic communities: A global perspective. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 15664–15669.
- Yemane, D., Field, J. G., & Leslie, R. W. (2005). Exploring the effects of fishing on fish assemblages using Abundance Biomass Comparison (ABC) curves. *ICES Journal of Marine Science: Journal du Conseil*, 62, 374–379.
- Zapata, F. A., Gaston, K. J., & Chown, S. L. (2003). Mid-domain models of species richness gradients: Assumptions, methods and evidence. *Journal of Animal Ecology*, 72, 677–690.
- Zapata, F. A., Gaston, K. J., & Chown, S. L. (2005). The mid-domain effect revisited. *The American Naturalist*, 166, E144–E148.

How to cite this article: Appolloni L, Bevilacqua S, Sbrescia L, Sandulli R, Terlizzi A, Russo, GF. Does full protection count for the maintenance of β -diversity patterns in marine communities? Evidence from Mediterranean fish assemblages. *Aquatic Conserv: Mar Freshw Ecosyst*. 2017;27:828–838. <https://doi.org/10.1002/aqc.2750>