



**Università di Trieste
Corso di Laurea in Geologia**

Anno accademico 2021 - 2022

Geologia Marina

Parte IV

**Modulo 4.4/5
Identifieri di movimento di fluidi: Idrati del metano**

Docente
A. Camerlenghi

B2 - Outline

Review of main mechanisms of fluid flow:

- Mud diapirs and mud volcanoes
- Gas chimneys
- Pockmarks
- Seafloor vents in general
- Polygonal fault systems
- Diagenetic fronts
- **Gas hydrates**

NATURAL GAS HYDRATES ARE CONSIDERED AS:

- **Natural controllers of climatic change**
- **Geo-hazards**
- **Potential energy source**
- **Controllers of deep sea eco-systems**
- **Potential safe and effective way of methane transport**
- **Problem in pipelines**
- ***Possible cause of ship and airplane disappearance in the Bermuda triangle***

METHANE

Is the simplest saturated hydrocarbon gas (alkane):



Other common hydrocarbon gases are:

ethane (C_2H_6), **propane** (C_3H_8) **butane** (C_4H_{10})

When the number of carbon atoms exceeds 5, the name of gases is from Greek numbers: **pentane**, **hexane**, **heptane**...

The general formula of hydrocarbon gases is:



Human-related sources

- **Landfills** (*waste disposal*)
- **Natural gas and petroleum systems** (*natural gas production*)
- **Coal mining**
- **Livestock enteric fermentation**
- **Livestock manure management**
- **Wastewater treatment**
- **Rice cultivation**

Natural sources


- **Wetlands**
- **Termites**
- **Oceans** (*anaerobic digestion in marine zooplankton and fish, and also from methanogenesis in sediment*)
- **Hydrates**
- **Juvenile**

HYDRATES OF NATURAL GAS

(discovered in 1810)


- Gas hydrates are a **solid phase** composed of **water** and low-molecular weight **gases** (predominantly methane)
- Water molecules form an ordered crystalline solid matrix which includes gas molecules with weak electro-static bounds
- The state of **saturation of hydrates is a function of pressure and temperature**. They form under conditions of low temperature, high pressure, and adequate gas concentration
- The ion content of water and the type of gas (methane, propane, pentane...) affect the phase diagram of the gas hydrate

Water molecule




Methane molecule

Hydrate crystal



Bohrmann and Torres, 2006, in Schulz, Horst D.; Zabel, Matthias (Eds.) Marine Geochemistry





RCOM - University of Bremen


Courtesy:

Carlo Giavarini, Filippo
Maccioni and Maria
Laura Santarelli

Università di Roma "La
Sapienza" Dipartimento
di Ingegneria Chimica.




PHASE DIAGRAM OF METHANE HYDRATES





Hydrate formation in arctic gas pipelines reduces or blocks methane flow. Chemical additives are necessary to prevent hydrate formation.



Alaska Highway Pipeline TransCanada

LIQUID NATURAL GAS TRANSPORT IS EXPENSIVE AND DANGEROUS



Methane is condensed into a liquid at almost atmospheric pressure


(Maximum Transport Pressure set around 25 kPa) by cooling it to approximately -163° C.

LNG is transported by specially designed **cryogenic sea vessels** and cryogenic road tankers, and stored in specially designed tanks.




self-preservation


At 1 Atm, CH₄ Hydrates are stable below -20° C.




Self-preservation is thought to be caused by a layer of ice from dissociated hydrate; this layer coats the hydrate and seals it from further dissociation.



PHASE DIAGRAM OF METHANE HYDRATES IN THE NATURAL ENVIRONMENT




METANE HYDRATES IN PERMAFROST






Source: International Permafrost Association, 1998.
Circumpolar Active-Layer Permafrost System (CAPS), version 1.0.


Permafrost is defined as soil, sediment or rock whose temperature is continuously below 0° C for at least two years

METHANE HYDRATES IN OCEANIC SEDIMENTS



Natural gas hydrates are distributed along the ocean margins world wide within the Gas Hydrate Stability Zone (GHSZ), that ranges from the seafloor down to a maximum of 700-800 m







Ruppel, C. D., and J. D. Kessler (2017), The interaction of climate change and methane hydrates, Rev. Geophys., 55, 126-168, doi:10.1002/2016RG000534.













UNIVERSITÀ
DEGLI STUDI
DI TRIESTE

Dipartimento
di matematica
e geoscienze


Corso di Geologia Marina 2021-22



OGS

Istituto Nazionale
di Oceanografia
e di Geofisica
Sperimentale





Van Dover et al. (2003). *Deep Sea Research*

METHODS OF DETECTION OF NATURAL GAS HYDRATES

- **SEISMICS**

(reflection, multichannel, OBS)

- **DRILLING**

(essentially through scientific drilling)

- **GEOCHEMICAL**


(inorganic- organic geochemistry)


- **MODELLING**

(laboratory models - data inversion)


- **BASIN-WIDE ANALYSES**

(modelling + data)



**SEISMIC STREAMER**


OCEAN BOTTOM SEISMOMETERS (OBS)




SEISMICS REFLECTION

The concept of
**Bottom
Simulating
Reflector
(BSR)**

The BSR is a seismic reflector that simulates the sea bottom reflector. It is produced by the acoustic impedance contrast between sediments with hydrates (above) and sediments with free gas (below).



SEISMIC PROFILE



Stratification in Marine Sediments




Turbidite (Gorgoglione Flysch),
Miocene, South Italy



Marne a Fucoidi (Sezione del Bosso)
Cretaceous (Albian), Central Apennines, Italy
FOTO MASSIMO GALLI

Makassar Straits, Indonesia



Data from TGS-NOPEC Geophysical Company, published in Max et al., 2006. Economic Geology of Natural Gas Hydrates, Springer.

Characteristics of the BSR

- **High reflection coefficient**
- **Reverse polarity with respect to the seafloor reflector**
- **Cuts through lithologic reflectors**
- **Its depth approximately coincides with the theoretical depth of the base of the hydrates stability zone.**
- **It may be underlined by another reflector produced by the base of the free gas layer (BGR)**

How can the hydrates be identified through drilling?

- Thermal anomaly in cores
 - Soupy layers
 - Chlorinity anomaly
 - Resistivity anomaly
- Seismic velocity anomaly



Thermal anomaly in cores




G. Bohrmann, Initial results from
M52/1 **MARGASCH**



Gas hydrate dissociation is an **endo-thermic** reaction (it absorbs heat from the surroundings >>> decrease in bulk sediment temperature)


SOUPY LAYERS: Sedimentological evidence of the presence of gas hydrates layers



Kastner et al., 1995


Soupy layers

Gas hydrate dissociation during core recovery destroys the sediment fabric, producing disturbed, liquified 'soupy' sediment layers in cores



Chlorinity anomaly


Gas hydrate dissociation during core recovery produces **residual fresh water** that dilutes the ionic concentration of the pore water



In situ Electrical Resistivity anomaly

The electrical resistivity of gas hydrates is **higher** than that of seawater.



The anomaly is identified with respect to a resistivity profile in a nearby borehole with similar sediment NOT containing hydrates



In situ seismic velocity anomaly

The compressional wave velocity of gas hydrates is **higher** than that of seawater.


The anomaly is identified with respect to a resistivity profile in a nearby borehole with similar sediment NOT containing hydrates



VELOCITY MODEL

Biot-Geerstma-Smit equations

$$V_p = \left\{ \left[\left(\frac{1}{C_m} + \frac{4}{3} \mu \right) + \frac{\phi_{eff} \rho_m}{k \rho_f} + \left(1 - \beta - 2 \cdot \frac{\phi_{eff}}{k} \right) \cdot (1 - \beta) \right] \cdot \frac{1}{\rho_m \left(1 - \frac{\phi_{eff}}{k} \frac{\rho_f}{\rho_m} \right)} \right\}^{1/2} \quad (1)$$



GLOBAL ESTIMATES


Estimation as amount of Carbon atoms · 10^{15} g

TERRESTRIAL LIVING ORGANISMS 830

WETLANDS AND OTHER 500

DISSOLVED IN WATER 980

SOILS 1.400





FOSSIL FULES 5.000

NATURAL GAS HYDRATES 10.000

Kvenvolden, various sources

HOWEVER.....



Global estimates of the volume of hydrate-bound gas in marine sediments versus the year in which the estimate was made.



Distribution of organic carbon in the Earth

(excluding dispersed organic carbon such as kerogen and bitumen)

- The distribution based on the estimate of 10000 Gt of methane carbon in gas hydrates (Kvenvolden, 1993).
- The distribution based on the revised estimate of the global gas hydrate inventory assuming the global volume of hydrate-bound gas at upper bound.
- The distribution based on the revised estimate of the global gas hydrate inventory assuming the global volume of hydrate-bound gas at lower bound.



In spite of changing global hydrates estimates.....

- Methane is the cleanest fossil fuel:



- Methane is presently the preferred fossil fuel for power generation in OECD countries,
- Methane demand is increasing also in other regions such as the Middle East, China and India
- Despite current high prices, the vast majority of new power generation on line in the review period will be gas-fired.


Methane Hydrates are considered as NON CONVENTIONAL HYDROCARBONS by most oil companies

- There is not an established extraction technology .
- The hydrates must be dissociated in situ by:
 - Heat injection
 - Injection of chemical inhibitors
 - depressurization
 -
- There are severe environmental implications and the methods will not be considered as economically feasible until the gas price will rise considerably.
- **USA, Japan, India, Korea** and other countries of the Pacific Area have started National Plans for the gas hydrates assessment in their Economic Exclusive Zones.

What Happens in Europe?

- No policy has been established so far
- Several EC projects have been dedicated to the matter from a purely scientific point of view (see next part of this talk)
- Oil companies and academia have focussed on this issue regarding the implications for submarine slope stability (see next part of this talk)
- **Regional evaluations of gas hydrate potential** of European Margins are in due course:
 - Atlantic European Margins
 - Mediterranean
 - Black Sea

HYDRATES PROSPECTIVITY in the Mediterranean Sea



D. Praeg, V. Unnithan, and A. Camerlenghi,
American Geophysical Union Fall Meeting,
December 2006


Due to:

- High temperature of bottom water
- High salinity of pore water
- Presence of a salt seal (Messinian) preventing deep fluid migration


The thickness of the Gas Hydrates Stability Zone is reduced in the Mediterranean Sea with respect to world Oceans.




Bathymetry



Bottom water temperatures (bwt)



Geothermal gradients



METHANE HYDRATES AND GLOBAL CLIMATE



MILANKOVITCH ORBITAL CYCLES


Frequencies of the main
Earth orbital variations

EARTH
ROTATION AXIS



Precession of the equinoxes
19 - 23 kyr

Earth axis
inclination
41 kyr

Orbital Excentricity
100 - 410 kyr



The mean Solar radiation received by the earth changes through time. Such changes are the primary cause of glacial interglacial epochs and other climatic stages.




Dome-C Station, Antarctica




Programma Nazionale di Ricerche in
Antartide (PNRA)

European Project for Ice Coring in Antarctica (EPICA)



Methane and CO₂ concentrations in the atmosphere since 800 ka from Environmental Protection Agency [2015]. Deep time data are from analysis of the European Project for Ice Coring in Antarctica (EPICA) Dome ice core and thus reflect Southern Hemisphere methane. Note the difference in time scales between the panels.

Ruppel, C. D., and J. D. Kessler (2017), The interaction of climate change and methane hydrates, Rev. Geophys., 55, 126-168, doi:10.1002/2016RG000534.



METHANE HYDRATES STORED IN PERMAFROST AND MARINE SEDIMENTS ARE HIGHLY SENSITIVE TO AMBIENT TEMPERATURE AND PRESSURE CHANGES

IN MARINE SEDIMENTS:

- WHEN OCEAN BOTTOM WATER TEMPERATURE RISES (DEGLACIACION) SOME METHANE CHANGES FROM HYDRATE FORM TO GASE FORM AND ESCAPES FROM THE SEAFLOOR INTO THE WATER COLUMN, AND FROM THERE TO THE ATMOSPHERE.
- HOWEVER, BECAUSE SEALEVEL CHANGES ACCORDING TO CLIMATIC CHANGES, THE INFLUENCE OF PRESSURE HAS TO BE TAKEN INTO ACCOUNT.


Anaerobic oxidation of methane (AOM) via sulfate reduction.

Microbial consortia of **methanotrophic archaea** and **sulfate-reducing bacteria** identified on gas hydrate-bearing samples (Boetius et al. 2000).

The archaea oxidize methane:



And sulfate reducing bacteria may act as indicated by the reaction:



Authigenic carbonate precipitation

Methane-derived Calcium Carbonate.




Bicarbonate
from bacterial action

alkaline
seawater

calcium ions
present in seawater

calcium
carbonate deposited



Bathymodiolus heckerae mussel beds. (A) Juvenile and adult mussels at Marker 'E'. (B) Dead mussels and octopus. (C) Extensive bed of live mussels of relatively uniform size, partially covered by bacterial mats, at Marker 'B'. (D) Dead mussels at the eastward periphery of Marker 'B'. (E) Mussels with a chiridotid holothurian and *Alvinocaris* sp. (F) Mussels with *Alvinocaris* sp. And ophiuroids. Scale bars: A-D : 10 cm, E; F : 5 cm.


Van Dover et al. (2003). Deep Sea Research

FORAMINIFERA (Marine zooplankton)

- **Protists: Unicellular marine organisms**
- **They live in all marine environments**
- **They have existed since the Cambrian**
- **Size: from 0.01 to 5 mm (> 50 mm)**

Shell composition:

- Agglutinated
- Tectine
- **Calcium Carbonate (CaCO_3)**



- Foraminifera form their skeleton in **isotopic equilibrium** with ambient seawater
- The isotopic composition of the skeleton of benthic foraminifera reflects the **micro-habitat** where they live


The $\delta^{13}\text{C}$ of foraminifera depends on the $\delta^{13}\text{C}$ of the Dissolved Inorganic Carbon (DIC) of the water in which they live:

Normal marine environment: $\delta^{13}\text{C}_{\text{DIC}} \approx 0\text{\textperthousand}$


(Peterson e Fry, 1987)

Methane seeps environment: $\delta^{13}\text{C}_{\text{DIC}} \approx -30/-60\text{\textperthousand}$


(Aharon *et al.*, 1992)




Negative excursions of $\delta^{13}\text{C}$ in marine sediments during warm periods can be explained ONLY with massive emissions of Methane from the seafloor




The Clathrate Gun Hypothesis




Methane hydrate dissociation as source of submarine slope instability




GLOBAL WARMING




THE STOREGGA SLIDE ON THE NORWEGIAN CONTINENTAL MARGIN(7000 yr)



Courtesy Petter Bryn





Courtesy Petter Bryn





Norsk Hydro
E&D Norway

Geophysical Operations

THE STOREGGA SLIDE CAUSED A HUGE TSUNAMI




The effect of gas hydrates





Mienert et al., 2005

Change in theoretical thickness of hydrate stability zone due to sea level change (small, negative) and bottom water temperature (locally large, positive). Biggest effect near upper headwall in Storegga.






THE NORTH SVALBARD SLIDE





Vanneste et al., EPSL, 2006

BLAKE RIDGE




(Dillon et al., 1998)

AT THE EDGE OF SCIENCE




4 scientific books



edited by J.-P. Honnorez & J. Mienert

Geological Society
Special Publication
No. 137

Published by The Geological Society

1998

COASTAL SYSTEMS AND CONTINENTAL MARGINS




Natural Gas Hydrate

In Oceanic and Permafrost Environments

Edited by Michael D. Max



Kluwer Academic Publishers

2000Charles K. Paull
William P. Dillon
Editors**2001****2003**

In Italiano

Energia immensa e sfida
ambientale. Gli idrati del
metanodi Carlo Giavarini (Autore)
Università La Sapienza,
2007

Fantascienza

Il quinto giorno (Der
Schwarm)Franz Schatting
2006