

**University of Trieste: GLOBAL CHANGE ECOLOGY a.a.
2021-2022**

**Conservation & Management in Marine Protected
Areas**


Dr. Stanislao Bevilacqua (sbevilacqua@units.it)

Effects of protection



Sheltering


This occurs when one or more target species increase their abundance, size or biomass within the protected areas with respect to fished areas.



Spillover


Buffering

This occurs when one or more target species exhibit less steep seasonal and/or interannual fluctuations within the protected area. Complex causes...reduction of post-recruitment mortality, increase of larval mortality (high density of predators)



Cascading effects

This occur when one or more target species have specific ecological role in stucturing marine communities. Protection, by increasing the abundance of this species allow them maintaning their role in controlling lower trophic levels, triggering cascading effects.




Effects on fish fauna


Giakoumi et al. 2017

a) Fully protected area

b) Partially protected area



Comparing effects between fish and invertebrates





Halpern, 2003

89 MPAs.


Density, size, biomass and diversity of fish fauna were significantly higher within than outside the reserve. Benthic invertebrates, however, showed significant difference only for density and size

Effects on different ecological compartments


Smallhorn-West et al. 2020



Relationship with reserve size





Size again...



Using 58 datasets from 19 European marine reserves, they showed that reserve size and age do matter: Increasing the size of the no-take zone increases the density of commercial fishes within the reserve compared with outside. Moreover, positive effects of marine reserve on commercial fish species and species richness are linked to the time elapsed since the establishment of the protection scheme. (Claudet et al, 2008)

Trophic cascades





Guidetti, 2006. *Ecol Appl*


Predation rates within reserves can be much more intense than outside

Increase of sea urchin predators due to protection reflects in decrease of sea urchins population within reserve boundaries, and the ensuing decrease of overgrazed substrates (Guidetti et al. 2008)


Effects on target species





Appolloni et al., 2017.



Maintainance of depth structure in fish assemblages. Abundance-biomass patterns typical of healthy conditions




Effects on fragile organisms





Diving frequentation in submarine caves. Effects on benthic invertebrates. (Guarnieri et al., 2012)

Mediterranean MPAs – subtidal rocky reefs




Fish biomass is significantly higher in well-enforced MPAs. Also, macroalgal stands (erect and canopy-forming species strongly varied, but were not related to protection. (Sala et al., 2012)



However, macroalgal stands were associated to low herbivore (sea urchins) pressure.

MPAs and resilience: a manipulative experiment



Date mussel (*Lithophaga lithophaga*) fishery

Banned in 1998 in Italy and in 2006 in EU
Caused the destruction of tens of km² of rocky bottoms in the Mediterranean, and especially in Italy, Croatia, Albania, Greece
Fishermen destroy the rocky surface, and everything living on the substrate, to reach the endolithic bivalve for collection
Still practiced, although illegal; costs of date mussels on the black market can range between 60-80 euros per Kg

Full protection



Simulating disturbance


Unprotected



Comparing trends in recovery



H₀: no difference in recovery between the no-take zone and controls



Temporal patterns of recovery



Human impact (date-mussel fishery) simulated within a no-take zone and 2 control areas (NW Mediterranean)

Recovery of macrobenthic assemblages followed during 20 months (5 times of sampling) in disturbed plots

Filled symbols = disturbed plots; empty symbols = undisturbed plots

Bevilacqua et al., 2006. J Animal Ecol




Recovery at the no-take zone was faster than at the unprotected control areas

Sea urchins


ANOVA

Source of variation	df	SS	MS	F	F versus
Time = Ti	2	0.08	0.04		
Location = Lo	2	1402	7.01	12086*	Ti x Lo
Controls = Cs	1	0.85	0.85	0.988ns	Ti x Cs
P-v-Cs	1	1317	1317	22706***	Residual
Ti x Lo	4	233	0.58	1.289ns	Residual
Ti x Cs	2	1.71	0.86	2.263ns	Res Cs
Ti x P-v-Cs	2	0.62	0.31	0.689ns	Residual
Residual	171	7697	0.45		
Res Cs	114	4349	0.38		
Res P	57	3348	0.59		

A. lixula




P. lividus






Does protection beget stability?

The MPA of Torre Guaceto (SE Adriatic Sea), instituted in 1991 and embedded into a human-dominated landscape, is a rare example of well-managed MPA where an adequate enforcement determined target fish recovery



This MPA provided the opportunity to follow the effects of protection on the stability of subtidal benthic assemblages, through the comparison of protected and unprotected locations, from 2002 to 2008

Protection, stability, and heterogeneity



SUBTIDAL ROCKY REEFS

The structure of subtidal sessile assemblages showed larger fluctuations outside the marine protected area than within the no-take zone where, in contrast, assemblage structure showed high temporal homogeneity.



Buffering effects on seagrass decline





Table 6. Classification of the status of *P. oceanica* beds based on shoot density following Pergent et al. [54].

Location	Patch	2006	2007	2008	2009
P1	1	undisturbed	disturbed	Undisturbed	undisturbed
P1	2	undisturbed	undisturbed	Undisturbed	undisturbed
P2	1	undisturbed	disturbed	Undisturbed	undisturbed
P2	2	undisturbed	undisturbed	Undisturbed	undisturbed
C1	1	undisturbed	very disturbed	very disturbed	disturbed
C1	2	undisturbed	very disturbed	very disturbed	undisturbed
C3	1	disturbed	undisturbed	Disturbed	Disturbed
C3	2	undisturbed	very disturbed	very disturbed	Disturbed

Seagrass beds under reduction in the area due to general increase in sedimentation rates and turbidity. However, the decline is less steep within the no-take areas, where additional direct human impacts (e.g., anchoring) are alleviated or excluded.



Further evidence




unprotected



Higher spatial heterogeneity, high temporal variability, decrease in canopy cover



ROCKY INTERTIDAL




Fraschetti et al., 2012. Mar Ecol Progr Ser



protected

Low spatial heterogeneity, high stability in canopy cover and associated understory assemblages



Fraschetti et al., 2012. Mar Ecol Progr Ser

Factors limiting protection effectiveness

Environmental

Poor recruitment from El Nino
(Preuss et al. 2009; Ferraris et al. 2005)

Environmental fluctuations
(Preuss et al. 2009; Powel et al. 2016)

Eutrophication (Moore et al. 2013)

Confounding habitat effects
(Dumas et al. 2010)

Discharge from river mouth
(Jupiter and Egli 2011)

Cyclone (Thiault et al. 2019)

Study design

Spillover into control sites
minimizing impact (Berdach 2003;
Ferraris et al. 2005; Preuss et al. 2009)

Habitat differences between
control and MPA sites (Wantiez et al.
1997; Jupiter et al. 2012)

Incorrect technique for question
(Jupiter et al. 2013)

Biological

Larval dispersal (Preuss et al. 2009)

Density dependent recruitment
(Dumas et al. 2012)

High natural variability
(Kulbicki et al. 2007)

Increased coral abundance attracts
Crown of thorns starfish
(Clements and Hay 2017)

Crown of thorns outbreak
(Thiault et al. 2019)

Low overall abundance of target
organisms (Dumas et al. 2010)

Complex life histories
(Dumas et al. 2010)

Changing predator dynamics (Goetz and
Fullwood 2013; Dell et al. 2015; Powel
et al. 2016))

Social

Insurmountable social barriers
(Bartlett et al. 2009b)

Poacher aggression
(Lalavanua et al. 2014)

Low overall fishing pressure
(Berdach 2003; Carassou et al. 2013)

Reserve design

Small reserve size (Preuss et al.
2009; Dumas et al. 2010;
Jupiter and Egli 2011)

Proximity to human populations
(Preuss et al. 2009; D'agata et al. 2016)

Insufficient time
(Dumas et al. 2010)

Unproductive habitat
(Preuss et al. 2009)

Poor visibility from village
(Jupiter and Egli 2011)

Management

Poaching/lack of compliance
(Bartlett et al. 2009b; Jupiter and Egli
2011; Moore et al. 2013; Lalavanua
et al. 2014; Albert et al. 2016;
Peters 2017; Thiault et al. 2019)

Overharvest of periodic closures
(Goetz et al. 2017)

Short periodic closure recovery time
(Jupiter et al. 2012; Goetz et al. 2015;
Goetz et al. 2016)

Smallhorn-West et al. 2020

