



**993SM - Laboratory of
Computational Physics
lecture 5
March 30, 2022**

Maria Peressi

Università degli Studi di Trieste - Dipartimento di Fisica
Sede di Miramare (Strada Costiera 11, Trieste)

e-mail: peressi@units.it

tel.: +39 040 2240242

Numerical integration - I

M. Peressi - UniTS - Laurea Magistrale in Physics
Laboratory of Computational Physics - Unit V

- **deterministic methods in ID**
equispaced points (trapezoidal, Simpson...),
others...
- **Monte Carlo methods**
(acceptance-rejection, sample mean,
importance sampling...)

Error handling:

sample mean

block average


reduction of the variance

Deterministic methods

Deterministic methods

Start from the geometrical interpretation of a definite integral:

$$F = \int_a^b f(x) dx$$




Divide the integration interval into “small” intervals:

$$\Delta x = \frac{b - a}{n},$$

$$x_n = x_0 + n \Delta x.$$

(Note: n intervals $\Leftrightarrow n + 1$ points)

Deterministic methods: rectangular rule



with error:
 $\mathcal{O}(h^2 f'), \propto 1/n^2$

applied iteratively over consecutive intervals:

$$F_n = \sum_{i=0}^{n-1} f(x_i) \Delta x.$$

with a total error:
 $\mathcal{O}(h f'), \propto 1/n$

: The rectangular approximation for $f(x) = \cos x$ for $0 \leq x \leq \pi/2$.


Deterministic methods: rectangular rule - error

$$I = \int_0^{\pi/2} \cos(x) dx = 1$$

$$F_n = \frac{\pi}{2n} \sum_0^{n-1} \cos x_i; \quad x_i = i \frac{\pi}{2n}$$

$$\Delta_n = F_n - I$$

n	F_n	Δ_n
2	1.34076	0.34076
4	1.18347	0.18347
8	1.09496	0.09496
16	1.04828	0.04828
32	1.02434	0.02434
64	1.01222	0.01222
128	1.00612	0.00612
256	1.00306	0.00306
512	1.00153	0.00153
1024	1.00077	0.00077



Rectangular approximations of the integral of $\cos x$ from $x = 0$ to $x = \pi/2$ as a function of n , the number of intervals. The error Δ_n is the difference between the rectangular approximation and the exact result of unity. Note that the error Δ_n decreases approximately as n^{-1} , that is, if n is increased by a factor of 2, Δ_n decreases by a factor 2.

Deterministic methods: generalities

- sum values of $f(x_i)$ with $x_i \in [a, b]$
- we want to have $F = \int_a^b f(x)dx$
as accurate as possible but with the
minimum number of calculations of $f(x_i)$

OK simple algorithms, but if the number of calculations is too high, improve the algorithm!

Deterministic methods: trapezoidal rule

In one interval:


$$\int_{x_i}^{x_{i+1}} f(x) dx = h \left[\frac{1}{2} f_i + \frac{1}{2} f_{i+1} \right]$$

with error:

$$\mathcal{O}(h^3 f''), \propto 1/n^3$$

Applied iteratively over consecutive intervals:


$$F_n = \left[\frac{1}{2} f(x_0) + \sum_{i=1}^{n-1} f(x_i) + \frac{1}{2} f(x_n) \right] \Delta x.$$



with a total error:

$$\mathcal{O}(h^2 f''), \propto 1/n^2$$

Deterministic methods: Simpson's rule



Parabolic interpolation procedure
between triplets of adjacent points

In one interval:

$$\int_{x_i}^{x_{i+2}} f(x) dx = h \left[\frac{1}{3} f_i + \frac{4}{3} f_{i+1} + \frac{1}{3} f_{i+2} \right] + \mathcal{O}(h^5 f^{IV}) \quad (\text{error} \propto 1/n^5)$$

Iteratively applied to the whole interval of integration (**odd** number of points!):

$$\int_{x_0}^{x_n} f(x) dx = h \left[\frac{1}{3} f_0 + \frac{4}{3} f_1 + \frac{2}{3} f_2 + \frac{4}{3} f_3 + \dots + \frac{2}{3} f_{n-2} + \frac{4}{3} f_{n-1} + \frac{1}{3} f_n \right] + \mathcal{O}(h^4 f^{IV}) \quad (\text{error} \propto 1/n^4)$$

Errors in deterministic methods

Error estimate for numerical integration with deterministic methods

$$\int f(x)dx = F_n + error$$

How to evaluate the error? Consider the Taylor expansion of the integrand function and then integrate:

$$f(x) = f(x_i) + f'(x_i)(x - x_i) + \frac{1}{2}f''(x_i)(x - x_i)^2 + \dots, (*)$$

$$\int_{x_i}^{x_{i+1}} f(x) dx = f(x_i)\Delta x + \frac{1}{2}f'(x_i)(\Delta x)^2 + \frac{1}{6}f''(x_i)(\Delta x)^3 + \dots(**)$$

$$\Delta x \equiv x_{i+1} - x_i$$

Error estimate for numerical integration: Rectangular approximation

$$\int_{x_i}^{x_{i+1}} f(x) dx \approx f(x_i) \Delta x$$

Compare with (**):

$$\int_{x_i}^{x_{i+1}} f(x) dx = f(x_i) \Delta x + \frac{1}{2} f'(x_i) (\Delta x)^2 + \frac{1}{6} f''(x_i) (\Delta x)^3 + \dots$$

error

(leading order in Δx)

For n intervals ($\Delta x = (b - a)/n$): error is $n(\Delta x)^2 \sim 1/n$

Error estimate for numerical integration: Trapezoidal approximation

$$\int_{x_i}^{x_{i+1}} f(x) dx \approx \frac{1}{2} [f(x_{i+1}) + f(x_i)] \Delta x$$

\downarrow \downarrow

$$f(x_{i+1}) \approx f(x_i) + f'(x_i)\Delta x + \frac{1}{2}f''(x_i)\Delta x^2 + \dots$$
$$\approx \frac{1}{2} \left[2f(x_i) + f'(x_i)\Delta x + \frac{1}{2}f''(x_i)\Delta x^2 + \dots \right] \Delta x$$

Compare with (**):


$$\int_{x_i}^{x_{i+1}} f(x) dx = f(x_i)\Delta x + \frac{1}{2}f'(x_i)(\Delta x)^2 + \frac{1}{6}f''(x_i)(\Delta x)^3 + \dots$$

error

(leading order in Δx)

For n intervals: error is $n(\Delta x)^3 \sim 1/n^2$

Numerical integration - deterministic methods: comparison of errors in ID



Deterministic methods - I

We use a piecewise polynomial interpolation:


- constant interpolation → 1 point → rectangular rule
- linear interpolation → 2 points → trapezoidal rule
- parabolic interpolation → 3 points → Simpson's rule
- ...
- higher-order polynomial → many points →

NOT CONVENIENT!

Warning: using higher degrees does not always improve accuracy!

(see also: Runge phenomenon (polynomial interpolation, oscillation at the edges of an interval), Gibbs phenomenon ...)

Deterministic methods - II



• $(x_i, f(x_i))$

Warning:

using high-order piecewise polynomial interpolation: possible strong oscillations between consecutive $(x_i, f(x_i))$, giving a bad interpolation of $f(x)$.

Here: $f(x)$ step function; - linear interp.; - cubic spline


Example: `gnuplot> p 'data', " smooth cspline`

Monte Carlo methods

Monte Carlo methods:

“acceptance-rejection” or “hit or miss”
(to calculate areas)

which is A_{pond} ?



- enclose the pond in a box of Area A_{box}
- throw pebbles uniformly and randomly in the box
- count the number of pebbles felt in the pond with respect to the number felt in the box
- Assuming a uniform distribution, the number of pebbles falling into the ponds is proportional to the area of the pond:


$$\frac{N_{\text{pond}}}{N_{\text{pond}} + N_{\text{box}}} = \frac{A_{\text{pond}}}{A_{\text{box}}}$$
$$\Rightarrow A_{\text{pond}} = \frac{N_{\text{pond}}}{N_{\text{pond}} + N_{\text{box}}} A_{\text{box}}$$

Monte Carlo methods:

“acceptance-rejection” or “hit or miss”
(to calculate areas)

$$\pi = ???$$

N random points in the unit square
- coordinates x_i, y_i -




Then, the number of points N_c lying within the quarter circle (i.e. fulfilling the relation $x^2 + y^2 \leq 1$) is compared to the total number N of points and the fraction will give us an approximate value of π :

$$\pi(N) = 4 \frac{N_c(N)}{N}$$

Monte Carlo methods:

“acceptance-rejection” or “hit or miss”
(to calculate definite integrals)

$$\int W(x) dx = ?$$



For $W(x)$ positive in the integration interval, the value of the area under $W(x)$ can be obtained by producing random points (i.e. (x,y) random pairs) uniformly distributed in a rectangle containing $W(x)$.


For each point (x,y) compare y with $W(x)$: if $y < W(x)$, the point is accepted. The area under $W(x)$ is the number of points accepted divided by the total number of points generated and multiplied by the area of the rectangle.

(remember: also used to generate random numbers x_i distributed according $W(x)$)

Other simple Monte Carlo methods

We can always write:

$$I = \int_a^b f(x) dx = (b - a) \langle f \rangle$$




i.e., the value of the integral of $f(x)$ between a and b equals the length of the interval $(b-a)$ times the average value of the function $\langle f \rangle$ over the same interval.

(If $f:[a,b] \rightarrow \mathbb{R}$ is a continuous function, then there exists a number c in $[a,b]$ such that $f(c) = \langle f \rangle$ (mean value theorem for integration))

how to estimate $\langle f \rangle$ efficiently and accurately?

A simple Monte Carlo method: “sample mean”



$$I = \int_a^b f(x) dx = (b - a) \langle f \rangle$$

The **sample mean** can be calculated by sampling the function (*if smooth enough...*) with a sequence of N uniform random numbers in [a,b]:

$$\langle f \rangle \approx \frac{1}{N} \sum_{i=1}^N f(x_i)$$

$$\int_a^b f(x) dx \approx (b - a) \frac{1}{N} \sum_{i=1}^N f(x_i) = (b - a) \langle f \rangle$$

Monte Carlo methods: error estimate

Example: MC estimate of π (exact value known)

We can use either acceptance-rejection or sample mean method: $I = 4 \int_0^1 \sqrt{1-x^2} = \pi = 3.1416\dots$

Since we know the “exact” result I , we can calculate the **error** in two ways:

1) the **actual error** from the difference with respect to the exact value:

$$\Delta_n = |F_n - I| \quad \text{with} \quad F_n = (b-a) \frac{1}{n} \sum_{i=1}^n f(x_i), \quad x_i \text{ random}$$

2) the numerical error from the **variance of the data** $\{f(x_i)\}$:

$$\sigma^2 = \langle f^2 \rangle - \langle f \rangle^2,$$

where


$$\langle f \rangle = \frac{1}{n} \sum_{i=1}^n f(x_i) \quad \text{and} \quad \langle f^2 \rangle = \frac{1}{n} \sum_{i=1}^n f(x_i)^2$$

Monte Carlo methods: error estimate

Results:

$$I = 4 \int_0^1 \sqrt{1-x^2} = \pi = 3.1416\dots$$

n	F_n	actual error	σ_n
10^2	3.0692	0.0724	0.85550
10^3	3.1704	0.0288	0.8790
10^4	3.1489	0.0073	0.8850



1) the **actual error** Δ_n decreases as $1/n^{1/2}$

2) the numerical error from the **variance of the data**, σ_n , is roughly constant and is much larger than the actual error

what is the correct error estimate?

Monte Carlo methods: error estimate

...typically you do not know which is the “actual error” (you do not know the “true” value and you cannot compare your result with that!)...
but **we would like to give an error to our numerical estimate...**
(to which extent is our numerical estimate reliable?)

Two methods to estimate the error numerically
from the variance of the data
(“reduction of variance”):

I) average of the averages

II) block average

MC error handling: method I

“average of the averages”

make additional runs of n trials each.

Let M_α be the average of each run :

run α	M_α	actual error
1	3.1489	0.0073
2	3.1326	0.0090
3	3.1404	0.0012
4	3.1460	0.0044
5	3.1526	0.0110
6	3.1397	0.0019
7	3.1311	0.0105
8	3.1358	0.0058
9	3.1344	0.0072
10	3.1405	0.0011

one run $\equiv n = 10^4$ trials each

Examples of Monte Carlo measurements of the mean value of $f(x) = 4\sqrt{1-x^2}$ in the interval $[0, 1]$. A total of 10 measurements of $n = 10^4$ trials each were made. The mean value M_α and the actual error $|M_\alpha - \pi|$ for each measurement are shown.

$$\text{Calculate: } \sigma_m^2 = \langle M^2 \rangle - \langle M \rangle^2 \quad \text{with } \langle M \rangle = \frac{1}{m} \sum_{\alpha=1}^m M_\alpha, \quad \langle M^2 \rangle = \frac{1}{m} \sum_{\alpha=1}^m M_\alpha^2$$
$$\implies \sigma_m = 0.0068$$

σ_m is consistent with the results for the actual errors

$\sigma = 0.0073$ (see slide 26)

MC error handling: method II

“block averages”

Instead of doing additional measurements, divide them into “s SUBSETS” and let S_k be the average within each subset :

subset k	S_k
1	3.14326
2	3.15633
3	3.10940
4	3.15337
5	3.15352
6	3.11506
7	3.17989
8	3.12398
9	3.17565
10	3.17878

The variance associated to the average of the subsets $\sigma_s^2 = \langle S^2 \rangle - \langle S \rangle^2$ gives $\sigma_s = 0.025$, but

σ_s/\sqrt{s} , which for our example is approximately $0.025/\sqrt{(10)} \approx 0.008$.

is consistent with the actual error

$\sigma = 0.0073$ (see slide 26)

Monte Carlo methods:

error estimate - variance reduction

summary

$$\sigma_n / \sqrt{n} \approx \sigma_m \approx \sigma_s / \sqrt{s}$$

from the variance of
the whole set of data

Note: for
uncorrelated data !

(proof)

the variance
of the

average of
the averages

from the variance
of the
block averages

the most convenient!
but: change block size
and check that
it does not change

Monte Carlo methods: summary

We have introduced :

* “acceptance-rejection”

* “sample mean” to estimate $\langle f \rangle \approx \frac{1}{N} \sum_{i=1}^N f(x_i)$

both OK for smoothly varying functions, but
not very efficient for rapidly varying functions

How to improve the efficiency of MC integration?

A trick for numerical integration: “reduction of variance”

(Note: same word, but different meaning w.r.t. previous slides on error handling)

Given a function $f(x)$ to integrate, suppose that $g(x)$ exists, whose integral is known and such that:

$$|f(x) - g(x)| \ll \varepsilon$$

Therefore:


$$F = \int_a^b f(x) dx = \int_a^b ((f(x) - g(x)) + g(x)) dx = \int (f(x) - g(x)) dx + \int g(x) dx$$




easy to calculate

Another simple Monte Carlo method: “importance sampling”

Mean value: easy to calculate for smoothly varying functions.
But not for functions rapidly varying.



smooth function



function with singularity

How to manage such cases?

Another simple Monte Carlo method: “importance sampling”

Mean value: easy to calculate for smoothly varying functions.

Idea: in order to calculate:

$$\langle f \rangle \approx \frac{1}{N} \sum_{i=1}^N f(x_i)$$

consider a **distribution function** $p(x)$ **easy to integrate analytically and close to** $f(x)$:

$$F = \int_a^b f(x) dx = \int_a^b \left[\frac{f(x)}{p(x)} \right] p(x) dx = \left\langle \frac{f(x)}{p(x)} \right\rangle \int_a^b p(x) dx$$

where $\left\langle \frac{f(x)}{p(x)} \right\rangle \approx \frac{1}{N} \sum_{i=1}^N \left[\frac{f(x_i)}{p(x_i)} \right]$

(particular case:
uniform distrib.
 $p(x)=1/(b-a)$...)

with $\{x_i\}$ distributed according to $p(x)$

Monte Carlo methods: “importance sampling”

Calculate:

$$F = \int_0^1 e^{-x^2} dx = 0.746824\dots$$

with “sample mean” with random numbers with uniform distribution or using the “importance sampling” with $p(x) = e^{-x}$

	$p(x) = 1$	$p(x) = Ae^{-x}$
n (trials)	4×10^5	8×10^3
F_n	0.7471	0.7469
σ	0.2010	0.0550
σ/\sqrt{n}	3×10^{-4}	6×10^{-4}
Total CPU time (s)
CPU time per trial (s)


← more efficient !

discover it yourself!

(use: call `cpu_time(...)`)

(pay attention to the normalization of $p(x)$...)

$\text{error}(\text{MC}) \sim I/\sqrt{N} \Rightarrow \text{see } \log(\text{error}) \text{ vs. } \log(N)$
but with different prefactors
for sample means vs importance sampling




Choice of the importance sampling function

Ex. 2

$$F = \int_0^1 e^{-x^2} dx.$$

$$F = \int_a^b f(x) dx = \int_a^b \left[\frac{f(x)}{p(x)} \right] p(x) dx = \left\langle \frac{f(x)}{p(x)} \right\rangle \int_a^b p(x) dx$$



(pay attention to the normalization of p(x)...)

Some programs:

on <https://moodle2.units.it/>


int.f90 (trapezoidal and Simpson integration) for Ex. 1

pi.f90 (Montecarlo integration for calculation of π) for Ex. 3

for the other exercises: write yourself the code!

↑
Ex. 2 and 4: homework!

Ex. 2: error(MC) $\sim 1/\sqrt{N}$: “true” error and statistical error



(credits: G. Lautizi, a.y. 2019-20)

$\text{error}(\text{MC}) \sim 1/\sqrt{N} \Rightarrow$ see $\log(\text{error})$ vs. $\log(N)$

