


Lesson 5

Cellular Chemistry, Reaction Thermodynamics and Enzymes



Metabolism

- The cell is an ideal continuous chemical reactor
- The ensemble of cellular chemical reactions = **metabolism**
- **Anabolic reactions** = build up molecules (*e.g.*, condensation)




- **Catabolic reactions** = break down molecules (*e.g.*, hydrolysis)





Factory, Maurice Utrillo, 1923

Condensation & hydrolysis



Condensation:


Covalent bonds form
with loss of water
Build molecules
Anabolic



Hydrolysis:

Covalent bonds break
with addition of water
Break down molecules
Catabolic


Metabolism & Energy



The metabolic map

Dots = molecules

Lines = chemical reactions
making up the metabolism of
the cell



Free energy difference


- Reactions are governed by **free energy** (usable energy) **G**

Reagents \rightleftharpoons Products

- What really matters is the **free energy difference** $\Delta G = \sum G_P - \sum G_R$
- ΔG stems from a fundamental law of thermodynamics


$$\Delta G = \Delta H - T\Delta S$$

- **H** = **Enthalpy** = total energy
- **T** = **Temperature**
- **S** = **Entropy** = useless energy




Free energy difference

- $\Delta G < 0$ ($G_P < G_R$) \rightarrow Energy released, reaction proceeds (**thermodynamically spontaneous, exergonic**)



Free energy difference

- $\Delta G < 0$ ($G_P < G_R$) \rightarrow Energy released, reaction proceeds (thermodynamically spontaneous, exergonic)
- $\Delta G > 0$ ($G_P > G_R$) \rightarrow Reactions requires energy to proceed (thermodynamically non-spontaneous, endergonic)




Free energy difference

- $\Delta G < 0$ ($G_P < G_R$) \rightarrow Energy released, reaction proceeds (thermodynamically spontaneous, exergonic)
- $\Delta G > 0$ ($G_P > G_R$) \rightarrow Reactions requires energy to proceed (thermodynamically non-spontaneous, endergonic)
- $\Delta G = 0$ ($G_P = G_R$) \rightarrow Chemical equilibrium ($R \rightarrow P = P \rightarrow R$)



$$K_{eq} = [B]/[A]$$

$$\Delta G = -RT \ln [B]/[A]$$



$$\Delta G = -RT \ln K_{eq}$$


$$R = 8.314 \text{ J}/(\text{mol K}) = 1.987 \text{ cal}/(\text{mol K})$$

Going to work in the cell factory

- Cells transfer energy from food molecules to run their processes
 - Synthesis
 - Complex molecules (DNA, proteins)
 - Organized structures (organelles)
 - Organization
 - Generation of specialized compartments to store and organize materials for specific tasks


Going to work in the cell factory

- Cells transfer energy from food molecules to run their processes
 - Synthesis
 - Complex molecules (DNA, proteins)
 - Organized structures (organelles)
 - Organization
 - Generation of specialized compartments to store and organize materials for specific tasks
- Creation of electrochemical gradients
 - Cells organize ions on either cell membrane sides to create electrochemical potentials
 - Mechanism that controls signaling and transfer of materials across the membranes




Going to work in the cell factory

- Cells transfer energy from food molecules to run their processes
 - Synthesis
 - Complex molecules (DNA, proteins)
 - Organized structures (organelles)
 - Organization
 - Generation of specialized compartments to store and organize materials for specific tasks
 - Creation of electrochemical gradients
 - Cells organize ions on either cell membrane sides to create electrochemical potentials
 - Mechanism that controls signaling and transfer of materials across the membranes
- **Transport and movement**
 - Cells transport molecules from one side of the cell to another
 - Some cells swim or crawl using special motility proteins



Where does cell energy really come from?

- Cells do not use the energy from food directly
- Adenosine triphosphate, the energy middleman




The ATP/ADP cycle

Simple molecules
glucose, amino acids, fatty acids

Catabolism – Exergonic reactions
transfer energy from complex
molecules to ATP

Anabolism – Endergonic reactions
transfer energy from ATP
to complex molecules


Complex molecules
polycarbs, proteins, lipids







C. Missig et US UP

The ATP/ADP cycle – energy coupling

- Cells make and break ATP all the time
- In endergonic reactions
 - Energy is provided by ATP hydrolysis to ADP and P_i
- In exergonic reactions
 - Energy is captured to restore ATP by ADP and P_i condensation




As easy as 1,2,3 – ATP, ADP and AMP





Free energy difference

- Even if a reaction is thermodynamically spontaneous ($\Delta G < 0$), it may not occur
 - It needs an “energetical push”
- **Activation energy E_a = energy barrier**





Enzymes

- Even if a reaction is thermodynamically spontaneous ($\Delta G < 0$), it may not occur
 - It needs an “energetical push”
- **Activation energy** E_a = energy barrier
- **Catalysts** = particular class of chemical substances that lower E_a and promote reactions
- **Enzymes** = biological catalysts (mostly proteins)




Enzymes

- There are approximately 1300 different enzymes found in the human cell
- Each enzyme catalyzes a specific chemical reaction
- **ENZYMES DO NOT CHANGE THE ΔG OF A REACTION BUT JUST SPEED UP THE REACTION RATE**
 - By breaking down the reaction into different steps, each of which has a low E_a




Enzymes




ENZYME CYCLE
(reversible)

Enzymes




ENZYME CYCLE
(reversible)

Enzymes




Enzymes




Enzyme specificity

Cellulose and starch are both glucose polymers



Cellulase breaks β -1,4 bonds in cellulose
People do not have this enzyme so we cannot digest grass!



Amylase breaks α -1,4 bonds in starch
People have this enzyme!




ENZYME SPECIFICITY

Metabolic pathways

- Cellular chemical reactions all catalyzed by enzymes
- Cellular chemical reactions are often linked into **pathways**
 - Ordered sequence of chemical reactions
- Pathways = the “cell production line”
 - Starting from point A → land up with a particular product that the cell really needs
- Reactions are organized into multistep pathways
- Cellular pathways are governed by internal feedback mechanisms
 - **Positive feedback** = make more product(s) along that particular pathway
 - **Negative feedback** = make less and/or stop producing product(s) along that particular pathways
- Cellular pathways can also be mastered by external control signals




Pathways and feedback loops




Positive feedback example

- Problem 1: when the cell gets a lot of product C it also needs a lot of product E
- Solution:
 - C (whatever the molecule is) speeds up the production of D
 - Increased D levels will increase the production of E



Negative feedback example

- Problem: the cell has enough D and now wants to produce E only by pathway 2
- Solution:
 - it turns off pathway 1
 - D inhibits production of B and hence of C
 - D is no longer produced via C
 - E is only produced via pathway 2



Real cellular pathway example

Phenylalanine metabolism and pathways
Every arrow = one or more enzymes!

