
Prof. Sabrina Pricl A.Y. 2023-2024

Lesson 1 Water, pH and buffers



- The fundamental molecule of life
- 60-95% of living human cells is H₂0
 - 55% in intracellular fluids
 - 45% divided between:
 - Plasma (8%)
 - Interstitial (between cells) and lymph (22%)
 - Connective tissue, cartilage and bones (15%)




Lymph (from Latin, lympha) is the fluid that flows through the lymphatic system, a system composed of lymph vessels (channels) and intervening lymph nodes whose function, like the venous system, is to return fluid from the tissues to the central circulation

- In biochemistry:
 - Transport medium across cell membranes
 - Body temperature maintenance
 - Solvent in the GI and excretion system


- In biochemistry:
 - Transport medium across cell membranes
 - Body temperature maintenance
 - Solvent in the GI and excretion system
- Healthy humans
 - Daily water intake/loss = 2L
 - Intake --> 45% from liquids, 40% from food and 15% chemical reactions
 - Loss -> 50% urines, 5% feces and 55% evaporation from lung and skin

- In biochemistry:
 - Transport medium across cell membranes
 - Body temperature maintenance
 - Solvent in the GI and excretion system
- Healthy humans
 - Daily water intake/loss = 2L
 - Intake --> 45% from liquids, 40% from food and 15% chemical reactions
 - Loss -> 50% urines, 5% feces and 55% evaporation from lung and skin
- Water balance must be preserved:
 - Loss >> intake --> dehydration
 - Intake >> loss --> edema




Water dissociation and pH

- $H_20 <-> H_30^+ + OH^-$
 - $[H_3O^+] = [OH^-] --> neutral$
 - $[H_3O^+] > [OH^-] --> acidic$
 - $[H_3O^+] = [OH^-] --> basic$
- $K_w = [H_3O^+] \times [OH^-] = 10^{-14} (25^{\circ}C)$
 - In human body @ 37° C --> $K_w = 2.4 \times 10^{-14}$



- pH = $-log[H_3O^+]$ --> in pure water pH = 7
- In human blood @ 37° C --> pH =7.4 --> [H_3O^{+}] = 3.98×10^{-8}
- $pK_w = 14$



Weak acids and bases

• In biology we have only weak acids and bases (incomplete dissociation)

$$CH_3COOH + H_2O <--> H_3O^+ + CH_3COO^-$$

$$NH_3 + H_2O < --> NH_4^+ + OH^-$$

$$K_a = [H_3O^+][CH_3COO^-]/[CH_3COOH]$$

$$K_b = [NH_4^+][OH^-]/[NH_3]$$

$$pK_a = -logK_a$$

$$pK_b = -logK_b$$


$$K_a \times K_b = K_w = 10^{-14} --> pK_a + pK_b = 14$$

CH₃COO⁻ conjugate base

NH₄⁺ conjugate acid

Buffers and pH control

 A solution that contains a conjugate acid-base pair of any weak acid or base in relative proportions to resist pH change when small amounts of either a (strong) acid or base are added is a buffer solution




- Stomach pH = 1-2
- GI tract pH = 8-9
- Blood pH = 7.4
 - Blood pH < 7.2 --> pathological condition = acidosis
 - Blood PH < 6.8 --> death
 - Blood pH > 7.6 --> pathological condition = alkalosis
 - Blood pH > 8.6 --> death

- Respiratory acidosis
 - Inefficient expulsion of CO₂, increased concentration of H₂CO₃, impaired-respiration pathologies (pneumonia, emphysema, asthma)


Respiratory alkalosis

 Excessive CO₂ removal, decreased concentration of H₂CO₃, hyperventilation


RESPIRATORY ACIDOSIS




RESPIRATORY ALKALOSIS



- Metabolic acidosis
 - Decreased concentration of HCO₃-, results from various kidney diseases, uncontrolled diabetes, or vomiting of non-acid fluids



- Metabolic alkalosis
 - Increased concentration of HCO₃-, results excessive vomiting of stomach acid



PROTEIN BUFFER SYSTEM

Protein buffer system helps to maintain acidity in and around cells

PHOSPHATE BUFFER SYSTEM

Phosphate buffer helps to maintain intracellular and urine pH

BICARBONATE BUFFER

- main extracellular buffer, main blood buffer
 - Helps in controlling CO₂ levels --> CO₂ + H₂O <--> H₂CO₃ <--> H⁺ + HCO₃
 - Coupled with CO₂ (blood) <--> CO₂ (lungs)

Buffer action and the pH of blood

- Normal blood pH = 7.4
 - Kept at this value by the buffering action of HCO3⁻, resulting from these two parallel physiological equilibria:
 - CO₂ + H₂O <--> H₂CO₃
 - H₂CO₃ <--> H⁺ + HCO₃⁻
- $K_{eq} = [H^+][HCO_3^-]/[CO_2]^* = 7.95 \times 10^{-7} --> pK_{eq} = 6.1$
- Rearranging:

$$[H^{+}] = K_{eq} \times [CO_{2}]/[HCO_{3}^{-}] --> -log[H^{+}] = -logK_{eq} -log[CO_{2}]/[HCO_{3}^{-}] --> pH = pK_{eq} + log[HCO_{3}^{-}]/[CO_{2}]$$

^{*} $[H_2O]$ = 55.6 M = const = included in the K_{eq} value

Quiz time

• A patient suffering from acidosis had a blood pH of 7.15 and a CO_2 concentration of 1.15 mM. If the reference range for pH = 7.4 are:

 $[HCO_3^-] = 22.0 - 26.0 \text{ mM (average} = 24 \text{ mM)}$

 $[CO_2] = 1.20 \text{ mM}$

and pK_{eq} for the bicarbonate buffer = 6.1:

- Q1. What was the patient's bicarbonate (HCO3⁻) concentration?
- Q2. What are the implications of this value to the buffer capacity of the blood?

Quiz time

R1.

pH = pK_{eq} + log[HCO₃⁻]/[CO₂] -->
$$7.15 = 6.1 + log[HCO3-]/(1.15 x 10-3)$$

$$10^{1.05} = [HCO_3^{-}]/(1.15 \times 10^{-3}) --> [HCO_3^{-}] = 12.9 \times 10^{-3} --> [HCO_3^{-}] = 12.9 \text{ mM}$$

R2.

Normal [HCO₃⁻] average value = 24 mM --> [HCO₃⁻] in patient lowered by 11.1 mM --> severely impaired buffer capacity --> any further, small acid production will have serious consequences for the patient