
Lesson 3 Nucleic acid polarity and structure



Macromolecules' Law & Order

- Macromolecules carry information as:
 - They have two ends (starting and terminal end)
 - They have a direction (from start to end)





Macromolecules' Law & Order

- Macromolecules carry information as:
 - They have two ends (starting and terminal end)
 - They have a direction (from start to end)
- Macromolecules possess ORDER and POLARITY
 - These two features constitute the FUNDAMEMTAL INFORMATION cellules can read and understand


Macromolecules' Law & Order

- Macromolecules carry information as:
 - They have two ends (starting and terminal end)
 - They have a direction (from start to end)
- Macromolecules possess ORDER and POLARITY
 - These two features constitute the FUNDAMEMTAL INFORMATION cellules can read and understand
- NUCLEIC ACIDS = carries of hereditary information
- PROTEINS = all other info/instructions







• Have two hands: 5' e 3




- Have two hands: 5' e 3' 5'P-S-P-S-P-S3' B1 B2 B3
- P-S = sugar-phosphate backbone joined by phosphodiester bonds



- Have two hands: 5' e 3' 5'P-S-P-S-P-S3' B1 B2 B3
- P-S = sugar-phosphate backbone joined by phosphodiester bonds
- 5' and 3' ends are CHEMICALLY DIFFERENT **ENDS** and cells can distinguish them from one another




- P-S = sugar-phosphate backbone joined by phosphodiester bonds
- 5' and 3' ends are **CHEMICALLY DIFFERENT ENDS** and cells can distinguish them from one another
- Polarity:
 - 5'end = P-5'C on sugar
 - 3'end = 3'OH on sugar



• Have two hands: 5' e 3'

5'P-S-P-S-P-S3' | | | B1 B2 B3


- P-S = sugar-phosphate backbone joined by phosphodiester bonds
- 5' and 3' ends are **CHEMICALLY DIFFERENT ENDS** and cells can distinguish them from one another
- Polarity:
 - 5'end = P-5'C on sugar
 - 3'end = 3'OH on sugar
- Base order along the polymer




• Have two hands: 5' e 3' **5'P-S-P-S-P-S3'**

5'P-S-P-S-P-S3' | | | | B1 B2 B3

- P-S = sugar-phosphate backbone joined by phosphodiester bonds
- 5' and 3' ends are **CHEMICALLY DIFFERENT ENDS** and cells can distinguish them from one another
- Polarity:
 - 5'end = P-5'C on sugar
 - 3'end = 3'OH on sugar
- Base order along the polymer
- 3' is the last base added (any new incoming base will be added to 3'OH)



Nucleic acids



Elements of Chemical and Molecular Biology – Lesson 3

• S-P backbone is not written, just the bases + polarity


- S-P backbone is not written, just the bases + polarity
- ALWAYS write 5' and 3' on each nucleic acid strand

 $5'B_1B_2B_3B_4B_5B_63' = 5'GAATCC3'$

- S-P backbone is not written, just the bases + polarity
- ALWAYS write 5' and 3' on each nucleic acid strand


 $5'B_1B_2B_3B_4B_5B_63' = 5'GAATCC3'$

- Base order = INFORMATION
- Polarity = 5' and 3' ends: shows
 - First to last nucleotide added
 - Direction to read information




Get to know your DNA structure


- DNA has a right-handed double-helix (DH) structure (2 nm diameter)
 - Ss and Ps lie on the outside of the helix (backbone)
 - Bs are stacked in the interior, in pairs
 - B pairs (BPs) are bound to each other by Hbonds
 - Every BP in the DH is separated from the next base pair by 0.34 nm (axial rise)
 - There are 10.4 BPs per helical trun
 - The distance to complete one helical turn (helical pitch) = 3.4 nm




- DNA has a right-handed double-helix (DH) structure (2 nm diameter)
 - Ss and Ps lie on the outside of the helix (backbone)
 - Bs are stacked in the interior, in pairs
 - B pairs (BPs) are bound to each other by Hbonds
 - Every BP in the DH is separated from the next base pair by 0.34 nm (axial rise)
 - There are 10.4 BPs per helical trun
 - The distance to complete one helical turn (helical pitch) = 3.4 nm
- The two strands of the helix run in opposite directions
 - the 5' carbon end of one strand will face the 3' carbon end of its matching strand
- This is referred to as **antiparallel orientation**
 - Key DNA PROPERTY (for DNA replication and in many nucleic acid interactions)



- Only certain types of base pairing are allowed:
- Specifically: A can only pair with T and G can only pair with C (base complementary rule)




- Only certain types of base pairing are allowed:
- Specifically: A can only pair with T and G can only pair with C (base complementary rule)
- In other words, the DNA strands are complementary to each other
 - If the sequence of one DNA strand is
 5'AATTGGCC3', the complementary strand would have the sequence 3'TTAACCGG5'




 RNA (ribonucleic acid) → mainly involved in the process of protein synthesis under the direction of DNA

- RNA (ribonucleic acid) → mainly involved in the process of protein synthesis under the direction of DNA
- RNA is usually single-stranded
- The RNA four nitrogenous bases are A, U, G, and C




- RNA (ribonucleic acid) → mainly involved in the process of protein synthesis under the direction of DNA
- RNA is usually single-stranded
- The RNA four nitrogenous bases are A, U, G, and C
- There are four major types of RNA
 - messenger RNA (mRNA), ribosomal RNA (rRNA), transfer RNA (tRNA), and microRNA (miRNA) (more later)



- The mRNA carries the message from DNA
 - If a cell requires a certain protein to be synthesized, the gene for this product is "turned on" and the messenger RNA is synthesized in the cellular nucleus (more later)

- The mRNA carries the message from DNA
 - If a cell requires a certain protein to be synthesized, the gene for this product is "turned on" and the messenger RNA is synthesized in the cellular nucleus (more later in the course)
- The RNA base sequence is complementary to the coding sequence of the DNA which it has been copied from but

- The mRNA carries the message from DNA
 - If a cell requires a certain protein to be synthesized, the gene for this product is "turned on" and the messenger RNA is synthesized in the cellular nucleus (more later in the course)
- The RNA base sequence is complementary to the coding sequence of the DNA which it has been copied from but
- If the DNA strand to be copied (template strand) has a sequence 3'TACGTCTA...3', the sequence of the complementary mRNA strand (transcript) is 5'AUGCAGAU...3'



Note: template strand reading is $3' \rightarrow 5'$ mRNA synthesis is $5' \rightarrow 3'$ (more later)