



Università degli studi di Trieste

LAUREA MAGISTRALE IN GEOSCIENZE

Classe Scienze e Tecnologie Geologiche

Curriculum: Esplorazione Geologica

Anno accademico 2023 - 2024

Analisi di Bacino e Stratigrafia Sequenziale (426SM)

Docente: Michele Rebesco



Analisi di Bacino e Stratigrafia Sequenziale (426SM)

Unit 1.2: Methods

Docente: **Riccardo Geletti**

Outline

- **Introduction**
 - The seismic section**
 - Seismic interpretation software**
 - Seismic data web sites**
 - Seismic trace display**
- **Raw data e final seismic section: elements of multichannel seismic processing**
- **Resolution: vertical and lateral**
 - Deconvolution**
 - Migration**
- **Velocity analysis and Depth migration**
- **Vertical Seismic Profiling (VSP) in seismic interpretation**
- **Coherent Noise in the seismic data: multiple reflections**
- **Some case studies**
- **Conclusion**
- **Questions**
- **Bibliography**

- **INTRODUCTION**

The Seismic method is the powerful geophysical techniques for imaging the Earth's interior.

This artificial source method involve the generation of seismic waves whose propagation velocities and transmission paths through the subsurface are mapped to provide information on the distribution of geological boundaries at depth.


An alternative method of investigation subsurface geology is, of course, by drilling boreholes, but these are expensive and provide information only at discrete locations. Nevertheless, seismic surveying does not dispense with the need for drilling because it can give a geological meaning to the seismic reflectors.

Seismic Reflection Interpretation


- Fundamental in applied research to geosciences
- Provides information regarding:
 - geometries of stratigraphic sequences
 - geometries of structural and tectonic elements
 - velocity of seismic waves
 - lithological characteristics

The seismic interpretation attributes geological meaning to geophysical data and produces reconstructions of:

2D sections, structural maps, fault systems, slumping and geo-hazard etc.




Crustal seismic section: example of seismic interpretation



Interpretation software


Seismic section



Position map of 2D seismic survey by an interpretation software

Seismic data websites:

- <http://unmig.sviluppoeconomico.gov.it/videpi/>
- <http://see-atlas.leeds.ac.uk:8080/home.jsp>



well

ViDEPI PROJECT
Visibility of petroleum exploration data in Italy

Home The project Data Maps Login

ViDEPI Project
Visibility of petroleum exploration data in Italy

The ViDEPI project has been designed to make all the documents concerning Italian oil exploration easily accessible. The documentation concerns expired, and therefore public, mining permits and concessions, filed since 1957 with UNMIG, National Mining Office for hydrocarbon and geothermal energy of the Ministry for Economic Development.

Oil exploration in Italy is subject to the **Law n. 6 of 11 January 1957**, which, among other things, regulates the foundation of UNMIG, National Mining Office for hydrocarbon and geothermal energy, **Directorate-general for mineral and energy resources**, based at the Ministry for Economic Development with branch offices in Bologna, Rome and Naples. Current regulations establish that operating oil Companies shall provide UNMIG with progressive technical reports on the activities carried out on their permits and concessions; the reports shall include copies of exemplifying documents, such as geologic maps, structural maps, final well logs, seismic lines, etc.

The law establishes that the filed documents shall become available to the public a year after the permit has expired. This has led to the creation of what is today, after 50 years, an important data base on our Country's subsurface. Before the implementation of the ViDEPI project, the documentation was available only on paper and difficult to consult, arranged as it was on the basis of the mining concession in which it had been acquired and filed by the various UNMIG offices.

2009-2018 - ViDEPI Project - Visibility of petroleum exploration data in Italy
Ministry for Economic Development DGRM - Italian Geological Society - Assomineraria (cc) BY - This work is licensed under a Creative Commons Attribution 3.0 Italy License

VSA Virtual Seismic Atlas
Sharing the geological interpretation of seismic data

Log In

SEARCH ABOUT SPONSORSHIP WHAT'S NEW HELP

Welcome to the Virtual Seismic Atlas.

The VSA has been created to share the geological interpretation of seismic data. By browsing freely through the site you will find seismic images and interpretations. And you can download higher resolution images for your own use - all without signing in. There are no membership fees, all we ask is that you request the intellectual property rights of the contributors who have posted images on the VSA. Just hit the link "EXPLORE THE VSA" to start!

Use the VSA to find images of subsurface structures, submarine landscapes, the structure of oceanic water bodies and much more. You can search for analogues of subsurface geology, rapidly contrasting geological features, finding areas of controversy. You'll find results from cutting-edge technology and breaking research alongside historically important images of the subsurface.

The VSA is used by thousands of geoscientists each week. It's a great place to promote your own science, and for companies to showcase datasets and expertise. It's simple to author new content and link to your research entrapages or related publications.

Use and cite new content. If you wish to post interpretations or new seismic images we can set you up with authoring permission. Find out more from the "About the VSA" link below.

The "Press Release" button for "New" content explains how the University of Leeds, the host for the VSA, collects and uses your personal data when you use the website.


EXPLORE THE VSA

Learn more about the VSA

- Explore
- About the VSA
- Support for the VSA
- Help

The VSA is an industry-academia partnership

UNIVERSITY OF ABERDEEN UNIVERSITY OF LEEDS NERC





ViDEPI PROJECT

Visibility of petroleum exploration data in Italy

Home The project Data Maps Login

Geographic consultation

Download KML files

In questa pagina sono presenti i collegamenti ad alcuni file in formato kml utilizzabili con il programma Google Earth.

KML files

1. Reconnaissance seismic campaigns of the offshore areas

- Zona B
- Zona C
- Zona D
- Zona E
- **Zona F**
- Zona G

2. CROP Atlas Project, seismic reflection profiles of the Italian crust

3. Seismic lines acquired in expired mining permits and concessions
4. Expired mining permits and concessions
5. Wells


To carry out a geographic search, you need to download Google Earth program on your pc. Click on earth.google.com/intl/en to install Google Earth.

Go to support.google.com/earth/2hl-en to learn how to use the program.


To view all the expired mining permit and concessions and wells, move the cursor to the right on the time bar.

The form and location of the elements is purely indicative, they are supplied only to inform users of the presence of data in a area.


2009-2018 - ViDEPI Project - Visibility of petroleum exploration data in Italy
 Ministry for Economic Development (MISE) - Italian Geological Society - Associazione
 (cc) BY - This work is licensed under a Creative Commons Attribution 3.0 Italy License



ViDEPI dataset




Position map of the seismic profiles of the Italian geophysical exploration projects MS and CROP acquired in 1969 – 1982 and 1991-1995 respectively




10 km

OGS - Istituto Nazionale di Oceanografia e di Geofisica Sperimentale	
SEISMOLOGIA CRUSTALE	
SEZIONE CRUSTALE, AREA DI TIRRENO ALTO	
L.S. 11.1	
1969	
PROGETTO	ESECUZIONE
OGS - Istituto Nazionale di Oceanografia e di Geofisica Sperimentale	OGS - Istituto Nazionale di Oceanografia e di Geofisica Sperimentale




1 s (TWT)


Examples of vintage crustal seismic sections MS 1 acquired in Tyrrhenian Sea in 1969




Example of a part of seismic line displayed in wiggle/variable area. On the right is shown a single trace that constitutes the section above (seismogram)


Seismogram: this is an example of a single trace of a seismic profile as shown in the figure above









Seismic section displayed in "variable area"



Seismic section displayed in “variable density” in grey scale




Seismic section displayed in "wiggles + variable area"




Seismic section displayed in grey scale variable density with the “enhanced reflections” in red colour


MultiChannel
Seismic reflection
data (MCS)

Raw data
(shot gathers)




Processed data
(stack section)





Shot gather


Water Bottom reflections



Example of «processing flow chart» that define two different output: Pre-Stack Time Migration (PSTM) and Pre-Stack Depth Migration (PSDM). This sequence has been applied to «crustal data» with «long offset streamer».

The main steps are the following:


- Reformatting
- Editing
- Sorting
- Gaining
- Deconvolution
- Velocity analysis
- NMO correction and stacking
- Migration



Seismic resolution


Vertical resolution Lateral resolution

Examples of seismic wave resolution





Overlap of a seismic wave and: A) a sonic log, B) a photo of a rock face of about 40 m

Very high resolution acoustic profile




Low resolution seismic profile
(crustal data)




Comparison between a seismic wave and: b) Big Ben of London, c) a well data





Seismic resolution

Very high resolution CHIRP profile (sweep 2-7 kHz)





Factors affecting horizontal and vertical seismic resolution



Swath bathymetry: Side Scan Sonar (above) & MultiBeam (left) data

CHIRP profile (Very High Resolution)

High penetration seismic line (low resolution)


Seismic resolution

Vertical Resolution


$$\lambda/4 = v/4f$$

v (m/s)	f (Hz)	$\lambda/4$ (m)
2000	50	10
3000	40	18
4000	30	33
5000	20	62


Threshold for vertical resolution



The relationship between velocity (v), dominant frequency (f) and wavelength ($\lambda = v/f$).



Definition of the Fresnel zone AA'



Lateral Resolution

$$r = (v/2)\sqrt{t_0/f}$$

t_0 (s)	v (m/s)	f (Hz)	r (m)
1	2000	50	141
2	3000	40	335
3	4000	30	632
4	5000	20	1118


Threshold for lateral resolution ($t_0 = 2z/v$, $r=OA$)




Seismic resolution

Example of high penetration (low resolution) multichannel seismic reflection profile (CROP project) in Central Adriatic Sea.

Vertical resolution = 75 m
Lateral resolution = 4840 m
($v=6$ km/s; $f=20$ Hz)


→ 40 km depth





Seismic resolution

Examples of two seismic sections acquired by different sources: (left) by 2 GI guns (11,6 l); (right) by an array of 16 air guns (70 l)




Factors affecting horizontal and vertical seismic resolution: processing solution


Vertical resolution

Deconvolution

Example of the effect of the deconvolution on the definition of seismic data (synthetic data). The deconvolution "shrinks" the wavelet as shown in the figure (synthetic data)




Before the deconvolution




After the deconvolution

Vertical resolution




Seismic stack section a) before and b) after deconvolution (Yilmaz, 2001)

Lateral resolution




Migration

- Consider a source-detector (s-d) on the surface of a medium of constant seismic velocity. For a given reflection time, the reflection point may be anywhere on the arc of a circle centred on the s-d position. On a non migrated seismic section the point is mapped to be immediately below the s-d.





- A planar-dipping record surface derived from a non-migrated seismic section (blu line) and its associated reflector surface (red line).




Stack section (a) before and (b) after migration. (from Yilmaz 2001)

Lateral resolution




- a) A sharp synclinal feature in a reflecting interface, and
- b) the resultant «bow-tie» shape of the reflection event on the non-migrated seismic section.

Lateral resolution: migration




a)



b)

Example of seismic reflection profile across two buried channels (a) non-migrated section with the presence of «bow-tie» effect (red ellipses) and b) after migration

2D migration




2D migration is an imperfect process on the «strike line»


2D and 3D migration

The image shows an example of a 3D geological model with two anticlinals (green and yellow object) and a direct fault (red and blue objects).

The seismic data along line 6 shows the comparative effects of 2D and 3D migration (from French, 1974). Only 3-D migration is able to provide a seismic profile faithful to the real situation of pending layers. However, the 2-D migration provides an often satisfactory result for interpretation.




From French, 1974




2D migration

Example of 2D migrated seismic profile with the presence of salt domes, some of which are lateral to the vertical plane of the section.


3D migration



Example of "3D cube" from Brown (1986)



Definition of the Fresnel zone AA'



Lateral Resolution

$$r = (v/2) \sqrt{t_0/f}$$

t_0 (s)	v (m/s)	f (Hz)	r (m)
1	2000	50	141
2	3000	40	335
3	4000	30	632
4	5000	20	1118

Threshold for lateral resolution ($t_0 = 2z/v$, $r=OA$)




FAULTS

Example of faults in the geological layers.

The faults represent points of inhomogeneity that originate diffractions. You can see the geological layers that are dislocated by the faults


Migration: diffraction collapse




(a)

(b)

Seismic section (a) before and (b) after migration: the diffractions have been collapsed and the faults are evident. The presence of diffractions can sometimes create difficulties of interpretation: the point of breakage of the reflector is not easily identifiable in the non-migrated section. In the case b) the section was migrated and the base of the salt layer is more evident than in stack section a)



Blow up of the previous seismic image




Velocity field of the seismic waves is the parameter that allow to get the geologic model from the seismic section


Velocity analysis

A: a common mid-point trace gather (left) and semblance coherence contour (right).

Peaks in coherence give the stacking velocity. (yellow line)




B: The same common mid-point trace gather of A with the normal move-out correction (NMO) after the velocity picking (yellow function). The interval velocity (red function) associated to this stacking velocity it will use in migration process.




The field velocity section can be obtained from the seismic data using the stack velocities. We are going to flatten the reflections in CDP gather as in the image, picking the maximum coherence in the semblance. In this way velocity functions are obtained which we will interpolate with the others in order to have a section.

Velocity field and depth migrated seismic section




Example of Pre Stack Depth Migrated section with its velocity field superimposed (see next slide for the interpretation)

Depth migrated seismic section




Example of Pre Stack Time/Depth Migration section and the interpretation superimposed with velocity function.

(from Micallef et al., 2018)




Time Migrated section with salt intrusion




Depth migrated section

Pull up velocity effect





(Above) Result of data processing in time domain: the pull-up event (red circles) of about 200 ms occurs beneath a salt dome and affects the underlying layers. (below) Result of Pre-Stack Depth Migration (PSDM) showing flattened pullup event. (modified from Dal Cin et al. 2016)



Vertical Seismic Profiling (VSP) in seismic interpretation


Example of correlation of well data (VSP) with the seismic section for geological interpretation

In some case we have a borehole across the seismic section and we can correlate these two type of data. From this we can therefore give a lithological meaning to the individual reflectors of the seismic profile passing through the well. In the image below and on the right, we can see an example of correlation between the seismic profile and the data obtained from the registration of the VSP in a borehole. The VSP was made after the geologic sampling in the same borehole.



- MULTIPLE REFLECTIONS -


The problem of coherent noise in marine seismic profiles



Long-path multiples appear as distinct events. Short-path multiples are added to primary reflections and tend to come from shallow subsurface phenomena

The marine seismic dataset is often characterized by the presence of multiple reflections that affect the quality of the section itself. The multiple are the coherent noise that it leads to the misinterpretation of real reflective horizons. There are a lot of type of coherence noise, but the first water bottom multiple reflections are the most frequently.

MULTIPLE REFLECTIONS






Example of seismic stack section with multiple reflections (M1 and M2)

Gas seeping features


Mud volcanoes and pockmarks

MB data with pockmarks




Schematic model of gas seeping related upward fluid migration

Seismic line & MB image with mud volcano




Sub Bottom Profilers (CHIRP) with pockmarks

Some example of gas seeping features. Sea bed fluid flow, also known as submarine seepages, involves the flow of gas and liquids through the seabed. This geological phenomenon has widespread implications in seabed slope instability, drilling hazard, hazards to seabed installations and so on. Seabed fluid flow affects seabed morphology (pockmarks, mud volcanoes). Natural fluid emissions also have a significant impact on the composition of the oceans and atmosphere: methane emissions have important implications for the global climate change.



Echosounder CHIRP profile where there are gas evidences






Crustal seismic profile CROP- M15

a) Image of the seismic reflection profile with the evidence of bright spots indicating the presence of gas in the Plio-Quaternary sediments, b) multibeam bathymetry (MB) and a pseudo 3D image (c) with CHIRP profile and MB where it is highlighting a system of active faults along which some pockmarks can be identified. The seismic line shows the presence of a deep diapiric structure that also deforms the sea floor





Detail of the CHIRP line with evidence of gas plume





In this slide there are some images of gas/fluid presences within the sediments. On the right, above you see an image of pockmarks on swath bathymetry of multibeam. Below, two chirp profiles. On the left side you see a seismic section where at a deep of 400/450 m there are seismic anomalies below superficial gas evidence in chirp profiles

Image of Pockmarks in MB data



Pseudo 3D image of a pockmarks visible on two orthogonal CHIRP profiles

Conclusion - 1

The Multi-Channel Seismic Reflection (MCS) method is:

- The most widespread method for the geophysical prospecting of the subsoil, fundamental in the exploration of hydrocarbon reservoir.
- It provides more detailed information than any other non-invasive method on stratigraphy, structure and properties of materials.
- It uses arrival times, amplitude and phase of the echoes from the discontinuity in the elastic properties present in the subsoil to obtain its position and physical properties (acoustic impedance, velocity propagation of seismic waves, elastic parameters, ...).

Disadvantages of the MCS method:

- High costs of data acquisition (R/V OGS Explora ship time - > 15-20 K€ a day)
- Complex signal processing required
- Numerous specialized people needed
- For a survey, numerous permits and authorizations are required

Conclusion - 2

The seismic reflection interpretation attributes geological meaning to geophysical data.

Interpretation provides information on:

- geometry of stratigraphic sequences and structural/tectonic elements
- seismic wave velocity
- Lithological characteristics

Applications for reconstructions of 2D section, structural maps, fault systems, slumping and seismic hazard.

The interpretation is made by a team of geologists / geophysicists / physicists with different skills who work in synergy.

“Interpretation is a combination of both art and science “ (Lines and Newrick, 2004)

Questions

1. What is a seismic section?
2. What is the difference between seismic and geological section?
3. What is the vertical scale in a seismic section?
4. Which is the difference between seismic stack section and migrated section?
5. What is a diffraction?
6. What is a «bow-tie» event?
7. What is a multiple reflection?
8. What are the advantages of a migrated section?
9. Which seismic parameter is fundamental in depth migration?
10. Which is the first reflection in a marine seismic section?
11. What is the *acoustic basement*?
12. What is a « bright spot» in a seismic section?
13. What are the seismic characters that identify the possible presence of gas in the sediments?
14. What are the gas seeping structures?
15. What is the best acoustic method to study these structures?

Bibliography:

- AN INTRODUCTION TO GEOPHYSICAL EXPLORATION, di Philip KEAREY, Michael BROOKS, Ian HILL; (2002) – Blackwell Science – ISBN 0-632-04929-4.
- MARINE GEOPHYSICS, E. J. W. JONES; (1999) – Wiley – ISBN 0-471-98694-1
- SEISMIC DATA ANALYSIS (Vol. I & II), Öz YILMAZ; (2001) – SEG – Vol. I ISBN 1-56080-098-4; Vol. II ISBN 1-56080-098-2
- Lines and Newrick - FUNDAMENTALS OF GEOPHYSICAL INTERPRETATION
- Sheriff and Geldart - EXPLORATION SEISMOLOGY
- Anstey - SEISMIC INTERPRETATION - The Physical Aspects
- Herron – FIRST STEP IN SEISMIC INTERPRETATION
- Hovland & Jadd - Seabed Pockmarks and Seepages
- Jadd & Hovland – Seabed Fluid Flow

Websites:

- ViDEPI: <http://unmig.sviluppoeconomico.gov.it/videpi/>
- Virtual Seismic Atlas: <http://see-atlas.leeds.ac.uk:8080/home.jsp>

Methods for the analysis of sedimentary basins

Part 2

Geophysical methods: indirect analysis of deposits in the subsoil
(+ or less ancient, on land and at sea) = today's lesson by Valentina volpi


Geological methods at sea: coring (superficial deposits) and drilling (deeper /
ancient / lithified deposits) = Course of Marine Geology

Geological methods on land: analysis of rocky outcrops, geological cartography,
measurement of stratigraphic sections... (exhumed rock deposits).

Geological / naturalistic methods: analysis of recent deposits: historical
cartography, trenches, ...

Coring of marine sediment


Sediment coring system



A sediment coring system is composed by a weight (corer head) that is used to force the barrel into the sea bottom. Inside the barrel there is a plastic liner that will hold the sampled sediments. At the base of the barrel the core catcher prevents the sampled sediments to flow out the barrel after sampling, and the cutter facilitate the detach of the sampling system from the ground.



This system can work with or without a triggering system




Courtesy: R.G. Lucchi

Drilling



IODP
INTERNATIONAL OCEAN
DISCOVERY PROGRAM

Joides Resolution



Outcrops

La Baronía




Large scale
(outcrop)

10 m



Mid scale
(bed)

50 cm



2,5 cm Small scale
(Sedimentary structure)

Martinius, 2011

DOI: 10.1007/978-94-007-0123-6_18

Three-scale approach

A three-scale approach should be attempted, wherever possible, for distinguishing the various facies types (especially the deep-water ones).

It includes:

large-scale (oceanographic and tectonic setting),
regional-scale (architecture and association)
and small-scale (sediment facies) observations.

Stow and Smillie, 2020 *Geosciences*, 10(2), 68;

<https://doi.org/10.3390/geosciences10020068>



Courtesy
of D.A.V.
Stow




Rebesco & Camerlenghi (2008)

Recognizing contourite deposits in ancient sedimentary series presently exposed on land, is a difficult task. The distinction between contourites and reworked turbidites is controversial.

Diagnostic criteria are their **facies and ichnofacies, texture and sequences, microfacies and composition**. **Sedimentary structures** are also “diagnostic indicators”, but for their interpretation its full context should always be considered. **Medium-scale criteria** (hiatuses and condensed deposits, variation in the thickness, geometry, palaeowater depth, geological context) can be definitive. **Large-scale criteria** (palaeoceanographic features and continental margin reconstructions) are essential, but generally more problematic to apply on outcrops.

Seismics characteristics



triple-scale approach that involves 3 “orders of seismic elements”.

Large scale (overall architecture): I-order elements (major changes in current strength and sediment supply): External geometry, Bounding reflectors, Gross internal character.


Medium scale (internal architecture): II-order seismic elements (reflecting smaller fluctuations): lens-shaped, upward-convex geometry; uniform stacking pattern; down-current migration or aggradation; downlapping reflector terminations

Small scale (internal acoustic character): III-order seismic elements: facies analysis (continuous, (sub)parallel, wavy, structureless), and attribute analysis (bedforms).

Sedimentary structures in cores

From Rebesco M.,
Camerlenghi A.,
(Eds.) 2008.
Contourites,
Developments in
Sedimentology, 60,
Elsevier, 666 pp.

Originally in:
Shanmugan, 2008;
Shanmugam et al.,
1993;
Shanmugam, 2012



Sedimentary structures in outcrops



<https://uhlibraries.pressbooks.pub/historicalgeologylab/chapter/chapter-4-sedimentary-structures/>




Flute casts from the central Alps, Switzerland. The view is from the underside of the rock. Image credit: Chris Spencer


Symmetrical ripples in Devonian-Mississippian age sandstone from Ohio, USA. These are all views from the top. Image credits: James St. John,

Jacob's staff






Sedimentary logs



How do you estimate grain size?



Geological mapping



Geologic Map of New Mexico


New Mexico Bureau of Geology and Mineral Resources
Celebrating 75 Years of Service
A DIVISION OF NEW MEXICO INSTITUTE OF MINING AND TECHNOLOGY
Peter A. Scholle, State Geologist
2013

Lithology

The lithology of a rock unit is a description of its physical characteristics visible in outcrop, or with low magnification microscopy. Physical characteristics include color, texture, grain size and composition.

The lithology is the basis of the subdivision of the rock sequences into single lithostratigraphic units for the purposes of mapping and correlating the areas.

Chronostratigraphic sequences



Facies

The facies is the sum of the overall characteristics of a rock unit, which reflect its origin and differentiate the unit from others around it.

Within the facies, we can distinguish eg lithofacies (stratigraphic unit distinct from the adjacent ones based on lithology),


Biofacies,

Microfacies

Ichnofacies


Seismic facies

...



Modern analog

Gironde estuary tidal bars: A modern analogue for deeply buried estuarine sandstone reservoirs



LEGEND:

- Fluvial-estuarine transition point bars
- Estuarine heterolithic point bars
- Tidal mudflats and marshes
- Estuary funnel tidal bars
- Estuarine mud
- Tidal inlet channel fill

- Tidal bars supratidal zone
- Tidal bars intertidal zone
- Tidal bars subtidal zone
- Bathymetric lines

- Cores location realized in this study in the Plassac tidal bar
- Cores location realized in 2010 in the Plassac tidal bar
- Core transect location in the Richard tidal bar

XX km: Distance from the estuary mouth

← - - → TMZ position during High-River Stage (HRS)


← - - → TMZ position during Low-River Stage (LRS)



Virolle et al., 2020.
Marine Petroleum
Geology

Historical data

River-mouth geomorphological changes over > 130 years, Pratellesi et al., 2018. Marine Geology



Difference surface between the 1882 and 1954 bathymetry; erosion areas are represented in blue and accretion ones in red (values are expressed in meters). Black and purple lines in the inset delineate, respectively, the 1882 and 1954 msl contour lines.

Trenches



stratigraphic soil layers, Vjosa Valley, Albania