

An underwater photograph showing a large school of small, silvery fish swimming in clear blue water above a dark, rocky reef. Sunlight rays penetrate the water from the top, creating a bright, shimmering effect. The fish are densely packed in some areas and more sparse in others, moving in various directions. The reef below is covered in green algae and other marine life.


**GLOBAL CHANGE ECOLOGY AND SUSTAINABILITY**  
**a.a. 2023-2024**

**Conservation and Management of Marine Ecosystems**  
**Prof. Stanislao Bevilacqua ([sbevilacqua@units.it](mailto:sbevilacqua@units.it))**

**Marine Biodiversity**

# Marine biodiversity

4/51



~34 animal phyla, 80% is marine or mostly marine

~almost all of them are benthic or have benthic taxa

(...and don't forget most of algae)

~1,500,000 known species on Earth

~300,000 are marine, ~85% of them are benthic




# How many species

About 240,000 marine species described

58,000-72,000 marine species sampled but still not described



There could be 0.7-1.0 million marine species



Appeltans et al. 2012

# Patterns

## Latitudinal gradients



Marine biodiversity peaks at tropical latitude (Snelgrove et al. 2016) and at shallower depths (Costello & Chaudhary 2017)



# Factors affecting biodiversity

- Geographic factors (latitude, depth)
- Productivity, climatic factors, history
- Predation, competition
- Disturbance, isolation, heterogeneity



The intermediate disturbance hypothesis (Connell 1978). Small-infrequent or large-frequent disturbance could reduce diversity, which is maximum at intermediate levels of disturbance


Stability-Time Hypothesis (Sanders 1968). This model says that physical instability in an environment prevents the establishment of diverse communities. However, if physically stable conditions persist for a long period of time, speciation and immigration will cause species diversity to increase gradually.




# Distribution, factors and processes

Sea temperature and solar radiation influence the distribution of benthic organisms, especially algae and corals and the associated fauna. Shifts in distribution (climate change), mass mortalities, bleaching


## Global distribution of macroalgal genera




## Solar radiation



## Global distribution of coral species





## Surface temperature






# Distribution, factors and processes





**Currents** play a crucial role in shaping the distribution of benthic species, which in most cases are sessile or sedentary and rely on water movements for their dispersal. Climatic factors, substrate availability, geographic barriers, pre- and post-settlements events drive community assembly generating differences in species distribution and community composition (beta-diversity).



Patterns of variation in  $\beta$ -diversity provide information on the structure (nodes and pathways) of units of ecological connectivity, and for conservation and resilience of marine ecosystems



# Productivity





**Productivity and high energy flow could sustain higher number of species with respect to less productive areas**

(maps from Costello & Chaudhary 2017)



# Temperature




(maps from Costello & Chaudhary 2017)

Rates of genetic divergence and speciation are both governed by metabolic rate and therefore show the same exponential temperature dependence. So, higher temperature increases speciation rates (Allen et al. 2006)




# Stability

A interoceanic gradient exists in benthic fauna: the Pacific Ocean (older) has much more species than the Atlantic Ocean.



**b**

Speciation rate




However, recent studies highlighted that speciation rates can be higher at higher latitudes

Rabosky et al. 2018



# Biodiversity in the last eon

## Biodiversity during the Phanerozoic



5 big mass extinctions. Biodiversity is increasing