
### **Climate change and global change**

**Climate change** refers to changes of climatic factors at a global scale (e.g., increasing SST, but also sea level rise, ice melting, and atmospheric phenomena)


**Global warming** is the increasing warming temperature at global scale in the last century, mostly due to fossil fuel use, referring to the baseline of 1950-1980 (Goddard Institute for Space Studies –NASA)

**Global change** refers to all changes that are occurring as a consequence human activities, including climate modifications, biodiversity loss, alterations of the natural environments and so on...


## Paleoclimate



We are in a Glacial Era started 40 millions years ago. Within a glacial era, glacial and interglacial periods alternate. The last glacial period started more than 100k years ago and finished about 10k years ago. Now we are in a interglacial period. However, we are less than 1 degree cooler than warmer peak in past interglacial periods. (Hansen and Sato, 2011)




### **Greenhouse effect**



CO<sub>2</sub> N<sub>2</sub>O CH<sub>4</sub> H<sub>2</sub>O CFC

### **Carbon dioxide emissions**



## Methane in the subpolar regions




nature communications

Article OPEN Published: 15 August 2018

V

### 21st-century modeled permafrost carbon emissions accelerated by abrupt thaw beneath lakes


Katey Walter Anthony <sup>™</sup>, Thomas Schneider von Deimling, Ingmar Nitze, Steve Frolking, Abraham Emond, Ronald Daanen, Peter Anthony, Prajna Lindgren, Benjamin Jones & Guido Grosse



## **Specific heat**

Specific heat: the amount of heat necessary to raise the temperature per unit mass by 1 degree Kelvin Air 0.25 Kcal/kg ° C Rock (average) 0.20 Kcal/kg ° C Seawater 0.95 Kcal/kg ° C

## **Earth Energy Imbalance**



More than 90% of energy imbalance of the planet is stored in the ocean, increasing ocean heat content (OHC), while the residual heat is manifest in melting of both land and sea ice, and in warming of the atmosphere and land surface. OHC is increasing due to greenhouse gases.

### Storming



### Magnitude of the Observed Extreme Precipitation during Hurricane Harvey Mark D. Risser 🕿, Michael F. Wehner

**Research** Letter

# Increase in strength and frequency of hurricanes, coastal flooding


**Geophysical Research Letters** 

Attributable Human-Induced Changes in the Likelihood and






## Increasing energy in weather phenomena



### Sea level rise




**Representative** Concentration Pathway (RCP) has been defined by **IPCC, as carbon dioxide atmospheric** concentration, to depict climate scerario by IPCC. Numbers (2.6, 4.5, 6.0, 8.5 are radiative forcing levels, delta between radiation adsorbed and dispersed back to space, in  $W/m^2$ ). **RCP2.6** carbon dioxide emission <sup>2100</sup> peak is now and then decline **RCP4.5 peak in 2040 RC6.0 peak in 2080 RCP8.5 continue to increase until** the end of century

### **El Nino**


**Atmospheric-oceanic coupled process El Nino - Southern Oscillation (ENSO)** 

Strong Trade winds EC Pacific high pressure, cold waters (upwelling) WC Pacific low pressure, precipitations, warm waters

Cycles of few years, during winter (December-January) Reduction in strength of Trade winds and inversion (El Nino)




### **ENSO (El Nino Southern Oscillation)**




**Normal conditions:** wind trades blow strong, the Humboldt current is strong, upwelling occurs on the S America coasts (Chile and Ecuador), high pressure is on S-central Pacific and low pressure (wet, warm) on the Australian and Indonesian coasts. Superficial waters in the east Pacific are cold. When T is 0.5 ° C or more below the seasonal average, we have **Ia Nina**.

### **ENSO (El Nino Southern Oscillation)**



**El Nino:** cyclic but irregular, every 2-7 years (5 on average) with max during winter (december). It is an increase in superficial water temperature in the central-SE Pacific of at least 0.5 ° C above the average T for at least 5 months. Wind trades are weak, the Humboldt current is weak, upwelling on the S America coasts (Chile and Ecuador) is strongly reduced or absent, high pressure is on the Australian and Indonesian coasts and low pressure (wet, warm) on the S-central Pacific coasts. Superficial waters in the east Pacific are warm.

## Thermal anomalies and melting ice



# NOAA

square km

noillin

Increasing SST, causing thermal anomalies, melting polar ice, and altering oceanic currents AVERAGE SEPTEMBER EXTENT


Data source: Satellite observations. Credit: <u>NSIDC</u>/NASA




#### RATE OF CHANGE

↓ 12.8


percent per decade



### **Mass mortalities**




### Acidification




Oceans absorb about one third of atmospheric CO<sup>2</sup>. So, increasing level of carbon dioxide in the air results in increasing levels in sea water. This lead to increase in carbonic acid, and H<sup>+</sup> ions that decrease ocean pH, which is generally slightly basic.

### **Effects on carbonate deposition**

Aragonite and calcite are the two crystalline forms of calcium carbonate, used by most of marine organisms with calcified structures (corals, molluscs, crustaceans, coralline algae, etc.).  $\Omega_{arag}$  was higher in the surface mixed layer. Higher hydrostatic pressure, lower water temperature, and more CO<sub>2</sub> buildup from biological activity in the absence of air-sea gas exchange helped maintain lower  $\Omega_{arag}$  in the deep ocean.




### **Effects**



### **Food webs**



Climate change could drive marine food web collapse through altered trophic flows and cyanobacterial proliferation



Habitat destruction for seals and bears with consequent loss of feeding grounds and refuge. Plankton can be affected with bottom-up cascading effects. Shift in composition of plankton producers could reduce energy transfer through trophic webs, leading to the decline of apical species populations.

### Shifting Patterns of Life in the Pacific Arctic and Sub-Arctic Seas


#### **Annual Review of Marine Science**

Vol. 4:63-78 (Volume publication date January 2012) First published online as a Review in Advance on September 19, 2011 https://doi.org/10.1146/annurev-marine-120710-100926


#### Jacqueline M. Grebmeier



### Invasions



### Impact on fisheries



### Harmful species



10 µm

Lagocephalus sceleratus

### The Mediterranean Sea


#### Its history and present challenges

Metamorphoses: Bioinvasions 2014 in the Mediterranean Sea

Increasing risk for human health due to the introduction of toxic or harmful species

B.S. Galil and Menachem Goren

### **Extinction scenarios**



Extinction rise with increases in annual mean global surface air temperature plotted under historical greenhouse gas emissions (petagrams of carbon per year) and divergent future scenarios, yielding radiative forcings of 2.6 W/m<sup>2</sup> [i.e., RCP/SSP1-2.6 versus 8.5 W/m<sup>2</sup> in 2100] Penn and Deutsch 2022