

An underwater photograph showing a large school of small, silvery fish swimming in clear blue water above a dark, rocky reef. Sunlight rays are visible filtering down from the surface.


**GLOBAL CHANGE ECOLOGY AND SUSTAINABILITY**  
**a.a. 2024-2025**

**Conservation and Management of Marine Ecosystems**  
**Prof. Stanislao Bevilacqua ([sbevilacqua@units.it](mailto:sbevilacqua@units.it))**

**Nets and cycles**

# Trophic chains and the importance of detritus

Detritus (90% of primary production)



# Detritus

*“non-predatory loss of organic carbon from each trophic level or inputs from external sources”*

(Wetzel et al., 1972)


So, everything non-living and organic, irrespective of its size, composition and origin




# Origin



Terrestrial supply  
Small plankton remains  
Moult  
Fecal pellets




Large animal remains  
(fish and mammal carcasses)



Algal and plant debris  
(kelp, seagrass, etc.)

# Classification of organic matter

- 
1. **Type:** Living (Organisms) 10%  
Detritus (fecal pellets, excretions, etc.) 90%
2. **Size:** DOM (<0.45  $\mu\text{m}$ )  
POM (>0.45  $\mu\text{m}$ )
3. **Sources:** *in situ* production 10-100%  
exogenous 0-90%
4. **Trophism:** Bioavailable 1-10%  
Recalcitrant 90-99%

# Type

Most of living organic matter in oceans comes from planktonic and benthonic bacteria, protists, phytoplankton, microzooplankton and meiofauna

Larger components are negligible in terms of amount and numbers

Microzooplankton are a group of heterotrophic and mixotrophic planktonic organisms between 20 and 200  $\mu\text{m}$  in size. Important contributors to the group are phagotrophic protists such as flagellates, dinoflagellates, ciliates, radiolarians, foraminiferans, etc., and metazoans such as copepod nauplii, rotiferans and meroplanktonic larvae, among others.



# Trophism

POM is composed by proteins, carbohydrates and fat acids

DOM is composed by a huge range of substances of molecular weight from very few until  $>100.000$  d, and includes, for instance,

*a. virus*

*b. carbohydrates (glucose, 50-60%)*

*e. aromatic compound (e.g., phenol, lignin, lipids)*

*f. amino acids*

*g. DNA and RNA*

DOM pool is largely produced by phytoplankton and decomposition of bacterial and virus action

DOC/POC ratio 10-20:1 in the water column


DOC  $<$  5% of TOC in sediments

Labile organic matter is easily and rapidly available to be remineralized by organisms, whereas recalcitrant organic matter is formed during decomposition and other processes (agglomeration), and is difficult to be degraded by bacteria unless during long periods.

Example: CRAM (carboxyl-rich alicyclic molecules) amino-sugars, amino acids, terpenoids, lignin)


# Carbon

1,85 billions Gt on Earth. Only 44.000 Gt are on the surface of the planet, while the remaining is in the nucleus and the mantle. 94% of carbon on the surface is stored into the ocean (water and sediments, mostly as bicarbonate and carbonate ions), 4.5% in the biosphere and 1.5% in the atmosphere.





# Carbon cycle and balance



Short term cycle (1-10<sup>3</sup>y).

The second part of the whole cycle consists of a long term cycle (10<sup>4</sup>-10<sup>6</sup>y), through weathering, volcanic activities, subduction of sediments and rocks, huge stocks in soil.


# BCP, ML, and MCP

**Table 1.** Definitions and major impacts of the BCP, ML and MCP.

| Concept               | Definition   | Major impacts and focus   |
|-----------------------|--|---|
| Biological pump       | A complex ecosystem process that transports particulate organic carbon from the epipelagic zone to the deep interior of the ocean and further to the ocean floor   | Sequestration of atmospheric CO <sub>2</sub> through vertical transportation of living biomass to marine sediments; focusing on sediment storage  |
| Microbial loop        | A 'feedback' pathway of loss of the primary production to the environment in the form of dissolved organic matter and the utilization of the latter by bacteria that feed the protozoa, which enter the food chain | The role of bacteria in sequestering nutrients from the environment, which are consumed by protozoa; focusing on organismal populations above thermocline   |
| Microbial carbon pump | A conceptual framework for understanding the role of microbial processes in the production of recalcitrant dissolved organic matter in the ocean water column  | Sequestration of atmospheric CO <sub>2</sub> through transformation of labile organic matter to recalcitrant organic matter; focusing on capacity of the ocean to store atmospheric CO <sub>2</sub> |




# Biological Carbon Pump



# Microbial loop


Farooq et al 1983



The microbial loop describes a trophic pathway in the marine microbial food web where dissolved organic carbon (DOC) is returned to higher trophic levels via its incorporation into bacterial biomass, and then coupled with the classic food chain formed by phytoplankton-zooplankton-nekton.

# Microbial Carbon Pump


Zhang et al 2018



The MCP mediates the transformation of labile carbon to RDOC, which constitutes the majority of DOC (662 Gt C) and persists in the ocean for up to 4000–6000 years

# Viral shunt

This process sustains a high prokaryotic biomass and provides an important contribution to prokaryotic metabolism, allowing the system to cope with the severe organic resource limitation of deep-sea ecosystems




The viral shunt, releasing on a global scale , **37-50 megatons of carbon per year**, is an essential source of labile organic detritus in the deep-sea ecosystems

# Terrestrial export of nutrients




# N cycle






# S cycle




# Trophic webs

1 law of thermodynamics

2 law of thermodynamics




# Energy flow



Efficiency of energy transfer is low, 10% on average at each trophic level. This is reflected in the biomass ratio between levels. Most of energy is lost in movement, excretions, fecal dejections, heat, so that moving from the basis (primary producers) to higher levels, the total sustainable biomass is drastically reduced. In some cases, in marine environments, the pyramid can be inverted because of differences in temporal turnover of organisms across levels


# Differences between land and sea

Because of these energy losses, most terrestrial ecosystems have no more than five trophic levels, and marine ecosystems generally have no more than seven. This is likely due to differences in the fundamental characteristics of land and marine primary organisms. In marine ecosystems, microscopic phytoplankton carry out most of the photosynthesis that occurs, while plants do most of this work on land. Phytoplankton are small organisms with extremely simple structures, so most of their primary production is consumed and used for energy by grazing organisms that feed on them. In contrast, a large fraction of the biomass that land plants produce cannot be used by herbivores for food, so proportionately less of the energy fixed through primary production travels up the food chain.




Growth rates may also be a factor. Phytoplankton are extremely small but grow very rapidly, so they support large populations of herbivores even though there may be fewer algae than herbivores at any given moment. In contrast, land plants may take years to reach maturity, so an average carbon atom spends a longer residence time at the primary producer level on land than it does in a marine ecosystem. In addition, locomotion costs are generally higher for terrestrial organisms compared to those in aquatic environments.


# Total primary production in the ocean




**Net Primary Productivity** (grams Carbon per m<sup>2</sup> per year)



# Trophic webs: coastal Mediterranean



# Trophic webs: pelagic



# Keystone species

A keystone species is an organism that helps define an entire ecosystem. Without its keystone species, the ecosystem would be dramatically different or cease to exist.

Keystone species have low functional redundancy. This means that if the species were to disappear from the ecosystem, no other species would be able to fill its ecological niche.

They could be predators or herbivores (or producers).

Keystone can have either small population size or large number of individuals. Generally, in the case of predators, small numbers can have strong effects on ecosystems.



# Paine's work

The term keystone species was first coined by Robert Paine (1966) after extensive studies examining the interaction strengths of food webs in rocky intertidal ecosystems in the Northwest Pacific. In his work, he studied a community dominated by the same species of mussels, barnacles, and the starfish, *Pisaster ochraceus*, which preys upon the other species as a top predator.



Pisaster



Thais

He had observed that the diversity of organisms in rocky intertidal ecosystems declined as the number of predators in those ecosystems decreased. He hypothesized that some of these consumers might be playing a greater role than others in controlling the numbers of species coexisting in these communities.



Gooseneck  
Barnacles



Limpets



Bivalves



Acorn  
Barnacles




Chitons

# Paine's experiment


He tested his hypothesis in an experiment in which an area of the intertidal was kept free of the starfish, comparing them with an undisturbed control area of equal size.

He observed dramatic changes after *Pisaster* was artificially removed compared with the control area that remained unchanged in its species number and distribution. After removal of starfishes the other species began to compete.



Within three months the barnacle, *Balanus glandula*, became dominant and after 9 months, it was replaced by another barnacle *Mitella* and the mussel *Mytilus*. The succession of species wiped out populations of benthic algae, causing some species, such as the limpet, to emigrate because of lack of food and/or space. After a year of the starfish's removal, species diversity significantly decreased in the study area from fifteen to eight species


# Small predators




Decline in predator snails *C. concholepas* allows mussels *Perumytilus* to overcompete barnacles, macroalgae and herbivorous snails *Fissurella*.




# Large predators



*Carcharhinus limbatus*



*Rhinoptera bonasus*




*Agropecten irradians*



Sharks play an important role, removing weak and/or sick fish. In this example, the decline of sharks in the eastern Atlantic (USA) lead to increasing population of rays (release of mesopredators), and to a decrease in scallops.

# Herbivores




Key role in maintaining coral reefs healthy removing algal turf, dead corals and preventing the dominance of algae after disturbance

# Keystone for trophic importance

Krill (Euphausiacea) are shrimp-like crustaceans that are extremely abundant in polar waters. In the Arctic they are abundant in waters on the Atlantic portion and in the Bering Sea. Krill can constitute up to 45% of zooplankton catches but krill are, more prominent in the Southern Ocean.

>10.000 ind m<sup>-1</sup>  
(William et al., 1983)



Antarctic krill *Euphausia superba* often dominates the zooplankton community in numbers and biomass. Krill are highly influential organisms, capable of grazing as much as 55% of the net primary production and sustaining the functioning of the whole marine ecosystem in the Antarctic (Flores et al., 2012). Many polar organisms, from zooplankton to whales rely on krill as a primary food resource. Its estimated biomass reach >400 million tons (Flores et al., 2012).




Crab-eater seal  
(*Lobodon carcinophaga*)


# Top-down and bottom-up

Top-down processes involve a predator/consumer that exert a control on populations of preys, which are at a lower trophic level. When this control propagate through the trophic web, it originates cascading effects down the food web.

Bottom-up processes, instead refers to changes in the environment, such as nutrient supply, or in the lower trophic levels, that reflects on the upper levels through the trophic web.




# Top-down from invasive predators



*Mnemiopsis leidyi* introduced in 1980 in the Black Sea. Blooms up to 7600 individuals per m<sup>3</sup>. Now spread in Mediterranean Sea, also in the Adriatic Sea. Wide range of tolerance to temperature and salinity. Predator of plankton, including fish eggs and larvae. Introduced with ballast waters in '80s the Black Sea in , native from western Atlantic Ocean.




# Effects on fish populations



Concurrent effects due to pollution, overfishing and finally the introduction of the ctenophores lead fish stocks and fisheries to collapse.


Double effect of predation of *Mnemiopsis*: direct reduction of recruitment of fish due to predation on egg and larval stages, and indirect effect

through competition due to predation on zooplankton




# Trophic cascades

Trophic cascades occur when top-down control causes drastic changes in the ecosystem through effects that propagate downwards the food web (Pace et al. 1999). In ecosystems that are strongly structured by predation, reducing top predator abundance can alter several lower trophic levels.




# Top-down and bottom up

Strong reductions in the abundance of a top predator (cod) has also been reported to cause a trophic cascade in the relatively simple Baltic food chain, involving a subsequent increase in the zooplanktivorous sprat, which in turn affected the biomass, species and stage composition, as well as the vertical distribution of zooplankton (Casini *et al.*, 2008).




# Top-down and bottom up



Changes in the Baltic Sea were primarily bottom-up, strongly structured by salinity, but top-down forcing related to changes in cod abundance also shapes the ecosystem. (Flinkman et al. 1998; Casini et al., 2011)

# Bottom up prevailing

Top-down control is likely to occur in simple trophic webs, where there is a strong predation control on lower trophic levels. This process is more likely to occur in closed basins and coastal areas. In pelagic food webs and open sea the dominant process is likely to be related to bottom up effects. One possibility, in accordance with a predominant view of oceanographers, is that these ecosystems are structured from the bottom-up (resource limitation) and top-down control by oceanic predators is truly rare. (Baum and Worm, 2009)



High degree of connectance among and within species

Prevalence of omnivory and dietary breadth

Ontogenetic diet shifts


Predator diversity may dampen cascading effects

Simultaneous exploitation of predator and prey can inhibit prey responses

Trophic cascade leading to regime shifts could be rare in open ocean ecosystems

# Benthic-pelagic coupling

## Benthic – pelagic coupling



Pelagic or planktonic species lay eggs, or have larval or juvenile stages in benthos

### Life cycles

Benthic species spent part of their life as adult, juvenile or larvae in plankton



Herbivores and predators from the water column feed on benthos

### Trophic webs

Benthic species have adults or juveniles feeding on plankton or on larval - juveniles of nekton



Planktonic species have resting stages in benthos. Organic matter (fecal pellets, dead organisms, etc.) fall on the bottom

### Organic matter

Resting stages disclose and turn back to the plankton. Benthic species feed on particles and could turn in the water column via life cycles



Nutrients and gases reach the bottom and can turn back as living matter or through upwelling

### Biogeochemical cycles




## Benthos



## Life cycles



Benthic species spent part of their life as adult, juvenile or larvae in plankton



Pelagic or planktonic species lay eggs, or have larval or juvenile stages in benthos



Life cycles connect pelagic and benthic domain as, depending on the life stage, species belong to benthos or plankton and pelagos

## Trophic webs

Predator-prey relationships across different compartments connect benthos, nekton and plankton allowing energy flow from the bottom to the water column and viceversa



Herbivores and predators from the water column feed on benthos



Benthic species have adults or juveniles feeding on plankton or on larval -juveniles of nekton







## Organic matter



Planktonic species have resting stages in benthos. Resting stages disclose and turn back to the plankton.



Organic matter (fecal pellets, dead organisms, etc.) fall on the bottom. Upwelling re-suspend nutrients in the upper layer triggering phytoplankton blooms



## Biogeochemical cycles

Nutrients and gases reach the bottom and can turn back as living matter or through upwelling. Shells of calcifying organisms, or silica shells of diatoms also export elements to the sea bottom.

