



Università degli studi di Trieste

LAUREA MAGISTRALE IN GEOSCIENZE

Classe Scienze e Tecnologie Geologiche

Curriculum: Esplorazione Geologica

Anno accademico 2024 - 2025

**Analisi di Bacino e
Stratigrafia Sequenziale (426SM)**

Docente: Michele Rebesco

Modulo 3.5

Contourites and sediment drifts

Docente: **Michele Rebesco**

Outline:

- Introduction
- Sediment drifts
- Sedimentary structures
- Examples from cores
- Facies Model
- Exploration & Production case studies

Introduction

Bottom currents, 'persistent' water current near the sea-floor, are influenced by a series of factors and pervasively affect the seafloor sediments.

Contourites, sediments deposited or significantly affected by bottom current, begin to be perceived as a fundamental component of deep sea depositional systems.


Contourite drifts generally composed of fine sediments, are large sedimentary accumulations produced by bottom currents, which can allow us to reconstruct their evolution and paleoceanography.

The **sedimentary facies** are manifold and the diagnostic sedimentary characteristics are still under discussion.

Bottom currents and contourites are of **great importance** for paleoclimatic reconstructions, investigations on continental slopes and their stability, the exploration of hydrocarbons and polymetallic nodules, definition of the "extended continental shelf", ecological health of deepwater ecosystems, accumulation of microplastics and contaminants .

Elongated contourite drift (contourites)


Asymmetric moat and mound geometry




Rebesco (2005)

Channel levee system (turbidites)


Symmetric gull-wing geometry (asymmetry controlled by Coriolis force)



Bottom currents are capable of building thick and extensive accumulations of sediments (“contourite drifts”). Similarly to channel-levee systems generated by turbidity currents, such large bodies normally have a noticeable mounded geometry, which is generally elongated parallel, or lightly oblique to the margin. Besides this, bottom currents and associated processes generate also a wide range of other depositional and erosional or non-depositional structures at different scales.




Methods to study contourite deposits



Three scale approach

Diagnostic criteria are their **facies and ichnofacies, texture and sequences, microfacies and composition**. **Sedimentary structures** are also “diagnostic indicators”, but for their interpretation its full context should always be considered. **Medium-scale criteria** (hiatuses and condensed deposits, variation in the thickness, geometry, palaeowater depth, geological context) can be definitive. **Large-scale criteria** (palaeoceanographic features and continental margin reconstructions) are essential, but generally more problematic to apply on outcrops.




Courtesy
of D.A.V.
Stow

Recognizing
contourite
deposits in
ancient
sedimentary
series presently
exposed on
land, is a
difficult task.
The distinction
between
contourites and
reworked
turbidites is
controversial.

Seismics characteristics

triple-scale approach that involves 3 “orders of seismic elements”.




Large scale (overall architecture): I-order elements (major changes in current strength and sediment supply): External geometry, Bounding reflectors, Gross internal character.


Medium scale (internal architecture): II-order seismic elements (reflecting smaller fluctuations): lens-shaped, upward-convex geometry; uniform stacking pattern; down-current migration or aggradation; downlapping reflector terminations

Small scale (internal acoustic character): III-order seismic elements: facies analysis (continuous, (sub)parallel, wavy, structureless), and attribute analysis (bedforms).


This updated compilation of **contourite occurrence** was done specifically for the present review, but was subsequently archived and visualised on the Marine Regions website (<http://www.marineregions.org>). It demonstrates that contourite features are ubiquitous within the oceanic basins (different settings and different water masses from the outer shelf to the abyssal plains). The highest numbers of described large contourite depositional and erosional features are located in the western side of the largest oceanic basins, but not exclusively.




ELONGATED, MOUNDED DRIFTS




SHEETED DRIFTS




CHANNEL-RELATED DRIFTS




CONFINED DRIFTS




PATCH DRIFTS




INFILL DRIFTS



FAULT-CONTROLLED DRIFTS



MIXED DRIFTS




All **contourite drifts** are characterised by a variable degree of mounding and somewhat evident elongation


Miramontes et al., 2021. Contourite and mixed turbidite-contourite systems in the Mozambique Channel (SW Indian Ocean): Link between geometry, sediment characteristics and modelled bottom currents. *Marine Geology* 437, 106502.

CONTOURITES


Deposition: Contourite drifts



Winnowing: Sandy contourites



Erosion: Erosional features




Faugères and Stow, 2008: Factors controlling drift location, morphology and depositional pattern

Large-scale features of drifts are controlled by a number of interrelated factors, including


- (1) the bathymetric framework (water depth and morphological context),
- (2) the current conditions (velocity, variability, and Coriolis force),
- (3) the sediment supply (amount, type, source, input, variability),
- (4) interaction with other depositional processes (in time and space),
- (5) sea level and sea-level fluctuations,
- (6) climate and climate change,
- (7) tectonic setting and activity and
- (8) the length of time over which these various processes and controls have operated and varied.

It is not a simple matter to disentangle these various controls as many clearly overlap and are interrelated. Neither is it always certain just what effect a particular control exerts.




Examples of large contourite drifts:

A) Eirik Drift, Greenland margin, northern hemisphere (Hunter et al, 2007);






B) Faro-Albufeira Drift, Gulf of Cádiz margin, northern hemisphere (courtesy of REPSOL Oil);





C) Agulhas Drift, Transkei Basin, southern hemisphere (Niemi et al., 2000).

Large-scale **erosional features** are also common in Contourite Depositional Systems, though less studied with respect to depositional ones. Most commonly they occur just in association with contourite drifts, but may also characterize a broad area of continental slopes. We propose here a reconsideration of the only systematic classification of large-scale erosional features attempted so far (Hernández-Molina et al., 2008; García et al., 2009).





Three-dimensional (3D) coherence volume showing **unidirectionally migrating deep-water channels** (C1 to C7). B) Facies and architecture within unidirectionally migrating deep-water channel 3 (C3). Five channel-complex sets (CCS1 to CCS5), are identified, each of which comprises **bottom-current reworked sands** (BCRS) in the lower part, grading upward into slumps and debris-flow deposits and, finally, into shale drapes. The BCRS are represented by subparallel and high-amplitude reflections with external lens shapes and are systematically nested in the direction of channel migration (Gong et al., 2013; with permission from the AAPG). BCRS from previous turbiditic deposits may represent a pragmatic alternative to the application of conventional turbidite concepts, and a new concept for understanding the origin and predicting the distribution of deep-water sandstones. BCRS frequently contain different seismic facies and sedimentary structures.




Various depositional and erosional bedforms are generated by bottom currents. They are highly variable in terms of sediment composition, morphology and dimension, from decimetres to kilometres. The detection of bedforms can be important for the reconstruction of bottom-current velocity and for geohazard assessment (where velocities can damage seafloor infrastructure, including pipelines and telecommunications cables).

From Stow *et al.*, 2013.

Sediment waves

Rebesco et al (2021) Bottom current-controlled Quaternary sedimentation at the foot of the Malta Escarpment (Ionian Basin, Mediterranean)
Marine Geology, 441, art. no. 106596,



- 1) Creep folds do not display lateral migration and sediments on either flank of individual folds are identical.
- 2) Creep folds show clear evidence of displacement along fault planes and individual reflections cannot be traced across the troughs.
- 3) Creep folds are oriented parallel to the slope and are generally arcuate in plane view, without bifurcation.

- 1) most turbidity current waves occurring in turbidite environments, are situated on channel levees, which are not identified in the “turbidite valley” (Gutscher et al., 2016; San Pedro et al., 2017) at the base of the sediment-starved Malta Escarpment;
- 2) turbidity current waves are scarce in basin floor environments, where their crest alignment is normally slope-parallel or oblique to the regional slope if found on levee backslopes, whereas in this case the crest alignment is almost perpendicular to the Malta Escarpment;
- 3) turbidity current waves usually show a progressive decrease in dimension downslope, which is not observed here;
- 4) in turbidity current wave fields, the stratigraphic interval over which the waves occur often shows a progressive downslope thinning (up to 40–60%) associated to upslope sourcing, which is not observed here;

Type of deposits		Grain size	Characteristics	Sed. Structures	Examples
CLASTIC CONTOURITES	Muddy contourites	5 - 11 ϕ < 0.31 mm	50 % Clays < 15 % of sands ≤ 20 - 30 % of bioclastic and / or carbonate components Homogeneous & highly bioturbated Poor sorting	Rare laminations Bioturbation Indistinct mottled appearance	Rockal Trough (Stow & Faugères, 2008)
	Silty contourites	4 - 8 ϕ 0.063-0.004 mm	40 - 60 % Silts Interbedded between muddy & sandy contourites Poor sorting Sharp to irregular tops and bases	Tractive structures (ripples, sed.waves) Bioturbational mottling Ichonofacies	Gulf of Cadz (Stow & Faugères, 2008)
	Sandy contourites	-1 - 4 ϕ 2 - 0.063 mm	Sheeted to wedge bedding Well-sorted deposits, but can be poor to moderate Mixed siliciclastic / biogenic composition Heavy mineral concentration Both positive & negative grading Gradational or erosive contacts	Tractive structures (e.g.: horizontal lamination, cross-lamination, ripples, etc) Bioturbation (sub.vertical burrows) Massive layers (structureless) Erosional or gradational contacts	Gulf of Cadz (Stow & Faugères, 2008)
	<i>Bottom current reworked sands (BCRS)</i>	Normally does not exceed fine sands	From previous turbiditic deposits Rhythmic layers Lenticular bedding Well sorted Coarsening upward sequences Sharp to gradational bottom contact and sharp (nonerosional) upper contact!	Tractive structures (e.g.: horizontal lamination, low-angle cross lamination; mud-offshoots in ripples, mud-drapes, flaser, etc)	Gulf of Cadz (Stow & Faugères, 2008)
	Gravel contourites	< -1 ϕ > 2 mm	Winnowing & erosion (channels, moats, etc) Irregular layer and lenses Poorly to very poorly sorted	Sandy gravel lag	Laena (Stow & Faugères, 2008)
VOLCANOCLASTIC CONTOURITES		Mud, silt or sands	Similar to the siliciclastic facies Composition is dominated by volcanoclastic material	Similar to the clastic contourites	Hawai (Paga-barradas, A., Pers. Comm.)
SHALE-CLASTS OR SHALE-CHIP LAYERS		Shale clasts generally mm in size	Developed in muddy & sandy contourite facies From substrate erosion by strong bottom currents Burrowing on the nondeposition surface	Clast axes sub-parallel both to bedding and to the current direction	Gulf of Cadz (Stow & Faugères, 2008)
CALCAREOUS CONTOURITES	Calcareous muddy & silty contourites	> 4 ϕ < 0.063 mm Silty clay to clayey silts	>70 % of bioclastic and / or carbonate components Dominant biogenic input Poorly sorted Distinct sand-size fractions (biogenic particles) Composition: pelagic to hemipelagic, including nanofossils & foraminifers as dominant elements Admixture of siliciclastic or volcanoclastic material	Bedding is indistinct, but may be enhanced by cyclic variations in composition / grain size. Bioturbation	Orcaia Basin (Bosellini & Cerrito, 2003)
	Calcareous sandy contourites	-1 - 4 ϕ 2 - 0.063 mm Sands	Equivalent of sandy contourites Both well-sorted to poorly sorted Particles from pelagic, benthic, off-shelf & off-reef sources Admixture of siliciclastic, volcanic & siliceous material	Thin-bedded cross-laminated foraminera contourite. Lenticularity Hardgrounds, non de positional surfaces Bioturbation & burrowing	Gulf of Cadz (Stow & Faugères, 2008)
	Calcareous gravel lag contourites	< -1 ϕ > 2 mm Gravel	Clasts or chips derived from erosion of the substrate		Gulf of Cadz (Stow & Faugères, 2008)
SILICEOUS BIOCLASTIC CONTOURITES		Mud, silt or sands	Rich in diatomaceous & radiolarian material	Laminated and / or cross-laminated sands	Hala-Bahama Basin (DSDP Site 534, Sheridan et al 1983)
CHEMOGENIC CONTOURITES	Manganiferous contourites		Manganiferous or ferro-manganiferous horizons Areas with ferro-manganese nodules & pavements	Bioturbation & burrowing	Illian Basin (Gornitz et al, 2003)
	Chemogenic gravel-lag contourites		Deep-water chenoherms (chemical - biogenic precipitates) of metal - carbonate chimneys, mounds & encrustations Winnowed and aligned into chemogenic gravel-lags	Strewn od debris Alineation of gravel-lag	Illian Basin (Gornitz et al, 2003)

Stow and Faugères (2008) + BCRS from Shanmugam, 2012

Martín-Chivelet et al., 2008 Traction structures

40 years of controversy

1950s: discovery of current ripples = establishment of the contourite facies

1970s: discovery of fine-grained turbidites = seed of controversy since similar sedimentary structures of contourites.

1980s: concept that bioturbation destroy traction structures = reinterpretation of contour-current deposits as fine-grained turbidites

1990s: few workers provided convincing evidence of traction structures = most workers reject this criterium

Setting the stage for sedimentary structures in contourites

In the time interval between deposition and significant lithification, burrowing can be sufficiently intense to destroy previous traction structures.

Traction structures are abundant on recent ocean floors.

Traction structures are more abundant and easily preserved in sandy contourites.

Thermohaline circulation in older, greenhouse times, was probably driven by active sinking of saline waters in intertropical seas

studies from boreholes or small outcrops might be biased by partial observation of bedform geometry, internal architecture and lateral arrangement.


turbidity currents are sediment gravity flows contour currents are water flows.

Various **sedimentary structures** have been described for contourites in present and ancient deposits (Martín-Chivelet *et al.*, 2008). However, in areas of intense bioturbation from benthic activity, the preservation potential of some of these structures can be low.



MAIN TYPES OF PRIMARY SEDIMENTARY STRUCTURES IN CONTOURITE DEPOSITS

Sketch	Sed. structures	Dominant grain size	Enviromental implications
1 cm	Horizontal or sinusoidal lamination, stripped, fine-grained deposits; "wispy" lamination	Fine sand, silt & mud < 2 ϕ < 0.250 mm	Low current strength Predominance of deposition from suspension
1 cm	Lenticular bedding starved ripples	Fine sand, silt & mud < 2 ϕ < 0.250 mm	Alternating flow conditions, low to moderate current strength, winnowing
1 cm	Wavy bedding, flaggy chalks	Fine sand, silt & mud < 2 ϕ < 0.250 mm	Alternating flow conditions, low to moderate current strength
1 - 5 cm	Flaser bedding, mud offshoots	Fine sand to silt 8 - 2 ϕ 0.004 - 0.250 mm	Alternating flow conditions, Current speed = 10 - 40 cm / s
1 - 5 cm	Climbing ripples (subcritical to supercritical)	Very fine to medium sands 4 - 1 ϕ 0.063 - 0.5 mm	Current speed = 10 - 40 cm / s High suspension load
10 - 50 cm	Large-scale cross-bedding, megaripples, dunes, sandwaves	Medium sands 2 - 1 ϕ 0.250 - 0.5 mm	Current speed = 40 - 200 cm / s Barchan dunes usually form at 40-80 cm / s
1 cm	Parallel lamination (upper stage plane beds), presence of primary current lamination	Very fine to medium sands 4 - 1 ϕ 0.063 - 0.5 mm	Current speed = 40 - 200 cm / s
1 cm	Minor erosive surfaces, mud rip-up clasts, upper sharp contacts	Sand, silt & mud < -1 ϕ < 2 mm	Alternating flow conditions, low to moderate current strength
1 - 5 cm	Sole marks: flutes, obstacle scours & longitudinal scours, cut & fill structures	Sand, silt & mud < -1 ϕ < 2 mm	Flow speed peaks
5 cm	Longitudinal ripples	Coarse sandy muds (20 % sand)	Low current speed = 2 - 5 cm / s Winnowing
1 - 10 cm	Bioturbation (strongly variable)	Sand, silt & mud	Low current speed Strong paleoecological control, Low to moderate accumulation rates
3 - 20 cm	Normal & reverse grading at different scales and within different types of deposits	From coarse sand to mud Usaully fine sand, silt & mud	Gradual changes in flow strength
0.1 - 2 cm	Pebble lags, furrows	Coarse sand, microconglomerate	Current speed over 200 cm / s

Clay and / or mud
 Silt
 Sand
 Pebbles & cobbles



Most of these structures are also present in other deep-water deposits (e.g. turbidites), but some have been suggested to be a clear diagnostic feature for bottom-current deposits, such as: *negative grading*; *longitudinal triangular ripples*; and *double mud layers and sigmoidal cross-bedding*, which are unique to deep-water tidal deposits in submarine canyons (Shanmugam, 2006; 2012).




Sedimentary structures in bottom-current reworked sands (BCRS): A) Discrete thin sand layers with sharp upper contacts; B) Rhythmic layers of sand and mud, inverse grading, and sharp upper contacts; C) Horizontal lamination with gradational upper contact; D) Convex-up and concave-up laminae; E) Flaser bedding; and F) Double mud.



Stow and Faurgères, 2008: Silty-Muddy contourites

Homogeneous, featureless, poorly bedded units in some cases showing cm-dm banding marked by subtle colour and core logging changes.

Highly bioturbated (mottled with burrows). Rare primary indistinct lamination (marked by colour change and/or irregular winnowed concentrations of coarser material). Rarely, remnants of thin cross-laminated beds. Silty-clay grain size and poor sorting, with dominantly siliciclastic composition with some biogenic fraction. Either local and far-travelled components



Sandy contourites

Either as thin irregular layers and much thicker units within the finer grained facies, may display either distinct or gradational contacts. Thoroughly bioturbated, appear massive (structureless). The mean grain size normally does not exceed fine sand (apart from coarser grained horizons and lags), and sorting is mostly poor to moderate, in part due to bioturbational mixing.



Both positive and negative grading may be present. A mixed siliciclastic–biogenic composition is typical, with evidence of abrasion, fragmented bioclasts and iron-oxide staining.




Laminated sandy contourites



Less common than their bioturbated counterparts and have been rarely documented, but do occur where high-energy (high-velocity) bottom currents are especially dominant and larger-scale bedforms (e.g. dunes) are evident on the sea floor. The few examples observed to date are thick to very thick-bedded and distinctly laminated. The lamination is relatively broad and diffuse, enhanced by slight colour variation, and parallel at the scale of the cores, although this may also be part of large-scale cross-bedding. Bioturbation is rare, but large sub-vertical burrows have been noted. The mean grain size is medium-grained sand, with moderately good sorting. The sediment has a mixed siliciclastic/biogenic composition, with evidence of abrasion, fragmented bioclasts and iron-oxide staining.

Gravel-rich contourites and gravel-bearing contourites




Common in drifts at high latitudes (ice-rafted debris). Under relatively low-velocity currents, IRD remains and is not subsequently reworked. This facies is often indistinguishable from glaciomarine hemipelagites.

Concentration of the coarser fraction occurs under higher-velocity currents and more extensive winnowing, yielding irregular layers and lenses of poorly to very poorly sorted, sandy gravel-lag.

Similar coarse-grained concentrations and gravel pavements are locally developed in response to high-velocity bottom-current activity in shallow straits, narrow contourite moats and passageways.

The creation of a definitive **facies model** for contourites poses major challenges.


The standard contourite facies model sequence (Stow and Faugères, 2008) was first proposed by Gonthier *et al.* (1984) and was derived from the Faro Drift within the middle slope of the Gulf of Cádiz.



Bioturbation



Discontinuous silt lenses




Gradational contact



Sharp (irregular) contact

This model implies a cyclic trend, encompassing 3 main facies, linked to variation in contour-current velocity.


Facies and facies sequences associated to contourites vary greatly, making any singular, systematic characterisation of facies rather difficult for the moment. Stow et al. (2002) slightly modified the standard sequence by using five principal divisions (C1–C5), and Stow and Faugères (2008) later proposed a model for partial sequences, which are equally or more common than the full bi-gradational sequence.




Examples of the principal sedimentary facies for the contourites recovered during **IODP Expedition 339** (Hernández-Molina et al., 2013).

Preliminary results are in agreement with the previously proposed idea that there is a greater variety of facies sequences for bottom current deposits than what is presently represented in the most commonly accepted contourite facies model. Additionally, remarkable interactions between contourite and turbidite processes have been reported that are completely new and different from the current facies models.


A Bigradational sequences




B Base-cut-out type



C Top-cut-out type



D Sandy contourites




Examples of the principal sedimentary facies for the contourites recovered during **IODP Expedition 339** (Hernández-Molina et al., 2013).

The main criticism for considering the Faro Drift deposits as the standard contourites facies sequence relates to two facts: this drift is predominantly muddy, and it is located in the distal part of a huge CDS. Moreover, other facies in other parts of the same depositional system have recently been reported, but it is difficult to apply the conceptual model to them.

Most of the contacts between the classical contourite facies (mottled, fine sand, and coarse sand) are sharp rather than transitional.

CADKS18



Gravelly contourite in the Gulf of Cádiz. Core log, X-ray, grain size, and indurated thin sections under natural light and fluorescence. (Mulder et al., 2013).

Preliminary contourite facies tract (?)

In the Campos Basin, bottom currents played a major role since late Cretaceous in reworking and redistributing turbidite fine sands derived from basin margins.

Currently available models for deep-water sedimentation are inadequate to describe and interpret the complexity of depositional patterns developed by the interaction of bottom currents and sediment gravity flows.



(CFA) Muddy fine sand with abundant mudstone clasts



(CFB) M-thick well-sorted horizontally laminated fine and very fine sand




Flow direction



(CFC) M-thick well-sorted fine and very fine sand with large ripples with internal sigmoidal laminae

(CFD) Alternating cm-thick packages of ripple-laminated fine-grained sand and bioturbated muddier units with sand streaks



Flow direction

(CFE) Cm-thick packages of lenticular rippled sand and sand streaks alternating with mudstones. Bioturbation is very common. These thin units strongly resemble contourite facies cycles of the classic Stow's model (Stow et al., 2002)




(CFF) Highly bioturbated terrigenous, mixed and biogenic (calcareous) mudstones



Mutti & Carminatti, 2015


Photos of Contourite Facies A-F



E & P case study: Santos Basin

Location map


Location map of Santos Basin within Brazilian Margin where sediment transported in suspension by turbidity currents was removed by bottom currents resulting in Contourites and Turbidites which are highly prolific oil reservoirs



Santos Basin sandy contourites


Example of barchan dunes, sand waves, sand furrows, sand furrows and ribbons in plan view: evidence of bottom current action

(Mutti et al., 2014)



Santos Basin: seismic section through sandy contourites


Seismic section showing that these large bedforms migrated on a flat sea-bottom under the action of high velocity bottom currents



(Mutti et al., 2014)

Santos Basin: well log and petrophysical properties


The well log shows the large thickness of the sand bodies and the photos show that sandstone is very well sorted (low depositional matrix) with good porosity and permability thanks to winnowing away of mud.



(Mutti et al., 2014)

Campos Basin Location map

Location map of Campos Basin within Brazilian Margin where there are clear evidence of bottom-current reworking of hydrocarbon-bearing turbidites



(Mutti et al., 2014)


Campos Basin: Contourites occur primarily in 3 settings:

within channels where bottom currents moved up and down




COARSE-GRAINED TURBIDITE BEDS REWORKED BY TIDAL BOTTOM CURRENTS (HYBRID FACIES ASSOCIATION)

in open slope settings, with quasi-permanent geostrophic thermohaline flow



CYCLICALLY STACKED THIN-BEDDED CONTOURITES

within channels or in turbidite fan remnants




CURRENT-LAMINATED SANDY CONTOURITES

patterns controlled by local and regional topography that moved large amounts of fine sand through migrating large-scale sand-dune fields and sand waves, and smaller bedforms.


(Mutti et al., 2014)

Espirito Santo Basin Location map



Location map of Espirito Santo Basin within Brazilian Margin where the preliminary contourite system tract of Mutti and Carminatti (2012) was conceived






(Mutti et al., 2014)



Espirito Santo: Basin hybrid facies associations

(Mutti et al., 2014)


Turbidite sands winnowed, eroded and redistributed by deep-marine bottom currents to form contourite sands which are difficult to explain with currently turbidite-dominated models for deep-water sedimentation




FROM MUTTI, 1992

Tanzania Location map




Location map of the Tanzanian margin where Hybrid turbidite-contourite systems have been identified



(Sansom, 2018)



Tanzania: migrating channel-levee complexes





In response to flow-stripping of the suspended load of the turbidity current, the northern levee migrates upstream with respect to contour current, and the channel displaced to the south

(Sansom, 2018)

Mozambique: Location map


(Fonnesu et al., 2019)



Location map of Area 4 off the coast of Northern Mozambique. Eocene and Oligocene Mamba and Coral systems are indicated including their straddling equivalents in the Area 1


Mozambique: Ocean floor bathymetry

(B) Shaded relief map overlain by bathymetry of modern deep-water fans and deep undercurrents flowing northwards along the continental slope.





The basin is located in the offshore of northern Mozambique, within the Mozambique Channel.

(D) Detail of seafloor features including large-scale scour marks (flutes and crescent) created by northerly directed bottom currents




Seismic section through the Coral sequence




Main stratigraphic surfaces (above) and seismic facies interpretation (below). Overall southwards migration and offset stacking pattern with channel complexes flanked by prominent asymmetric drift-mounds on the N side of the fans.


The reservoir shows a systematic internal organization linked to 4th and 5th order base level



(Fonnesu et al., 2019)



Block-diagram of depositional processes affecting the Coral system



(Fonnesu et al., 2019 after Fonnesu, 2013)

Most of the **future perspectives** in contourite research strongly depend on continuous technological advances.

Use of numerical or sand-box modelling, indurated thin sections, Ichnological Digital Analysis Images Package, CT scanning, HR 3D seismics, observations from AUV, seismic oceanography, fingerprinting of water masses using isotopic tools are steadily expanding techniques.


A more intensive collaboration between physical oceanographers and geologists!!!

Scale will be an especially important factor:

- high-resolution to elucidate the relationship currents and smaller contourite deposits;
- increased resolution to detail spatial and temporal variability within a single deposit;
- larger-scale perspective on CDS sharing the same basin, water masses and time scale.

The advances expected in contourite research should lead to the establishment of better diagnostic criteria for contourite identification.

Future discoveries?



CONCLUSIONS

Contourite processes are not as simple as initially thought.

Contourite nomenclature might need to be reconsidered.


New facies models must be established.

More work is needed to understand sandy contourites.


Integrated studies will be essential for an holistic perspective.

Pervasiveness of bottom-water circulation to be reconsidered.

Exercise: Shan Liu, F. Javier Hernández-Molina, Gemma Ercilla, David Van Rooij, 2020, Sedimentary evolution of the Le Danois contourite drift systems (southern Bay of Biscay, NE Atlantic): A reconstruction of the Atlantic Mediterranean Water circulation since the Pliocene, *Marine Geology* 427, 106217



Rebesco et al (2002) Geological Society Memoir, 22 (1), pp. 353-371.



Seismic record of a cyclic turbidite-contourite system in the Northern Campos Basin, SE Brazil
 Pandolpho et al. 2021, Marine Geology

Depositional Environment	Seismic Facies	Attributes	Line Drawing	Description	Depositional Architecture
Upper and mid-slope				Sigmoidal clinoforms, medium to low amplitude facies	Plastered Contourite Drifts
Mid- and Lower slope and SP Plateau				Continuous to discontinuous, high-amplitude reflections, V- or U- shaped	Basal Lag of Channels
Mid-slope				Discontinuous, wavy high- to low- amplitude reflections	Sediment Waves
Lower slope, continental foot and SP Plateau				Chaotic to transparent, discontinuous, low amplitude facies with deformed internal architecture	Mass Transport Deposits (MTD)