





MOLECULAR NEUROPHYSIOLOGY -lesson 1-

Prof. Giada Cellot
cellot@sissa.it



The plasticity of neuronal circuits: synaptic changes and co-related behavioral aspects

- Hebbian (NMDA mediated) LTP - Spike timing dependent plasticity
- Anti-Hebbian LTP

- Functional implications of hippocampal synaptic plasticity: memory and learning
- Optogenetic manipulation of synaptic plasticity: impact on behavior



- Synaptic plasticity in the amygdala
- Associative learning: the fear conditioning paradigm

- Synaptic plasticity during the development: when GABA is depolarizing

| | |
|----------|---------------------------------|
| Lesson 1 | November 19 th h9-11 |
| Lesson 2 | November 26 th h9-11 |
| Lesson 3 | December 3 rd h9-11 |
| Lesson 4 | December 10 th h9-11 |

NEURAL ACTIVITY CARVES CONNECTIVITY DURING THE MATURATION OF THE SYNAPTIC CIRCUIT

At specific synapses:



Synaptic reinforcement is obtained by repeated activation of the synapse

SYNAPTIC PLASTICITY: EXPERIENCE DEPENDENT change in the EFFICACY of synaptic communication between neurons → Long term potentiation (LTP)

“When an axon of cell A is near enough to excite a cell B and repeatedly or persistently takes part in firing it, some growth process or metabolic change takes place in one or both cells such that A's efficiency, as one of the cells firing B, is increased”

D. O. Hebb, 1949





CELLS that FIRE TOGETHER, WIRE TOGETHER

Synaptic plasticity in the hippocampus can be studied *in vitro*


HIPPOCAMPAL
SLICES

- Composed of defined populations of identifiable neurons
- Suitable for investigation (electrophysiology)
- Easy pharmacological manipulation, limitation of animals used in research


Brain is dissected in 300-400 μm thick slices that can be recorded for hours in oxygenated artificial cerebrospinal fluid



Phases of a LTP experiment



1. Baseline responses
(low frequency stimulation
0.5 Hz)



3. Post-induction
responses
(low frequency
stimulation 0.5 Hz)


2. Protocol to induce potentiation = INDUCE STRONG STIMULATION OF SYNAPSES (Electrical stimulation: Tetanus: 1s @100 Hz; or theta burst stimulation: interval of 200 ms)

1. **Early-LTP:** kinase dependent LTP






Potentiation is downregulated
after few seconds

2. **Late-LTP:** protein synthesis dependent LTP



Potentiation is downregulated
after 5-6 hours

HEBBIAN LTP mechanisms





LTP INDUCTION: always postsynaptic
 LTP EXPRESSION: pre- or/and postsynaptic

LTP induced morphological modification of synapses

DENDRITIC SPINES: tiny protrusions from dendrites, form functional contacts with neighboring axons of other neurons (Smith et al, 2014)

- Actin cytoskeleton
- **POST SYNAPTIC DENSITY (PSD):** high density packed ion channels, receptors, kinases anchored by scaffolding proteins


Acute hippocampal slice after TBS




Toni et al, Nature, 1999

DENDRITIC SPINES

Hippocampal cultures exposed to chemical LTP




Classification
based on head /neck size




Pchitskaya & Bezprozvanny, Front Syn Neurosci, 2020
doi: 10.3389/fnsyn.2020.00031

NMDA receptor as detector of coincidence




COOPERATIVITY, ASSOCIATIVITY, INPUT SPECIFICITY can be all explained on the assumption that the induction of LTP requires pre-synaptic activation with **COINCIDENT ACTIVATION** of post-synaptic cell



Lynch et al., 1983



NMDA antagonists or Buffering intracellular calcium with EGTA block the induction of LTP

Hebb's postulate:
CELLS that FIRE TOGETHER, WIRE TOGETHER




LTP can be induced by PAIRING pre- and post-synaptic NEURONS

Not only high frequency stimulation (tetanus: 1 s @100 Hz) of the afferent fibers can be used to induce LTP, but also **PAIRING PROCEDURE** in which pre-synaptic stimuli at low frequency (1-3 Hz) are paired with postsynaptic depolarization.



Bliss & Lomo, 1973







PATCH CLAMP TECHNIQUE

It allows to set the potential of the recorded neuron at precise values




Spike Timing-Dependent Plasticity (STDP)

The pairing procedure becomes very efficient when postsynaptic neuron fires later respect to presynaptic cell (in a time window <20 ms)




Modified from Markram et al., Science, 1997

STDP and back-propagating action potentials




Stuart et al., Trends Neuroscience, 1997

- Action potentials can back-propagate in the dendritic tree, as a wave of depolarization.
- The physiological meaning of backpropagation is to have a “trace” of memory of recent cell history.
- Thus, if presynaptic firing coincides with postsynaptic AP backpropagation, LTP can occur



“HEBBIAN” LTP



Plasticity that occurs when pre-synaptic cell firing coincides with postsynaptic neuron depolarization





“anti-HEBBIAN” LTP

Plasticity that occurs when pre-synaptic cell firing coincides with postsynaptic neuron hyperpolarization (prevented by postsynaptic depolarization)

THE “anti-HEBBIAN” LTP




WHOLE CELL CONFIGURATION




PERFORATED PATCH CONFIGURATION
(by using antibiotics such as gramicidin)
Not to dialyze intracellular milieu

THE “anti-HEBBIAN” LTP




Calcium permeable AMPA receptors

- More abundant in interneurons
- Do not express the GluR2 subunit (-> permeable to calcium)
- At depolarized potential, the pore of the channel is blocked by polyamines (spermine, spermidine) present in the cytoplasm of neurons
- Also called “rectifying AMPAR”



Role of metabotropic glutamatergic receptors in LTP

“HEBBIAN” LTP ↔ “anti-HEBBIAN” LTP



- G-proteins
- Activation of phospholipase C
- Calcium release from internal stores