

Ordinary (0-form) symmetries

Symmetry transf. in QFT

$$\langle U_g(\Sigma) \Phi^i(y) \rangle = R(g)^i_j \langle \Phi^j(y) \rangle$$

Since the sym. generators are CONSERVED / COMPUTE WITH HAMILTONIAN,
 $U_g(\Sigma)$ is "topological" (as we will see)

In Field Theory, if S is invariant under sym group G , then
 there exists a CONSERVED CURRENT $\partial_\mu j^\mu = 0$

j.s.t. if we take local transf

$$S[\Phi^i + \epsilon(x) M^i_j \Phi^j] - S[\Phi^i] = - \int \epsilon(x) \partial_\mu j^\mu(x) \quad (*)$$

\uparrow generator \rightarrow

In QFT \rightsquigarrow WI

then is a current associated with any gen.

$$i \langle \partial_\mu j^\mu(x) \Phi^i(y) \rangle = \delta^4(x-y) M^i_j \langle \Phi^j(y) \rangle \quad (o)$$

Dim. $\langle \partial_\mu j^\mu(x) \Phi^i(y) \rangle = N \int \mathcal{D}\Phi \partial_\mu j^\mu(x) \Phi^i(y) e^{iS[\Phi]} =$

$$\stackrel{(*)}{=} -N \int \mathcal{D}\Phi \frac{\delta}{\delta \epsilon(x)} S[\Phi^k + \epsilon(x) M^k_j \Phi^j] \Big|_{\epsilon=0} \Phi^i(y) e^{iS[\Phi]} =$$

$$= -\frac{1}{i} \frac{\delta}{\delta \epsilon(x)} \left(N \int \mathcal{D}\Phi \Phi^i(y) e^{iS[\Phi^k + \epsilon M^k_j \Phi^j]} \Big|_{\epsilon=0} \right)$$

$\equiv \Phi^i \rightarrow \Phi^k = \Phi^i - \epsilon M^k_j \Phi^j$

$$= i \frac{\delta}{\delta \epsilon(x)} N \int \mathcal{D}\Phi \left(\Phi^i(y) - \epsilon(y) M^i_j \Phi^j(y) \right) e^{iS[\Phi]} \Big|_{\epsilon=0}$$

$$= -i \delta^4(x-y) M^i_j \langle \Phi^j(y) \rangle //$$


it's not
 ANOMALY:
 $\partial_\mu^x \langle j^\mu(x) j^\nu(x_2) \rangle$
 $= \frac{i}{\hbar} \epsilon^{\alpha\beta} \partial_\beta \delta(x_1 - x_2)$

$\int dx_1 \downarrow = 0$

We can now integrate the WI (o) and obtain

$$i \langle [Q, \Phi^i(y)] \rangle_{\text{eq. time}} = M^i_j \langle \Phi^j(y) \rangle \quad (\text{canonical quantization})$$

Dim. Integrate $i \langle \partial_\mu j^\mu(x) \Phi^i(y) \rangle = \delta^4(x-y) M^i_j \langle \Phi^j(y) \rangle$
over the domain $\Omega_\Sigma \equiv [y^0 + \epsilon, y^0 - \epsilon] \times \mathbb{R}^3$




$$\int_{\mathbb{R}^3} \partial_i j^i = 0$$

$$\begin{aligned} \text{LHS: } \int_{\Omega_\Sigma} d^4x \partial_\mu j^\mu(x) &= \int d^3x (j^0(y^0 + \epsilon, \bar{x}) - j^0(y^0 - \epsilon, \bar{x})) = \\ &= Q(y^0 + \epsilon) - Q(y^0 - \epsilon) \end{aligned}$$


$$\begin{aligned} \langle (Q(y^0 + \epsilon) - Q(y^0 - \epsilon)) \Phi^i(y) \rangle &= \langle 0 | T(Q(y^0 + \epsilon) - Q(y^0 - \epsilon)) \Phi^i(y) | 0 \rangle = \\ &= \langle [\hat{Q}(y^0), \hat{\Phi}^i(y)] \rangle \quad // \end{aligned}$$

How does it work for extended objects?



Rewriting ordinary sym. transf.

$$i \langle Q(\Sigma) \Phi^i(y) \rangle = \text{Link}(\Sigma, y) M^i_j \langle \Phi^j(y) \rangle$$



Charge Q on a time slice is generalized (Euclidean signature) to a charge $Q(\Sigma)$ on a 3d CLOSED subspace Σ

$$Q(\Sigma) \equiv \int_\Sigma *j$$

The commutation relations to LINK of Σ and y .
How do we derive this relation?

↓

Let's integrate $w_1(\cdot)$ on Ω_Σ


$$\text{LHS: } \int_{\Omega_\Sigma} \partial_\mu j^\mu dx = \int_{\Omega_\Sigma} d*j = \int_\Sigma *j = Q(\Sigma)$$

$$\hookrightarrow i \langle Q(\Sigma) \Phi^i(y) \rangle = \underbrace{\int_{\Omega_\Sigma} d^4 \delta^4(x-y)}_{\text{Link}(\Sigma, y)} M^i_j \langle \Phi^j(y) \rangle$$

← TOPOLOGICAL INVARIANT

Also this is
TOPOLOGICAL
due to conserv. law:

under a contin. deform. $\Sigma \rightarrow \Sigma' = \Sigma + \partial\Omega_0$ $y \in \Omega$



Inside Ω_0 there is no INSERTION of local operators

\uparrow in correlators
 $= 0 \iff \partial_{\mu} j^{\mu} = 0$

$$Q(\Sigma') = Q(\Sigma) + \int_{\partial\Omega_0} *j = Q(\Sigma) + \int_{\Omega_0} d*j = Q(\Sigma)$$

By exponentiating infinitesimal generators:

$$\langle U_g(\Sigma) \Phi^i(y) \rangle = R(g)^i_j \langle \Phi^j(y) \rangle \quad (\text{if LINKED})$$



charged operator (0-dim \rightsquigarrow 0-form sym.)

TOPOLOGICAL unitary operator depending on $g \in G$ & Σ

$$\left[\frac{d}{d\alpha} U_{e^{i\alpha}}(\Sigma) \Big|_{\alpha=0} = i Q(\Sigma) \right]$$

[The insertion of the top. op. $U_g(\Sigma)$ can be removed at the cost of transforming all the local operators inside Σ . Equivalently, we can say that if we deform the support passing through one loc. op. position, we act on it by g .]

Discrete symmetries




- $g \in G$ discrete $\leftarrow [U_g, P^\mu] = 0$
- U_g : unitary operator commuting with Hamiltonian & momentum

$$\langle U_g \Phi^i(y) U_g^{-1} \rangle = R(g)^i_j \langle \Phi^j(y) \rangle$$

\uparrow
 related to a TOPOL. OPERATOR $U_g(\Sigma)$ sit.

$$\langle U_g(\Sigma) \Phi^i(y) \rangle = R^i_j(g) \langle \Phi^j(y) \rangle \quad (\text{if linked})$$

• $[U_g, P^\mu] = 0 \Rightarrow U_g$ can continuously move, i.e. is Topol.



• $U_g(\Sigma) U_{g'}(\Sigma) = U_{gg'}(\Sigma)$

Summary ORDINARY SYMMETRIES

$g \in G \iff$ Topol. op. $U_g(\Sigma)$ G cont/disc.

with

$\langle U_g(\Sigma) \Phi^i(y) \rangle = R^i_j(g) \langle \Phi^j(y) \rangle$ (if linked) (*)

\uparrow topological Σ is $(d-1-0)$ -obj.
 \nwarrow not necessary topol. $\Phi^i(y)$ is $\underline{0}$ -dim \Rightarrow "0-form symmetry"

Reduced the problem of finding symmetries to the problem of finding topological operators.

This can be generalized to topological $(d-1-p)$ -op.

i.e. p -form symmetries (we are now going to illustrate an example with $p=1$).

Observation: interpret (*) as

$\langle U_g(\Sigma) \Phi^i(y) \rangle = R^i_j(g) \langle \Phi^j(y) \rangle + 0 = R^i_j(g) \langle \Phi^j(y) \rangle + \langle U_g(\Sigma') \Phi^i(y) \rangle$
 with $\text{Link}(\Sigma', y) = 0$



1-form symmetries in Maxwell theory

$$S[A] = -\frac{1}{2e^2} \int F \wedge *F = -\frac{1}{4e^2} \int d^4x F^{\mu\nu} F_{\mu\nu} \quad (*)$$


$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad A_\mu \quad \text{COMPACT } U(1) \text{ gauge field}$$

$\int_{\Sigma_2} F \in 2\pi\mathbb{Z}$
 \uparrow
 proper normalization of $U(1)$ connection
 $\rightarrow e$ is a minimal coupling.
 \Rightarrow ELECTRIC CHARGES ARE QUANTIZED
 (seen through gauge transf. not-connected to identity)

Gauge transformations:

$$A \rightarrow A + \lambda \quad \text{with } \lambda \text{ a (globally defined) closed 1-form (locally } \lambda = d\alpha)$$

- $\alpha(x)$ is a $U(1)$ parameter, meaning that the well defined local transf. are $e^{i\alpha(x)} \rightarrow \alpha \sim \alpha + 2\pi$ (*)
- let's take an S^2 and integrate F over it



$$\int_{S^2} F = \int_{S^+} F + \int_{S^-} F = \int_{S^+} A^+ - \int_{S^-} A^- = \int_{S^1} \lambda =$$

$$= \alpha(2\pi) - \alpha(0) = 2\pi n$$

\uparrow
 $n \in \mathbb{Z}$
 $U(1)$ gauge group (*)

\Rightarrow For $U(1)$ gauge groups

$$\int_{\Sigma_2} F \in 2\pi\mathbb{Z} \quad \forall \Sigma_2 \quad \neq \quad \int_{\gamma} \lambda \in 2\pi\mathbb{Z} \quad \forall \gamma$$

If $\int_{\gamma} \lambda = 0 \rightarrow \lambda = d\alpha$ e α e-hen def. \leadsto small gauge transform.

\Rightarrow For $U(1)$ gauge group, Wilson lines have integer charge $n \in \mathbb{Z}$:
 $W_n(\gamma) = e^{in \int_{\gamma} A} \rightarrow e^{in \int_{\gamma} \tilde{\lambda}} W_n(\gamma)$

S-duality

Let's rewrite the action as

$$S[F, \tilde{A}] = \frac{1}{2e^2} \int F \wedge *F + \frac{1}{2\pi} \int F \wedge d\tilde{A}$$

\nearrow now indep. variable
 \uparrow new 1-form
 \nwarrow Lagrange multiplier

• e.o.m. of \tilde{A} : $dF = 0 \rightarrow$ Bianchi id.

• e.o.m. of F : $\frac{1}{e^2} *F = \frac{1}{2\pi} \tilde{F}$ with $\tilde{F} = d\tilde{A}$

\Rightarrow Integrating over $\tilde{A} \rightarrow S[A] = \frac{1}{2e^2} \int F \wedge *F$ with $F = dA$

Integrating over $F \rightarrow S[\tilde{A}] = \frac{1}{2\tilde{e}^2} \int \tilde{F} \wedge * \tilde{F}$ with $\tilde{F} = d\tilde{A}$
 and $\tilde{e}^2 = \frac{4\pi^2}{e^2}$

\leadsto Example of DUALITY: the same theory has two equivalent presentations.

1-form symmetries

The e.o.m. of (*) are

$$\frac{1}{e^2} \partial_\mu F^{\mu\nu} = 0 \quad \text{and} \quad \partial_\mu (\star F)^{\mu\nu} = 0 \quad \star F_{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu\sigma\rho} F^{\sigma\rho}$$
$$\uparrow \qquad \qquad \qquad \uparrow$$
$$d\star F = 0 \qquad \qquad \qquad dF = 0$$

F and $\star F$ are two-forms that are closed
 \Rightarrow they define two 1-form symmetries with

currents $J_e = \frac{1}{e^2} F$ and $J_m = \frac{1}{2\pi} \star F$

• The correspondingly CONSERVED CHARGES are

- Electric flux

$$Q_e(\Sigma_2) = \frac{1}{e^2} \int_{\Sigma_2} \star F \sim \int_{\Sigma_2} \vec{E} \cdot d\vec{S} \leftrightarrow U(1)_e^{(1)}$$

- Magnetic flux

$$Q_m(\Sigma_2) = \frac{1}{2\pi} \int_{\Sigma_2} F \sim \int_{\Sigma_2} \vec{B} \cdot d\vec{S} \leftrightarrow U(1)_m^{(1)}$$

• Under S-duality $J_e \leftrightarrow J_m$

$$Q_e \leftrightarrow Q_m$$

- Both $Q_e(S^2)$ & $Q_m(S^2)$ are TOPOLOGICAL under contin. deformations of Σ_2 .

\Rightarrow there should be corresponding symmetries (whose related conserved quantities are the topol. q's)



• Sym. operators $\begin{cases} \rightarrow U_e(\alpha_e, \Sigma_2) = e^{i\alpha_e Q_e(\Sigma_2)} \\ \rightarrow U_m(\alpha_m, \Sigma_2) = e^{i\alpha_m Q_m(\Sigma_2)} \end{cases}$

with $\alpha_e \sim \alpha_e + 2\pi$ and $\alpha_m \sim \alpha_m + 2\pi$.

WILSON LINES

Crucial observation: shifting $A \rightarrow A + \xi$ with ξ a NON-QUANTIZED closed 1-form, still keeps the action invariant ($F \mapsto F$).

However, the Wilson lines are not invariant:

$$W_h(\gamma) \mapsto e^{i\beta} W_h(\gamma) \quad \text{with} \quad \beta = \int_{\gamma} \xi \in \mathbb{R} / \frac{2\pi\mathbb{Z}}{\cong U(1)}$$

\Rightarrow this is not a redundancy, but a global symmetry of the type just described, i.e. a 1-form symmetry.

\rightsquigarrow we want the associated W.l.

In order to derive the W.L., we use the usual trick: we change variable in P.I., applying a symmetry transformation with non-const. parameter. For 1-form sym, this corresponds to ξ being NON-CLOSED:

$$\delta S = -\frac{1}{e^2} \int \xi \wedge d * F \quad \text{---} \frac{-1}{e^2} \int \underbrace{d\xi \wedge * F}_{d(\xi \wedge * F) + \xi \wedge d * F}$$

$$0 = \int e^{iS} i \delta S W + \int e^{iS} \delta W = \int e^{iS} \left(-\frac{iW}{e^2} \int \xi \wedge d * F + i n \left(\int_{\gamma} \xi \right) W \right) \rightarrow \int_{\gamma} = \int \xi \wedge \delta^{d-1}(\gamma)$$

(d-1)-form s.t. $\int_{\gamma} \eta \wedge \delta^{d-1}(\gamma) = \int_{\gamma} \eta \wedge \theta_{\gamma}$

$$\Rightarrow \left\langle \frac{1}{e^2} d * F W_n(\gamma) \right\rangle = n \int \delta^{d-1}(\gamma) \langle W_n(\gamma) \rangle \quad (*)$$

Consider a topologically trivial submanif. Σ_{d-2} .

It is the boundary of a D_{d-1} .

$$\int_{D_{d-1}} \delta^{d-1}(\gamma) = \gamma \cdot D_{d-1} = \text{Link}(\gamma, \Sigma_{d-2})$$

Let's integrate (*) over D_{d-1} , remembering $J_e = \frac{1}{e^2} F$:

$$\left\langle Q_e(\Sigma_{d-2}) W_n(\gamma) \right\rangle = n \text{Link}(\Sigma_{d-2}, \gamma) \langle W_n(\gamma) \rangle$$

$$\frac{1}{e^2} \int_{\Sigma_{d-2}} * F$$

\Downarrow

This shows what is the top op. $U_e(\Sigma_{d-2})$ and that the Wilson lines are charged under this symmetry.

→ Symmetry transformation

q_E (electr. charge of Wils. lin.)



$$\ln_{d=4} \langle U_{e^{i d_E}}(\Sigma_2) e^{i n \int_{\gamma} A} \rangle = e^{i d_E n \text{Link}(\Sigma_2, \gamma)} \langle e^{i q_E \int_{\gamma} A} \rangle$$

$$U_{e^{i d_E}}(\Sigma_2) = e^{i d_E Q_E(\Sigma_2)}$$

Sym. group is $U(1)$

$d_E + 2\pi \sim d_E$ due to quant. of $q_E = n$.

$\Sigma_2 = S^2$:



Summary:

- Sym op: $U_{e^{i d_E}}(S^2) = e^{i d_E Q_E(S^2)}$ 2d topol. op.

- Charged op: $e^{i q_E \int_{\gamma} A}$ $q_E = n \in \mathbb{Z}$

- Sym. group: $e^{i d_E} \in U(1)$

↓

"ELECTRIC 1-form SYMMETRY"

't Hooft Loop $T_n(\gamma)$

- Probe magh. part. (monopole)
- Closed line \leftrightarrow gauge invariance of dual photon
- $q_M = n \in \mathbb{Z}$
- obtain same formal expression as before when we dualize electric \leftrightarrow magnetic.



"MAGNETIC 1-form SYMMETRY"

Alternatively: inserting $T_n(\gamma_{d-3})$ in a correlator modifies P.I. domain, specifying b.c.

$$\int_{S^2} \frac{F}{2\pi} = n \quad \text{with} \quad \text{Link}(S^2, \gamma_{d-3}) = 1$$

$$\langle U_m(\alpha_m, \Sigma_2) T_n(\gamma_{d-3}) \rangle = e^{i n \alpha_m \text{Link}(S^2, \gamma_{d-3})} \langle T_n(\gamma_{d-3}) \rangle$$

↑
now $\int \frac{F}{2\pi}$ is
fixed to a specific
number \Rightarrow it goes
outside P.I.

Generalisations

G p -form symmetry in d dim :

- Sym op. $U_g(\Sigma_{d-p-1})$


- Charged objects $W(q, \gamma_p)$

- Sym. transf. $\langle U_g(\Sigma_{d-p-1}) W(q, \gamma_p) \rangle = R(g)^q \langle W(q, \gamma_p) \rangle$
if linked



Take-home message : Existence of sym = Existence of **TOPOLOGICAL OPERATORS**

Adding matter

Let's remember how 1-form sym work:



If we now add charged fields $\phi(x)$ with charge q there will be gauge invariant lines that can end on the location x of the charged operator: in fact the gauge transf. on the extremum x of the WL is compensated by the gauge transf. of $\phi(x)$



To be consistent:

$$e^{iq\theta} = 1$$

i.e. $\theta = \frac{2\pi k}{q}$

$$U(1)^{(1)} \rightarrow \mathbb{Z}_q^{(1)}$$

This is called "SCREENING" by lower dim. operators.
(Now line is uncharged.)

NON-ABELIAN GAUGE THEORIES

$$S = -\frac{1}{2g^2} \int \text{Tr}(F \wedge *F) \quad \text{Gauge group } G.$$

P.L. integrates over all G -bundles & their connections modulo gauge transformations.

G -bundl :

- cover $\{U_i\}$
- transition functs $g_{ij} : U_i \cap U_j \rightarrow G$ s.t. $g_{ij} g_{jk} g_{ki} = \mathbb{1}$

Connection A :

- local 1-forms $A_i \in \Omega^1(U_i, \mathfrak{g})$ glued as

$$A_j = g_{ij}^{-1} A_i g_{ij} + i g_{ij}^{-1} dg_{ij}$$

This should not be confused with gauge transf.; these are defined as $U_i : U_i \rightarrow G$ that acts as

$$A_i \mapsto U_i A_i U_i^{-1} + i U_i dU_i^{-1} \quad g_{ij} \mapsto U_i g_{ij} U_j^{-1}$$

$$\begin{aligned} & (g_{ij}^{-1})^t (U_i A_i U_i^{-1} + i U_i dU_i^{-1}) g_{ij}^t + i g_{ij}^{-1} dg_{ij} \\ & U_j g_{ij}^{-1} U_i^{-1} \quad \downarrow \quad i U_j g_{ij}^{-1} dU_i^{-1} U_i g_{ij} U_j^{-1} \end{aligned}$$

$$\begin{aligned} & U_j (g_{ij}^{-1} A_i g_{ij} + i g_{ij}^{-1} dg_{ij}) U_j^{-1} \quad i U_j g_{ij}^{-1} U_i^{-1} (dU_i^{-1} g_{ij} U_j^{-1} + U_i dg_{ij}^{-1} U_j^{-1} + U_i g_{ij} dU_j^{-1}) \\ & + i U_j dU_j^{-1} = A_j \end{aligned}$$

Wilson lines.

Let us consider the Wilson lines

$$W_R(\gamma) = \text{tr}_R P e^{i \int_{\gamma} A}$$


- To define the path ord. exp on an arbitrary curve γ we cut γ in small arcs $\gamma_i \subset U_i$ and

use A_i to compute $\text{hol}_{\gamma_i}(A_i) = P e^{i \int_{\gamma_i} A_i}$.

- Under gauge transf. $\text{hol}_{\gamma_i} \rightarrow U_i(x_{in}) \text{hol}_{\gamma_i} U_i(x_{fin})^{-1}$

$$P e^{i \int_{\gamma} A} = \prod_{k=1, \dots} \text{hol}_{\gamma_{ik}}(A_{ik}) g_{i_k i_{k+1}}(x_{i_{k+1}})$$

and it transforms to $U_i(x_{in}) P e^{i \int_{\gamma} A} U_i(x_{fin})^{-1}$.



1-form symmetries.

Let us consider abelian case first:

- 1-form sym. $A \mapsto A + \lambda$ λ closed 1-form

• On patch U_i : $\lambda|_{U_i} = d\eta_i$

- We can make a local gauge transform. $U_i = e^{-i\eta_i}$

that corresponds to modifying the transition functions as

$$(*) \quad g_{ij} \mapsto e^{-i\eta_i} g_{ij} e^{i\eta_j} = g_{ij} t_{ij} \quad t_{ij} = e^{i(\eta_i - \eta_j)}$$

keeping A invariant.

$$\rightarrow e^{i \int_{\gamma} A} \mapsto e^{i \int_{\gamma} A} \left(e^{-i\eta_i(x_{in})} e^{i\eta_j(x_{fin})} \right) = e^{i \int_{\gamma} A}$$

In abelian theory the action of 1-form sym is (\star) .
 This can be easily generalised to non-abelian gauge theories

- 1-form sym: $g_{ij} \mapsto g_{ij} t_{ij}$
- To preserve the cocycle condition t_{ij} must commute with any possible $g_{ij} \Rightarrow t_{ij} \in Z(G)$ (\star) .

Moreover, it must happen that $t_{ij} t_{jk} t_{ki} = 1$

$[(\star) Z(G) = \{ z \in G \mid z g z^{-1} = g \ \forall g \in G \}$ is
 the CENTER of the group G . It is ABELIAN.]

- Consider the Wilson line $W_R(\gamma) = \text{tr}_R P e^{i \int_{\gamma} A}$
 with R an irrep then $t_R \in Z(G)$ is
 a matrix proportional to the identity with
 prop. factor being the phase $\phi_R(t) = \frac{\text{Tr}_R(t)}{\text{Tr}_R(1)}$

The Wilson line then transforms as

$$W_R(\gamma) \mapsto \phi_R(g) W_R(\gamma)$$

$$\text{with } g = \prod_{\substack{U_{ij} \text{ t.c.} \\ U_{ij} \cap \gamma \neq \emptyset}} t_{ij} \in Z(G)$$

\Rightarrow $W_R(\gamma)$ are line ops changed
 under the 1-form sym $Z(G)^{(1)}$.

- For $G = SU(N)$ $Z(G) = \mathbb{Z}_N$ and

$$W_R(\gamma) \mapsto e^{2\pi i q \frac{a}{N}} W_R(\gamma)$$

with $a=0, 1, \dots, N-1$ labels \mathbb{Z}_N group-elm. and $q=0, \dots, N-1$ is the N -ality of the rep.


- There exist a SURFACE OPERATOR

$U(\Sigma_2)$ that will be the generator of a \mathbb{Z}_N one-form symmetry; in fact

$\langle U(\Sigma_2) W_N(\gamma) \dots \rangle$ and $\langle W(\gamma) \dots \rangle$ turn out to differ by a factor $e^{2\pi i \frac{q}{N} \text{Link}(\Sigma_2, \gamma)}$



- Why \mathbb{Z}_N instead of a continuous sym. like in abelian gauge theories?

Let's remember how 1-form sym works:



If we now add charged fields $\phi(x)$ with charge q there will be gauge invariant lines that can end on the location x of the charged operator: in fact the gauge transf. on the extremum x of the WL is compensated by the gauge transf. of $\phi(x)$

Now,



To be consistent:

$$e^{iq\theta} = 1$$

i.e. $\theta = \frac{2\pi k}{q}$

$$U(1)^{(1)} \rightarrow \mathbb{Z}_q^{(1)}$$

Wilson lines corresponding to probes with charges $\notin q\mathbb{Z}$ cannot end on $\phi(x)$ and have in fact a non-trivial transformation under $\mathbb{Z}_q^{(1)}$

- In Maxwell theory there is no charged field, then WL for all probes have non-trivial $U(1)^{(1)}$ transformation.

- However, in YM there are ADJOINT FIELDS, i.e. the gluons gauge bosons.

Probes in the adjoint rep produces WL that can end on the location of an adjoint field; then one can unlink the Σ_2 from the line and the corresponding WL must have zero charge

Only weights $\bar{\mu}$ that are not in $\Lambda_{\text{root}}(\mathfrak{g})$ give WL transforming non-trivially $\rightsquigarrow \mathbb{Z}_N^{(1)}$ -sym.