





η^1 -alkenyl

R




M

η^1 -alkynyl




η^1 -aryl

Hydride β -elimination




Stabilizing groups:

methyl, benzyl ($\text{CH}_2\text{C}_6\text{H}_5$), neopentyl (CH_2CMe_3), trimethylsilylmethyl (CH_2SiMe_3)




C–H *agostic* interactions

Typical synthetic procedure





Alternative option: Grignard Reagents



C-C = 137 pm vs 134 pm in ethene

η^2 -alkene
***side-on* coordination**







metallacyclopropane

X-ray structure of $\text{Ru}(\eta^2\text{-C}_2\text{H}_4)(\text{PMe}_3)_4$



C–C = 144 pm vs 134 pm in ethene




π -symmetry orbital for isolated ethene, σ -symmetry orbital when ethene is coordinated η^2 (*side-on*)



Dewar – Chatt – Duncanson Model





Alkene-to-M donation
(a)






M-to-alkene back-donation
(b)

Dewar – Chatt – Duncanson Model





π -backdonation




metallacyclopropane

η^2 -alkene bond - fluxionality






4-electron donor




C—C in Co_2C_2 -unit = 136 pm vs 120 pm in C_2Ph_2




The two C_2Co planes, *i.e.* the two η^2 bonds, are roughly orthogonal


Non-conjugated Dienes




Cycloocta-1,5-diene, cod











Butadiene -
lays on the xy
plane, above the
metal










(a)






(b)








empty $(n+1)s$ and $(n+1)p_z$




dz^2
 dx^2-y^2 $e_g =$





dxz
 dyz



dxz
 dyz
 dxy $t_{2g} =$



dxy ,
 dx^2-y^2




dz^2







η^1 -mode




η^3 -mode



η^5 -mode




empty $(n+1)s$ and $(n+1)p_z$





***neo*-Menthylcyclopentadienyl**


Allyl group,
 $\text{CH}_2=\text{CH}-\text{CH}_2^-$



$\eta^1-(\text{CH}_2\text{CH}=\text{CH}_2)$









π -acceptor

3π




π -donor


2π




σ -donor

1π









syn




anti


syn-anti exchange mechanism









(b)



(c)





Cyclobutadiene is anti-aromatic (4π -electrons), and unstable as a free molecule; but its dianionic form is stable because of being aromatic (6π -electrons) - cyclobutadiene is stabilized by significant metal to ligand π -back donation to the vacant ligand orbitals.