

# Chimica Computazionale

## Hybrid multiscale schemes

Emanuele Coccia



UNIVERSITÀ  
DEGLI STUDI  
DI TRIESTE



**DSCF**


Dipartimento di  
**Scienze Chimiche  
e Farmaceutiche**

PhotoInduced Quantum Dynamics (PIQD) Group



# Hybrid multiscale methods



- Physical and chemical properties in **large** systems ( $10^4$ - $10^6$  atoms)  $\rightarrow$  quantum (QM) methods **not applicable**
- Quantum meets classical:
  - QM for a (small) subregion of the system
  - Classical for the rest of it (**environment**)





# Hybrid multiscale methods

## Biological/biochemical applications




### ENZYMATIC CATALYSIS



### ION CHANNELS



### METAL-ENHANCED SPECTROSCOPIES



# Nobel prize in Chemistry 2013



© Nobel Media AB. Photo: A. Mahmoud

**Martin Karplus**

Prize share: 1/3



© Nobel Media AB. Photo: A. Mahmoud

**Michael Levitt**

Prize share: 1/3



© Nobel Media AB. Photo: A. Mahmoud


**Arieh Warshel**

Prize share: 1/3


---

The Nobel Prize in Chemistry 2013 was awarded jointly to Martin Karplus, Michael Levitt and Arieh Warshel "for the development of multiscale models for complex chemical systems"


# Hybrid multiscale methods



# Hybrid multiscale methods




# Hybrid multiscale methods



- How the **classical part** is modelled?




# Hybrid multiscale methods





- How the **classical part** is modelled?
- How **partition** is done?

# Hybrid multiscale methods



- How the **classical part** is modelled?
- How **partition** is done?
- How QM and classical subregions **interact**?

# Modelling the classical environment

<p>The molecular cavity containing the target and the dielectric constant</p>	<p>Continuum</p> <p><math>\epsilon</math></p>  A diagram showing a target molecule (represented by a small cluster of atoms) embedded within a blue, irregularly shaped volume representing the molecular cavity. The volume is labeled with the Greek letter epsilon ( $\epsilon$ ), indicating the dielectric constant of the continuum environment.	<p>Atomistic</p>  A diagram showing a target molecule (represented by a ball-and-stick model) surrounded by a collection of small blue dots, representing the configuration of environment atoms around the target and the MM force field.	<p>The configuration of the environment atoms around the target and the MM force field</p>
---	---	--	--

- **Atomistic**: use force fields from molecular mechanics (MM)
- **Continuum**: the environment is a polarizable medium


# Partitioning the system

- Local character of most chemical reactions in condensed phases
- Distinction between a “reaction center” and a “spectator”
- Expensive but accurate QM for the small “reaction center”
- Cheaper classical methods for the “spectator” region


# Partitioning the system

- Local character of most chemical reactions in condensed phases
- Distinction between a “reaction center” and a “spectator”
- Expensive but accurate QM for the small “reaction center”
- Cheaper classical methods for the “spectator” region
- Try to avoid to cut bonds at the boundaries
- But this too is covered by the models (see next slides)

# QM/MM methods




- Three types of interaction:




- Three types of interaction:
  - Among atoms in the QM region





- Three types of interaction:
  - Among atoms in the QM region
  - Among atoms in the MM region (electrons **ignored** in the MM region!)




- Three types of interaction:
  - Among atoms in the QM region
  - Among atoms in the MM region (electrons **ignored** in the MM region!)
  - Among QM and MM atoms ←

- $E_{QM/MM}(QM/MM)$ : coupling energy between QM and MM subregions


- $E_{QM/MM}(QM/MM)$ : coupling energy between QM and MM subregions
- **Subtractive** QM/MM coupling

$$E_{QM/MM}(QM/MM) = E_{MM}(\text{full}) + E_{QM}(QM) - E_{MM}(QM)$$



- $E_{QM/MM}(QM/MM)$ : coupling energy between QM and MM subregions
- **Subtractive** QM/MM coupling


$$E_{QM/MM}(QM/MM) = E_{MM}(\text{full}) + E_{QM}(QM) - E_{MM}(QM)$$



- **Advantage:**
  - Straightforward implementation (no QM/MM communication)

- $E_{QM/MM}(QM/MM)$ : coupling energy between QM and MM subregions
- **Subtractive** QM/MM coupling

$$E_{QM/MM}(QM/MM) = E_{MM}(\text{full}) + E_{QM}(QM) - E_{MM}(QM)$$



- **Advantage:**
  - Straightforward implementation (no QM/MM communication)

## Drawbacks:

- Required force field for QM subregion
- No polarization of the QM electron density by MM environment


- Additive QM/MM coupling

$$\hat{H}_{\text{full}}(\text{full}) = \hat{H}_{\text{QM}}(\text{QM}) + \hat{H}_{\text{MM}}(\text{MM}) + \hat{H}_{\text{QM/MM}}(\text{QM/MM})$$

- MM region only at classical level
- $E_{\text{QM/MM}}(\text{QM/MM})$  explicitly computed
  - Mechanical embedding
  - Electrostatic embedding

$$\begin{aligned} E_{\text{QM/MM}} &= E_b + E_{nb} \\ E_b &= E_{\text{bond}} + E_{\text{angle}} + E_{\text{dihe}} \\ E_{nb} &= E_{\text{VdW}} + E_{\text{el}} \end{aligned}$$

## Mechanical embedding





$$\hat{H}_{QM/MM} = \hat{H}_b + \hat{H}_{nb}$$

$$\begin{aligned}
 \hat{H}_{QM/MM} &= \hat{H}_b + \hat{H}_{nb} \\
 \hat{H}_b &= \sum_k^{QM \text{ atoms}} \sum_m^{MM \text{ atoms}} E_{bond}(r_{km}) \\
 &+ \sum_{k,m,l}^{at \text{ least one QM}} E_{angle}(\theta_{kml}) \\
 &+ \sum_{k,m,l,p}^{at \text{ least one QM}} E_{dihe}(\theta_{kmlp})
 \end{aligned}$$

$$\begin{aligned}
 \hat{H}_{QM/MM} &= \hat{H}_b + \hat{H}_{nb} \\
 \hat{H}_b &= \sum_k^{QM \text{ atoms}} \sum_m^{MM \text{ atoms}} E_{bond}(r_{km}) \\
 &+ \sum_{k,m,l}^{at \text{ least one QM}} E_{angle}(\theta_{kml}) \\
 &+ \sum_{k,m,l,p}^{at \text{ least one QM}} E_{dihe}(\theta_{kmlp}) \\
 \hat{H}_{nb} &= \sum_k^{QM \text{ atoms}} \sum_m^{MM \text{ atoms}} \left[ \frac{Z_k q_m}{r_{km}} + 4\epsilon_{km} \left( \frac{\sigma_{km}^{12}}{r_{km}^{12}} - \frac{\sigma_{km}^6}{r_{km}^6} \right) \right]
 \end{aligned}$$

- Electrostatic embedding: polarization **included**

$$\hat{H}_{nb,el} = \hat{H}_{nb} - \sum_i^{\text{electrons}} \sum_m^{\text{MM atoms}} \frac{q_m}{r_{im}}$$

- Electrostatic embedding: polarization **included**

$$\hat{H}_{nb,el} = \hat{H}_{nb} - \sum_i^{\text{electrons}} \sum_m^{\text{MM atoms}} \frac{q_m}{r_{im}}$$

- Electrons see MM atoms as special nuclei with non-integer and possibly negative charges

- Electrostatic embedding: polarization **included**


$$\hat{H}_{nb,el} = \hat{H}_{nb} - \sum_i^{\text{electrons}} \sum_m^{\text{MM atoms}} \frac{q_m}{r_{im}}$$

- Electrons see MM atoms as special nuclei with non-integer and possibly negative charges
- Risk of **overpolarization** at boundaries (electron spill-out)
- Smeared charges


$$\Omega_m(r) = \sqrt{\frac{q_m}{\pi\alpha^3}} \exp\left[\frac{(r - r_m)^2}{2\alpha^2}\right]$$

# QM/MM

- **Capping bonds** at the QM/MM boundary
- Monovalent link atom at an appropriate position along the bond vector between the QM and MM atoms



- Link atom **only** present in the QM calculation
- Link atom **frozen** at a given position
- Do not cut double or triple bonds



$$\begin{aligned}
\langle \Psi | \hat{H}_{\text{full}} | \Psi \rangle &= \langle \Psi | \hat{H}_{\text{QM}} | \Psi \rangle + \langle \Psi | \hat{H}_{\text{MM}} | \Psi \rangle + \langle \Psi | \hat{H}_{\text{QM/MM}} | \Psi \rangle \\
&= \langle \Psi | - \sum_i^N \frac{1}{2} \nabla_i^2 - \sum_i^N \sum_k^K \frac{Z_k}{r_{ik}} + \sum_{i<j} \frac{1}{r_{ij}} + \sum_{k<l} \frac{Z_k Z_l}{r_{kl}} | \Psi \rangle \\
&+ \langle \Psi | \Psi \rangle E_{\text{MM}} + \langle \Psi | - \sum_i^N \sum_m^M \frac{q_m}{r_{im}} | \Psi \rangle + \langle \Psi | \Psi \rangle (E_b + E_{nb})
\end{aligned}$$




$$\begin{aligned}
\langle \Psi | \hat{H}_{\text{full}} | \Psi \rangle &= \langle \Psi | \hat{H}_{\text{QM}} | \Psi \rangle + \langle \Psi | \hat{H}_{\text{MM}} | \Psi \rangle + \langle \Psi | \hat{H}_{\text{QM/MM}} | \Psi \rangle \\
&= \langle \Psi | - \sum_i^N \frac{1}{2} \nabla_i^2 - \sum_i^N \sum_k^K \frac{Z_k}{r_{ik}} + \sum_{i<j} \frac{1}{r_{ij}} + \sum_{k<l} \frac{Z_k Z_l}{r_{kl}} | \Psi \rangle \\
&\quad + \langle \Psi | \Psi \rangle E_{\text{MM}} + \langle \Psi | - \sum_i^N \sum_m^M \frac{q_m}{r_{im}} | \Psi \rangle + \langle \Psi | \Psi \rangle (E_b + E_{nb}) \\
&= \langle \Psi | - \sum_i^N \frac{1}{2} \nabla_i^2 - \sum_i^N \sum_k^K \frac{Z_k}{r_{ik}} - \sum_i^N \sum_m^M \frac{q_m}{r_{im}} + \sum_{i<j} \frac{1}{r_{ij}} + \sum_{k<l} \frac{Z_k Z_l}{r_{kl}} | \Psi \rangle \\
&\quad + E_{\text{MM}} + E_b + E_{nb}
\end{aligned}$$

SCF calculation **affected** by the MM charges

# Polarizable continuum models

# QM/continuum

- Continuum models for solutions
- QM solute(s), solvent as a polarizable medium



- Medium with a dielectric constant  $\epsilon$

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:
  - how **size** and **shape** of the hole are defined

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:
  - how **size** and **shape** of the hole are defined
  - how the cavity **dispersion term** is computed

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:
  - how **size** and **shape** of the hole are defined
  - how the cavity **dispersion term** is computed
  - how the **charge distribution** of the solute is described



$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:
  - how **size** and **shape** of the hole are defined
  - how the cavity **dispersion term** is computed
  - how the **charge distribution** of the solute is described
  - how the **dielectric medium** is given

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:
  - how **size** and **shape** of the hole are defined
  - how the cavity **dispersion term** is computed
  - how the **charge distribution** of the solute is described
  - how the **dielectric medium** is given
- $\epsilon$  fully describes the solvent
- Also spatial and frequency dependencies are used, i.e.  $\epsilon(\mathbf{r})$  and  $\epsilon(\omega)$

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:
  - how **size** and **shape** of the hole are defined
  - how the cavity **dispersion term** is computed
  - how the **charge distribution** of the solute is described
  - how the **dielectric medium** is given
- $\epsilon$  fully describes the solvent
- Also spatial and frequency dependencies are used, i.e.  $\epsilon(\mathbf{r})$  and  $\epsilon(\omega)$
- Sphere or ellipsoid allow for an analytical electrostatic interaction
- Interlocking spheres on each nucleus (**vdW surface**)

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{dispersion}} + \Delta G_{\text{elec}}$$

- Models differ:
  - how **size** and **shape** of the hole are defined
  - how the cavity **dispersion term** is computed
  - how the **charge distribution** of the solute is described
  - how the **dielectric medium** is given
- $\epsilon$  fully describes the solvent
- Also spatial and frequency dependencies are used, i.e.  $\epsilon(\mathbf{r})$  and  $\epsilon(\omega)$
- Sphere or ellipsoid allow for an analytical electrostatic interaction
- Interlocking spheres on each nucleus (**vdW surface**)
- Born** model

$$\Delta G_{\text{solv}} = -\frac{1}{2} \left( 1 - \frac{1}{\epsilon} \right) \frac{q^2}{r}$$

- Self-consistent reaction field (SCRF)
- Poisson equation (electrostatic potential  $\phi$ , solute charge distribution  $\rho$ )

$$\nabla \cdot (\epsilon(\mathbf{r}) \nabla \phi(\mathbf{r})) = -4\pi \rho(\mathbf{r})$$

- Self-consistent reaction field (SCRF)
- Poisson equation (electrostatic potential  $\phi$ , solute charge distribution  $\rho$ )

$$\begin{aligned}\nabla \cdot (\epsilon(\mathbf{r}) \nabla \phi(\mathbf{r})) &= -4\pi \rho(\mathbf{r}) \\ \nabla^2 \phi(\mathbf{r}) &= -\frac{4\pi}{\epsilon} \rho(\mathbf{r})\end{aligned}$$

- Self-consistent reaction field (SCRF)
- Poisson equation (electrostatic potential  $\phi$ , solute charge distribution  $\rho$ )

$$\begin{aligned}\nabla \cdot (\epsilon(\mathbf{r}) \nabla \phi(\mathbf{r})) &= -4\pi \rho(\mathbf{r}) \\ \nabla^2 \phi(\mathbf{r}) &= -\frac{4\pi}{\epsilon} \rho(\mathbf{r}) \\ 4\pi \epsilon \sigma(\mathbf{r}_s) &= (\epsilon - 1) \mathbf{F}(\mathbf{r}_s)\end{aligned}$$

- Self-consistent reaction field (SCRF)
- Poisson equation (electrostatic potential  $\phi$ , solute charge distribution  $\rho$ )

$$\begin{aligned}\nabla \cdot (\epsilon(\mathbf{r}) \nabla \phi(\mathbf{r})) &= -4\pi \rho(\mathbf{r}) \\ \nabla^2 \phi(\mathbf{r}) &= -\frac{4\pi}{\epsilon} \rho(\mathbf{r}) \\ 4\pi \epsilon \sigma(\mathbf{r}_s) &= (\epsilon - 1) \mathbf{F}(\mathbf{r}_s)\end{aligned}$$

- Extra term in the Hamiltonian

$$\begin{aligned}\hat{H}_{\text{SCRF}} &= \hat{H} + \phi_\sigma \\ \phi_\sigma(\mathbf{r}) &= \int \frac{\sigma(\mathbf{r}_s)}{|\mathbf{r} - \mathbf{r}_s|} d\mathbf{r}_s\end{aligned}$$



- Mixed solvent models
- First solvation shell explicitly modelled

- Mixed solvent models
- First solvation shell explicitly modelled
- Configurations sampling issue


- Mixed solvent models
- **First solvation shell** explicitly modelled
- Configurations sampling issue
- Parametrization of the continuum model against **experimental data**
- Mixed models may yield substantially better results than pure continuum models, at the price of an **increased** computational cost

- Mixed solvent models
- **First solvation shell** explicitly modelled
- Configurations sampling issue
- Parametrization of the continuum model against **experimental data**
- Mixed models may yield substantially better results than pure continuum models, at the price of an **increased** computational cost
- Solvation energy from few (neutral solute) to hundreds of kcal/mol (ions) in water

- Mixed solvent models
- **First solvation shell** explicitly modelled
- Configurations sampling issue
- Parametrization of the continuum model against **experimental data**
- Mixed models may yield substantially better results than pure continuum models, at the price of an **increased** computational cost
- Solvation energy from few (neutral solute) to hundreds of kcal/mol (ions) in water
- Inclusion of solvent effects may change the geometry, charge distribution and conformational preferences


# Example: Reactivity of charged species

- $S_N2$  substitution



- Kinetics

- $k = 1.3 \times 10^{10} \text{ M}^{-1}\text{s}^{-1}$  (gas phase)
- $k = 3.3 \text{ M}^{-1}\text{s}^{-1}$  (acetone)
- $k = 5.0 \times 10^{-6} \text{ M}^{-1}\text{s}^{-1}$  (water)



# Example: Reactivity of charged species


- DFT calculations ( $\omega$ B97X-D functional, 6-31+G(d) basis set)
- SMD model for solvation (screening model based on density)

Species	$\Delta G_{(g)}$	$\Delta G_{(solution)}$
$Cl^- + CH_3Br$	0.0	0.0
$Cl^- \bullet CH_3Br$	-48.8	10.9
TS	13.3	100.7
$ClCH_3 \bullet Br^-$	-34.9	38.5
$ClCH_3 + Br^-$	-18.7	31.0

- Computed kinetics
  - $k = 2.85 \times 10^{10} \text{ M}^{-1}\text{s}^{-1}$  (gas phase, 2.3 times larger than exp)
  - $k = 1.4 \times 10^{-5} \text{ M}^{-1}\text{s}^{-1}$  (acetone,  $2.4 \times 10^{-5}$  smaller)
- Inaccuracy due to approximations in SMD

# Example: acid-base equilibria


- Dissociation of a **weak acid** HA in solvent (water)





# Example: acid-base equilibria

- Dissociation of a **weak acid** HA in solvent (water)



$$K_A = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}][\text{H}_2\text{O}]}$$

Thermodynamic cycle

$$\begin{aligned}\Delta G_{(aq)}^0 &= \Delta G_{(g)}^0 + \Delta G_{\text{solv}}(\text{A}^-) + \Delta G_{\text{solv}}(\text{H}_3\text{O}^+) \\ &\quad - \{ \Delta G_{\text{solv}}(\text{HA}) + \Delta G_{\text{solv}}(\text{H}_2\text{O}) \}\end{aligned}$$

$$\text{p}K_A = -\log_{10} K_A = \frac{-\Delta G_{(aq)}^0}{RT \ln 10}$$

