

# DFT in a Nutshell

Kieron Burke<sup>[a,b]</sup> and Lucas O. Wagner<sup>\*[a,b]</sup>

The purpose of this short essay is to introduce students and other newcomers to the basic ideas and uses of modern electronic density functional theory, including what kinds of approximations are in current use, and how well they work (or not). The complete newcomer should find it orients them well, while even longtime users and aficionados might find


something new outside their area. Important questions varying in difficulty and effort are posed in the text, and are answered in the Supporting Information. © 2012 Wiley Periodicals, Inc.

DOI: 10.1002/qua.24259

## Electronic Structure Problem

For the present purposes, we define the modern electronic structure problem as finding the ground-state energy of nonrelativistic electrons for arbitrary positions of nuclei within the Born-Oppenheimer approximation.<sup>[1]</sup> If this can be done sufficiently accurately and rapidly on a modern computer, many properties can be predicted, such as bond energies and bond lengths of molecules, and lattice structures and parameters of solids.

Consider a diatomic molecule, whose binding energy curve is illustrated in Figure 1. The binding energy is given by



**Figure 1.** Generic binding energy curve. For  $N_2$ , values for  $R_0$  and  $D_e$  are given in Table 1. [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]

$$E_{\text{bind}}(R) = E_0(R) + \frac{Z_A Z_B}{R} - E_A - E_B \quad (1)$$

where  $E_0(R)$  is the ground-state energy of the electrons with nuclei separated by  $R$ , and  $E_A$  and  $Z_A$  are the atomic energy and charge of atom  $A$  and similarly for  $B$ . The minimum tells us the bond length ( $R_0$ ) and the well-depth ( $D_e$ ), corrected by zero-point energy ( $\hbar\omega/2$ ), gives us the dissociation energy ( $D_0$ ).

The Hamiltonian for the  $N$  electrons is

$$\hat{H} = \hat{T} + \hat{V}_{\text{ee}} + \hat{V}, \quad (2)$$

where the kinetic energy operator is

$$\hat{T} = -\frac{1}{2} \sum_{j=1}^N \nabla_j^2, \quad (3)$$

the electron–electron repulsion operator is

$$\hat{V}_{\text{ee}} = \frac{1}{2} \sum_{i \neq j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}, \quad (4)$$

and the one-body operator is

$$\hat{V} = \sum_{j=1}^N v(\mathbf{r}_j). \quad (5)$$

For instance, in a diatomic molecule,  $v(\mathbf{r}) = -Z_A/r - Z_B/|\mathbf{r} - \mathbf{R}|$ . We use atomic units unless otherwise stated, setting  $e^2 = \hbar = m_e = 1$ , so energies are in Hartrees (1 Ha = 27.2 eV or 628 kcal/mol) and distances in Bohr radii (1  $a_0$  = 0.529 Å). The ground-state energy satisfies the variational principle:

$$E = \min_{\Psi} \langle \Psi | \hat{H} | \Psi \rangle, \quad (6)$$

where the minimization is over all antisymmetric  $N$ -particle wavefunctions. This  $E$  was called  $E_0(R)$  in Eq. (1).\*

Many traditional approaches to solving this difficult many-body problem begin with the Hartree–Fock (HF) approximation, in which  $\Psi$  is approximated by a single Slater determinant (an antisymmetrized product) of orbitals (single-particle wavefunctions)<sup>[2]</sup> and the energy is minimized.<sup>[3]</sup> These include configuration interaction, coupled cluster, and Møller–Plesset perturbation theory, and are mostly used for finite systems, such as molecules in the gas phase.<sup>[4]</sup> Other approaches use reduced descriptions, such as the density matrix or Green's function, but leading to an infinite set of coupled equations that must somehow be truncated, and these are more common in applications to solids.<sup>[5]</sup>

More accurate methods usually require more sophisticated calculation, which takes longer on a computer. Thus, there is a compelling need to solve ground-state electronic structure problems reasonably accurately, but with a cost in

[a, b] K. Burke, L. O. Wagner

Department of Chemistry, University of California, Irvine, California 92697

Department of Physics, University of California, Irvine, California 92697

E-mail: lwagner@uci.edu

\*Explain why a vibrational frequency is a property of the ground-state of the electrons in a molecule.

© 2012 Wiley Periodicals, Inc.

**Kieron Burke** is a professor of heretical physical and computational chemistry at the University of California in Irvine. He spent the last two decades of his life in functionals. [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]



**Lucas Wagner** began working in 2009 as a graduate student with Kieron Burke starting off curious, plucky, and impressionable in the wide world of DFT. Lucas studies simple (mostly 1d) models and enjoys the combination of analytical and numerical studies involved. [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]



computer time that does not become prohibitive as the number of atoms (and therefore electrons) becomes large.

## Basic DFT

The electronic density  $n(\mathbf{r})$  is defined by the requirement that  $n(\mathbf{r}) d^3r$  is the probability of finding any electron in the volume  $d^3r$  around  $\mathbf{r}$ . For a single electron with wavefunction  $\phi(\mathbf{r})$ , it is simply  $|\phi(\mathbf{r})|^2$ . In density functional theory (DFT), we write the ground-state energy in terms of  $n(\mathbf{r})$  instead of  $\Psi$ . The first DFT was formulated by Thomas<sup>[6]</sup> and Fermi.<sup>[7]</sup> The kinetic energy density at any point is approximated by that of a uniform electron gas of noninteracting electrons of density  $n(\mathbf{r})$ , which for a spin-unpolarized system is:<sup>†</sup>

$$T^{\text{TF}} = a_5 \int d^3r n^{5/3}(\mathbf{r}), \quad a_5 = 3(3\pi^2)^{2/3}/10. \quad (7)$$

The interelectron repulsion is approximated by the classical electrostatic self-energy of the charge density, called the Hartree energy:

$$U = \frac{1}{2} \int d^3r \int d^3r' \frac{n(\mathbf{r}) n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}. \quad (8)$$

Because the one-body potential couples only to the density,

$$V = \langle \hat{V} \rangle = \int d^3r n(\mathbf{r}) v(\mathbf{r}). \quad (9)$$

The sum of these three energies is then minimized, subject to the physical constraints:

$$n(\mathbf{r}) \geq 0, \quad \int d^3r n(\mathbf{r}) = N. \quad (10)$$

This absurdly crude theory gives roughly correct energies (errors about 10% for many systems) but is not nearly good

enough for most properties of interest (for example, molecules do not bind<sup>[8]</sup>). For same-spin, noninteracting fermions in 1d, the corresponding kinetic energy is

$$T^{\text{TF 1d}}[n] = \frac{\pi^2}{6} \int dx n^3(x) \quad (11)$$

and makes only a 25% error on the density of a single particle in a box. Hours of endless fun and many good and bad properties of functional approximations can be understood by applying Eq. (11) to standard text book problems in quantum mechanics,<sup>‡</sup> and noting what happens, especially for more than one particle.<sup>§</sup>

But modern DFT began with the proof that the solution of the many-body problem can be found, in principle, from a density functional. To see this, we break the minimization of Eq. (6) into two steps. First minimize over all wavefunctions yielding a certain density, and then minimize over all densities. Because the one-body potential energy depends only on the density, we can define separately<sup>[9,10]</sup>

$$F[n] = \min_{\Psi \rightarrow n} \langle \Psi | \{ \hat{T} + \hat{V}_{\text{ee}} \} | \Psi \rangle, \quad (12)$$

where the minimization is over all antisymmetric wavefunctions yielding a given density  $n(\mathbf{r})$ .<sup>¶</sup> This is transparently a *functional* of the density, meaning it assigns a number to each density, as was first proven by Hohenberg and Kohn.<sup>\*\*[11]</sup> Then

<sup>†</sup>Calculate the TF kinetic energy for a 1d particle of mass  $m = 1$  in (a) a harmonic well ( $v(x) = x^2/2$ ) and (b) in a delta-well ( $v(x) = -\delta(x)$ ). Give the % errors.

<sup>§</sup>Deduce the exact energy for  $N$  same-spin fermions in a flat box of width 1 bohr. Then evaluate the local approximation to the kinetic energy for  $N = 1, 2$ , and 3, and calculate the % error.

<sup>¶</sup>Derive  $F[n]$  for a single electron. It has no electron–electron interaction, and is known as the von Weizsäcker kinetic energy.

<sup>\*\*</sup>Technically, HK only proved fact this for  $v$ -representable densities, i.e. densities which are the ground-state of some external one-body potential.

<sup>†</sup>Evaluate the TF kinetic energy of the H atom and deduce the % error. Repeat using spin-DFT.

$$E = \min_n \left\{ F[n] + \int d^3r n(\mathbf{r}) v(\mathbf{r}) \right\}, \quad (13)$$

where the minimization is over all reasonable densities satisfying Eq. (10). Hohenberg and Kohn proved (i) that all properties are determined by  $n(\mathbf{r})$ , i.e., they are functionals of  $n(\mathbf{r})$ ,<sup>††</sup> (ii)  $F[n]$  is a universal functional, independent of  $v(\mathbf{r})$ , and (iii) the exact density satisfies

$$\frac{\delta F}{\delta n(\mathbf{r})} = -v(\mathbf{r}), \quad (14)$$

where  $\delta F/\delta n(\mathbf{r})$  is the functional derivative of  $F$  with respect to  $n(\mathbf{r})$ . In fact, these days we use *spin* DFT,<sup>[12]</sup> in which all quantities are considered functionals of the up,  $n_\uparrow(\mathbf{r})$ , and down,  $n_\downarrow(\mathbf{r})$ , spin densities separately. This makes approximations more accurate for odd electron systems and allows treatment of collinear magnetic fields. All functionals written without spin dependence, such as the ones discussed thus far, are assumed to be referring to a spin-unpolarized system.

The next crucial step in developing the modern theory came from (re)-introducing orbitals. Kohn and Sham<sup>[13]</sup> vastly improved the accuracy of DFT by imagining a fictitious set of noninteracting electrons that are defined to have the same density as the interacting problem. They are still spin- $\frac{1}{2}$  fermions obeying the Pauli principle, so like in HF theory, their wavefunction is (usually) a Slater determinant, an antisymmetrized product of orbitals of each spin,  $\phi_{j\sigma}(\mathbf{r})$ ,  $j = 1, \dots, N_{\sigma}$ ,  $\sigma = \uparrow, \downarrow$ . These KS electrons satisfy a noninteracting Schrödinger equation:

$$\left\{ -\frac{1}{2} \nabla^2 + v_{S,\sigma}(\mathbf{r}) \right\} \phi_{j\sigma}(\mathbf{r}) = \epsilon_{j\sigma} \phi_{j\sigma}(\mathbf{r}), \quad (15)$$

and the  $\epsilon_{j\sigma}$  are called KS eigenvalues and  $\phi_{j\sigma}(\mathbf{r})$  are KS orbitals.<sup>‡‡</sup> By evaluating Eq. (3) on the KS Slater determinant, the KS kinetic energy is the sum of the orbital contributions:<sup>§§</sup>

$$T_S = -\frac{1}{2} \sum_{\sigma=\uparrow,\downarrow} \sum_{j=1}^{N_\sigma} \int d^3r \phi_{j\sigma}^*(\mathbf{r}) \nabla^2 \phi_{j\sigma}(\mathbf{r}). \quad (16)$$

If we write the energy in terms of KS quantities:

$$E = T_S + U + V + E_{XC}, \quad (17)$$

$E_{XC}$  is defined by Eq. (17) and called the exchange-correlation (XC) energy.<sup>¶¶</sup> KS showed that one could extract the unknown KS potential if one only knew how the terms depend on the density. Writing the Hartree potential as

$$v_H(\mathbf{r}) = \frac{\delta U}{\delta n(\mathbf{r})} = \int d^3r' \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}, \quad (18)$$

then

$$v_{S,\sigma}(\mathbf{r}) = v_\sigma(\mathbf{r}) + v_H(\mathbf{r}) + v_{XC,\sigma}(\mathbf{r}), \quad (19)$$

where

$$v_{XC,\sigma}(\mathbf{r}) = \frac{\delta E_{XC}}{\delta n_\sigma(\mathbf{r})}. \quad (20)$$

This is a formally exact scheme for finding the ground-state energy and density for any electronic problem.<sup>\*\*\*</sup> In Figure 2,




Figure 2. Exact KS potential of He atom along  $z$  axis; for neutral atoms, the KS potential goes asymptotically to  $1/r$  for large  $r$  (data from Umrigar and Gonze, *Phys. Rev. A* 1994, 50, 3827, © American Physical Society, reproduced by permission). [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]

we emphasize the exactness of the KS scheme by plotting the exact KS potential for a He atom (which is trivial to find, once the exact density is known from an accurate many-body calculation<sup>[14]</sup>).<sup>†††</sup> Two noninteracting electrons, doubly occupying the 1s orbital of this potential, have a density that matches that of the interacting system exactly.<sup>‡‡‡</sup> But in practical calculations, we always use approximations to  $E_{XC}$  and hence to  $v_{XC}(\mathbf{r})$ .

Traditionally,  $E_{XC}$  is broken up into exchange (X) and correlation contributions:<sup>§§§</sup>

$$E_{XC} = E_X + E_C. \quad (21)$$

The exchange energy is  $\hat{V}_{ee}$  evaluated on the KS Slater determinant minus the Hartree energy, and typically dominates.<sup>¶¶¶</sup> In terms of the orbitals:

\*\*\*Why is a KS calculation much faster than direct solution of the Schrödinger equation?

†††Write the formula that extracts  $v_S(\mathbf{r})$  from  $n(\mathbf{r})$  for the helium atom. Can you explain why this does *not* tell us the vital  $v_{XC}[n](\mathbf{r})$  for any spin-unpolarized two-electron systems?

‡‡‡Why is the helium KS potential less deep than the original potential,  $-2/r$ ?

§§§What is  $E_{XC}$  for a one-electron system?

¶¶¶What is the expectation value of the Hamiltonian ( $\hat{H} = \hat{T} + \hat{V} + \hat{V}_{ee}$ ) evaluated on the KS Slater determinant? Use this to prove that the DFT definition of correlation energy is never positive.

††Again, HK proved this only for  $v$ -representable densities.

‡‡Is the sum of the KS eigenvalues equal to the total energy?

§§Apply Eq. (12) to the KS system to define  $T_S$  without ever mentioning  $v_S(\mathbf{r})$ . Then prove that  $T \geq T_S$  always.

¶¶Give the signs of  $E$ ,  $T$ ,  $V_{ee}$ ,  $U$ ,  $E_X$ , and  $E_C$  for real systems (i.e., atoms, molecules, and solids).

$$E_X = -\frac{1}{2} \sum_{\sigma, i, j} \int d^3r \int d^3r' \frac{\phi_{i\sigma}^*(\mathbf{r}) \phi_{j\sigma}^*(\mathbf{r}') \phi_{i\sigma}(\mathbf{r}') \phi_{j\sigma}(\mathbf{r})}{|\mathbf{r} - \mathbf{r}'|}. \quad (22)$$

This is precisely the same orbital expression given in HF, though it makes use of the KS orbitals (which are implicit functionals of the density<sup>[15]</sup> rather than the HF orbitals.<sup>\*\*\*\*</sup> Then the correlation energy is everything else, i.e., defined to make Eq. (17) exact.<sup>+++</sup>

## Real Calculations

Practical calculations use some simple approximation to  $E_{XC}[n_\uparrow, n_\downarrow]$ . The KS equations are started with some initial guess for the density, yielding a KS potential via Eq. (20). The KS equations are then solved and a new density is found. This cycle is repeated until changes become negligible, i.e., this a self-consistent field calculation.

The standard approximations are very simple. The local (spin) density approximation (often just called LDA) is<sup>[13]</sup>:

$$E_{XC}^{LDA} = \int d^3r e_{XC}^{unif}(n_\uparrow(\mathbf{r}), n_\downarrow(\mathbf{r})) \quad (23)$$

where  $e_{XC}^{unif}(n_\uparrow, n_\downarrow)$  is the XC energy density of a uniform gas of spin densities  $n_\uparrow$  and  $n_\downarrow$ . The X contribution was first written by Dirac,<sup>[16]</sup> and is found by inserting plane waves (the KS orbitals of a uniform gas) into Eq. (22).<sup>\*\*\*\*</sup> In the unpolarized case, the result is<sup>ssss</sup>

$$e_X^{unif}(n) = -a_X n^{4/3}, \quad a_X = 3(3\pi^2)^{1/3}/(4\pi) \quad (24)$$

while  $e_C^{unif}(n_\uparrow, n_\downarrow)$  has been calculated and accurately parametrized.<sup>[17]</sup> More accurate energetics are usually obtained with a generalized gradient approximation (GGA), which includes dependence on the gradient of the density

$$E_{XC}^{GGA} = \int d^3r e_{XC}^{GGA}(n_\uparrow(\mathbf{r}), n_\downarrow(\mathbf{r}), |\nabla n_\uparrow(\mathbf{r})|, |\nabla n_\downarrow(\mathbf{r})|), \quad (25)$$

where now the energy density is given by some approximate form. The third standard form of approximation is a hybrid, which mixes in a fixed fraction of the exact exchange energy

$$E_{XC}^{hyb} = a (E_X - E_X^{GGA}) + E_X^{GGA}, \quad (26)$$

where  $a$  is a universal parameter (often about 1/4) and  $E_X$  is the exact exchange defined in Eq. (22).

\*\*\*\*Derive a formula for  $E_X[n_\uparrow, n_\downarrow]$  in terms of  $E_X[n]$ , evaluated on various densities. Does the same formula apply to the KS kinetic energy, but with  $E_X$  replaced by  $T_S$ ? How about for  $E_C$ ?

+++This definition differs slightly from that of quantum chemistry.

\*\*\*\*Show that if electrons repelled via a contact repulsion,  $\delta(\mathbf{r} - \mathbf{r}')$ , the exchange is given exactly by LDA, and give its expression, including spin-dependence.

ssssUse dimensional analysis to explain the powers of the density in the local approximations of the kinetic energy in 1d and 3d, as well as the exchange energy in 3d.

Although LDA is uniquely defined, there are two basic flavors of the more sophisticated functionals. There are those that are derived without fitting to reference data on atoms and molecules, using information from only the slowly varying electron and known exact conditions on the functional. The standard GGA of this type is PBE,<sup>[18]</sup> while the hybrid is PBE0, mixing 25% exact exchange.<sup>[19]</sup> Of the empirical type, B88 is the standard GGA for exchange,<sup>[20]</sup> LYP for correlation,<sup>[21]</sup> combining to form BLYP. The most commonly used functional today is a hybrid called B3LYP.<sup>[22]</sup>

The original LDA became a standard tool in solid-state physics, yielding excellent lattice parameters, and fairly good bulk moduli. But LDA typically overbinds by about 1 eV/bond, which is too large an error to be useful in quantum chemistry. GGAs reduce this error to about 0.3 eV/bond, and hybrids reduce it another factor of 2. Table 1 shows typical results. Em-

**Table 1.** Differences in calculated [using Turbomole (<http://www.turbomole.com/>)<sup>[23]</sup>] versus experimental values (from Ref. [24]) for well depth ( $D_e$ ) in kcal/mol and equilibrium distance ( $R_0$ ) in pm of  $N_2$ .

	$\Delta = \text{calc} - \text{expt}$						Expt.
	HF	LDA	PBE	BLYP	PBE0	B3LYP	
$D_e$	-110	40.4	16.2	13.0	-1.73	1.06	227.0
$R_0$	-3.2	-0.33	0.41	0.40	-0.93	-0.80	109.8

pirical parametrized functionals are usually about a factor of 2 better than nonempirical ones, but with less systematic errors.<sup>[25]</sup> Although hybrids are popular in chemistry, where HF codes have been including exact exchange for decades, they are much less popular for solids, due to the singularities in HF for metals.

In contrast, many limitations of these functionals have been identified. Perhaps the most well-known is the gap problem. The fundamental gap is  $I - A$ , the difference between the ionization potential and the electron affinity of a system. This is usually larger than the KS gap, the difference between the KS HOMO and LUMO energy eigenvalues. Calculations with LDA and GGA yield fundamental gaps of insulating solids close to the KS gap, and so are too small (by about a factor of 2). This is related to self-interaction error (that is, the functionals are incorrect for one electron). Hybrid functionals often do better. Other well known failures of these approximations include the lack of asymptotic van der Waals forces, which can be important for soft matter.<sup>[26]</sup>

There are many suggestions on how to improve these standards. Presently, meta-GGAs, which use KS kinetic energy densities to approximate  $E_{XC}$ , can produce accurate energetics without using HF exchange.<sup>[27]</sup> Alternatively, including only the short-ranged contribution to exchange can also avoid the difficulties of applying HF to solids.<sup>[28]</sup> But so far, none of these is even close to replacing the standard functionals in amount of use.

Computer codes also need some basis set to represent the KS orbitals. Basis functions localized on atoms are used for molecular systems in quantum chemistry. Most codes now use Gaussians, but some use Slater-type orbitals (exponentials



Table 2. Acronyms.

Abbr.	Expansion
B88	Becke 88 $E_x$ functional
BLYP	GGA using B88 and LYP
B3LYP	BLYP hybrid
DFT	Density functional theory
GGA	Generalized gradient approximation
HOMO	Highest-occupied molecular orbital
KS	Kohn-Sham
LDA	Local density approximation
LUMO	Lowest-unoccupied molecular orbital
LYP	Lee-Yang-Parr $E_c$ functional <sup>[21]</sup>
PBE	Perdew-Burke-Ernzerhof GGA functional
PBE0	PBE hybrid
TF	Thomas-Fermi
X	Exchange
XC	Exchange-correlation

centered on atoms) and localized basis sets. In solids and materials, periodic boundary conditions are convenient for extended systems, making plane waves a natural choice. Core electrons are often treated as frozen or by pseudo-potentials, saving calculational effort. A few codes use numerical grids.<sup>[5]</sup>

Electronic KS-DFT has been implemented in a huge variety of codes, and is being applied in many areas of science. Modern computers allow routine calculation of hundreds of atoms, far more than with traditional quantum chemical methods, but with a reasonable (not very high) degree of accuracy and reliability. In traditional areas of condensed-matter physics and quantum chemistry, many experimental groups run codes themselves, and many purely computational groups exist. For example, new catalysts have been found by calculation, then demonstrated in the lab, patented, and are being further developed for industrial production.<sup>[29]</sup> But such calculations and calculators are also appearing in materials science, geology, and biochemical departments. Recent uses also include protein science, soil science, and astrophysics. Liquids can be simulated using *ab initio* molecular dynamics (AIMD).<sup>[30]</sup> There is increasing interest in matter under high pressure and temperature (but not high enough to be pure plasma) and here DFT simulation is making a large impact.<sup>[31]</sup>

In its original form, electronic DFT focused purely on the ground-state energy and density. Several approaches exist to tackle excited states, but the most popular is time-dependent density functional theory (TDDFT), based on a formal proof for time-dependent one-body potentials.<sup>[32]</sup> In the linear response regime, this leads to a simple scheme for calculating transition frequencies which has become popular for molecules. The response scheme uses the ground-state KS orbitals as its starting point. In Figure 2, the (doubly occupied)  $1s$  orbital is at  $-24.57$  eV, precisely  $-I$ , the ionization potential for the system. The  $2p$  orbital is at  $-3.429$  eV, so the first allowed optical transition of the KS electrons has frequency  $21.14$  eV.<sup>[33]</sup> TDDFT corrects these transitions to the exact value  $20.97$  eV,<sup>[34]</sup> but once again requires knowledge of an unknown functional, called the XC kernel, which depends on the perturbing frequency.<sup>[35]</sup> Present applications all use the adiabatic approximation, which ignores the frequency dependence and allows

Table 3. Notation of representative symbols.

Symbol	Meaning
$D_e$	Electronic well-depth (see Fig. 1)
$E_c$	Correlation energy
$E_x$	Exchange energy [defined in Eq. (22)]
$e_{xc}^{unif}$	Exchange-correlation energy density of uniform gas
$F[n]$	Hohenberg-Kohn universal functional (12)
$n_-(\mathbf{r})$	Density of spin – electrons
$R_0$	Equilibrium bond length (see Fig. 1)
$T$	True kinetic energy [expectation of Eq. (3)]
$U$	Hartree energy [defined in Eq. (8)]
$V_{ee}$	Electron–electron repulsion [expectation of Eq. (4)]
$V_s(\mathbf{r})$	KS potential (20)
$Z$	Atomic charge

the kernel to be extracted from a ground-state XC approximation. Typical results give moderate accuracy for transition frequencies (0.4 eV errors), but excellent results for the properties of molecules in excited states.<sup>[36]</sup> Dramatic failures of these approximations include double excitations, charge transfer, and a missing contribution to the optical response of solids.<sup>[37]</sup>


## Useful Resources

Tables 2 and 3 give a list of acronyms and notation used in this essay. Perhaps the best pedagogical introduction of recent years is ‘A Primer in DFT.’<sup>[38]</sup> For physicists with some background in many-body theory, there’s nothing better than Dreizler and Gross,<sup>[39]</sup> or the more recent Engel and Dreizler.<sup>[40]</sup> Analogously, if you are a chemist with knowledge of quantum chemical methods, try Parr and Yang.<sup>[41]</sup> A more pragmatic approach is that of Koch and Holthausen<sup>[42]</sup> who, in Part B, discuss the accuracy of different approximate functionals for different molecular properties. For TDDFT, the standard reference is that edited by Marques,<sup>[37]</sup> and there is a new introduction by Ullrich.<sup>[43]</sup>

To check you have followed the essay, you should certainly try the questions posed in the text, which are collected and answered in the Supporting Information.

**Keywords:** density functional theory, electronic structure, local density approximation

How to cite this article: K. Burke, LO. Wagner, *Int. J. Quantum Chem.* **2013**, *113*, 96–101. DOI: 10.1002/qua.24259

 Additional Supporting Information may be found in the online version of this article.

- [1] M. Born, R. Oppenheimer, *Ann. Phys.* **1927**, *389*, 457.
- [2] J. C. Slater, *Phys. Rev.* **1929**, *34*, 1293.
- [3] D. R. Hartree, W. Hartree, *Proc. R. Soc. London. Ser. A – Math. Phys. Sci.* **1935**, *150*, 9.
- [4] A. Szabo, N. S. Ostlund, *Modern Quantum Chemistry*; MacMillan: New York, **1982**.
- [5] R. M. Martin, *Electronic Structure: Basic Theory and Practical Methods*; Cambridge University Press, **2004**.
- [6] L. H. Thomas, *Math. Proc. Camb. Phil. Soc.* **1927**, *23*, 542.

- [7] E. Fermi, *Z. Phys. A: Hadrons Nucl.*, **1928**, 48, 73.
- [8] E. Teller, *Rev. Mod. Phys.* **1962**, 34, 627.
- [9] M. Levy, *Proc. Natl. Acad. Sci. USA* **1979**, 76, 6062.
- [10] E. H. Lieb, *Int. J. Quantum Chem.* **1983**, 24, 243.
- [11] P. Hohenberg, W. Kohn, *Phys. Rev.* **1964**, 136, B864.
- [12] U. von Barth, L. Hedin, *J. Phys. C: Solid State Phys.* **1972**, 5, 1629.
- [13] W. Kohn, L. J. Sham, *Phys. Rev.* **1965**, 140, A1133.
- [14] C. J. Umrigar, X. Gonze, *Phys. Rev. A* **1994**, 50, 3827.
- [15] S. Kümmel, L. Kronik, *Rev. Mod. Phys.* **2008**, 80, 3.
- [16] P. A. M. Dirac, *Math. Proc. Cambridge Philos. Soc.* **1930**, 26, 376.
- [17] J. P. Perdew, Y. Wang, *Phys. Rev. B* **1992**, 45, 13244.
- [18] (a) J. P. Perdew, K. Burke, M. Ernzerhof, *Phys. Rev. Lett.* **1996**, 77, 3865;  
(b) J. P. Perdew, K. Burke, M. Ernzerhof, *Phys. Rev. Lett.* **1997**, 78, 1396(E).
- [19] K. Burke, M. Ernzerhof, J. P. Perdew, *Chem. Phys. Lett.* **1997**, 265, 115.
- [20] A. D. Becke, *Phys. Rev. A* **1988**, 38, 3098.
- [21] C. Lee, W. Yang, R. G. Parr, *Phys. Rev. B* **1988**, 37, 785.
- [22] A. D. Becke, *J. Chem. Phys.* **1993**, 98, 5648.
- [23] R. Ahlrichs, M. Bär, M. Häser, H. Horn, C. Kölmel, *Chem. Phys. Lett.* **1989**, 162, 165.
- [24] M. Ernzerhof, G. E. Scuseria, *J. Chem. Phys.* **1999**, 110, 5029.
- [25] D. Rappoport, N. R. M. Crawford, F. Furche, K. Burke, Computational Inorganic and Bioinorganic Chemistry, In Which functional should I choose? E. I. Solomon, R. B. King, R. A. Scott, Eds.; John Wiley & Sons, Inc., **2009**.
- [26] M. Dion, H. Rydberg, E. Schröder, D. C. Langreth, B. I. Lundqvist, *Phys. Rev. Lett.* **2004**, 92, 246401.
- [27] J. P. Perdew, A. Ruzsinszky, G. I. Csonka, L. A. Constantin, J. Sun, *Phys. Rev. Lett.* **2009**, 103, 026403.
- [28] J. Heyd, G. E. Scuseria, *J. Chem. Phys.* **2004**, 121, 1187.
- [29] A. Nilsson, L. G. M. Pettersson, J. Nørskov, Eds. Chemical Bonding at Surfaces and Interfaces; Elsevier: Amsterdam, **2008**.
- [30] R. Car, M. Parrinello, *Phys. Rev. Lett.* **1985**, 55, 2471.
- [31] S. Root, R. J. Magyar, J. H. Carpenter, D. L. Hanson, T. R. Mattsson, *Phys. Rev. Lett.* **2010**, 105, 085501.
- [32] E. Runge, E. K. U. Gross, *Phys. Rev. Lett.* **1984**, 52, 997.
- [33] C. J. Umrigar, A. Savin, X. Gonze, Are Unoccupied Kohn-Sham Eigenvalues Related to Excitation Energies? Plenum: New York, **1997**.
- [34] A. Kono, S. Hattori, *Phys. Rev. A* **1984**, 29, 2981.
- [35] K. Burke, J. Werschnik, E. K. U. Gross, *J. Chem. Phys.* **2005**, 123, 062206.
- [36] P. Elliott, F. Furche, K. Burke, Excited States from Time-Dependent Density Functional Theory; Wiley: Hoboken, NJ, **2009**; pp. 91–165.
- [37] M. A. L. Marques, N. T. Maitra, F. M. S. Nogueira, E. K. U. Gross, A. Rubio, Eds. Fundamentals of Time-Dependent Density Functional Theory, *Lecture Notes in Physics*, Vol 837; Springer: Heidelberg, **2012**.
- [38] C. Fiolhais, F. Nogueira, M. Marques, A Primer in Density Functional Theory; Springer-Verlag: New York, **2003**.
- [39] R. M. Dreizler, E. K. U. Gross, Density Functional Theory: An Approach to the Quantum Many-Body Problem; Springer-Verlag: Berlin, **1990**.
- [40] E. Engel, R. M. Dreizler, Density Functional Theory: An Advanced Course; Springer: Berlin, **2011**.
- [41] R. G. Parr, W. Yang, Density Functional Theory of Atoms and Molecules; Oxford University Press: Oxford, **1989**.
- [42] W. Koch and M. C. Holthausen, A Chemist's Guide to Density Functional Theory, 2nd ed.; Wiley-VCH: Weinheim, **2002**.
- [43] C. A. Ullrich, Time-Dependent Density-Functional Theory; Oxford University Press: Oxford, **2012**.

Received: 25 May 2012  
Revised: 25 May 2012  
Accepted: 4 June 2012  
Published online on 12 July 2012

# Erratum: DFT in a Nutshell

Kieron Burke<sup>[a,b]</sup> and Lucas O. Wagner<sup>\*[a,b]</sup>

Published in *International Journal of Quantum Chemistry* **2013**, 113, 96–101.

The authors notified us of a number of errors as included below. We apologize for any inconvenience this may have caused.

- On Page 4, equation 23 should read as:

$$E_{XC}^{LDA} = \int d^3r e_{XC}^{unif}(n_{\uparrow}(\mathbf{r}), n_{\downarrow}(\mathbf{r}))$$

- On Page 4, second column, second sentence should read as,

"... using information from only the slowly varying electron **gas** and known exact conditions on the functional."

(change in bold text)

- On Page 5, second column, in Table 3 (Notation of representative symbols), the sixth definition should read as:

$n_{\downarrow}$  Density of spin  $\downarrow$  electrons.

(This is done to be consistent with the usage in the paper.)

- On Page 5, second column, the missing Acknowledgment section (before Keywords) should read as:

"The authors kindly thank the Department of Energy for support through grant DE-FG02-08ER46496."

---

Received: 14 July 2012  
Accepted: 16 July 2012  
Published online 26 February 2013  
DOI: 10.1002/qua.24411

---

[a, b] K. Burke, L. O. Wagner  
Department of Chemistry, University of California, Irvine, California 92697  
Department of Physics, University of California, Irvine, California 92697  
E-mail: lwagner@uci.edu