


# Lesson 5

## Cellular Chemistry, Reaction Thermodynamics and Enzymes



# Metabolism

- The cell is an ideal continuous chemical reactor
- The ensemble of cellular chemical reactions = **metabolism**
- **Anabolic reactions** = build up molecules (*e.g.*, condensation)




- **Catabolic reactions** = break down molecules (*e.g.*, hydrolysis)




Factory, Maurice Utrillo, 1923

# Condensation & hydrolysis



## Condensation:


Covalent bonds form  
with loss of water  
Build molecules  
Anabolic



## Hydrolysis:

Covalent bonds break  
with addition of water  
Break down molecules  
Catabolic


# Metabolism & Energy



# The metabolic map

Dots = molecules

Lines = chemical reactions  
making up the metabolism of  
the cell



# Free energy difference

- Reactions are governed by **free energy** (usable energy) **G**

Reagents  $\rightleftharpoons$  Products

- What really matters is the **free energy difference**  $\Delta G = \sum G_P - \sum G_R$
- $\Delta G$  stems from a fundamental law of thermodynamics


$$\Delta G = \Delta H - T\Delta S$$

- **H** = **Enthalpy** = total energy
- **T** = **Temperature**
- **S** = **Entropy** = useless energy




# Free energy difference

- $\Delta G < 0$  ( $G_P < G_R$ )  $\rightarrow$  Energy released, reaction proceeds (**thermodynamically spontaneous, exergonic**)



# Free energy difference

- $\Delta G < 0$  ( $G_P < G_R$ )  $\rightarrow$  Energy released, reaction proceeds (thermodynamically spontaneous, exergonic)
- $\Delta G > 0$  ( $G_P > G_R$ )  $\rightarrow$  Reactions requires energy to proceed (thermodynamically non-spontaneous, endergonic)



# Free energy difference

- $\Delta G < 0$  ( $G_P < G_R$ )  $\rightarrow$  Energy released, reaction proceeds (thermodynamically spontaneous, exergonic)
- $\Delta G > 0$  ( $G_P > G_R$ )  $\rightarrow$  Reactions requires energy to proceed (thermodynamically non-spontaneous, endergonic)
- **$\Delta G = 0$**  ( $G_P = G_R$ )  $\rightarrow$  Chemical equilibrium ( $R \rightarrow P = P \rightarrow R$ )



$$K_{eq} = [B]/[A]$$

$$\Delta G = -RT \ln [B]/[A]$$

┆

$$\Delta G = -RT \ln K_{eq}$$


$$R = 8.314 \text{ J}/(\text{mol K}) = 1.987 \text{ cal}/(\text{mol K})$$

# Going to work in the cell factory

- Cells transfer energy from food molecules to run their processes
  - **Synthesis**
    - Complex molecules (DNA, proteins)
    - Organized structures (organelles)
  - **Organization**
    - Generation of specialized compartments to store and organize materials for specific tasks


# Going to work in the cell factory

- Cells transfer energy from food molecules to run their processes
  - Synthesis
    - Complex molecules (DNA, proteins)
    - Organized structures (organelles)
  - Organization
    - Generation of specialized compartments to store and organize materials for specific tasks
- **Creation of electrochemical gradients**
  - Cells organize ions on either cell membrane sides to create electrochemical potentials
    - Mechanism that controls signaling and transfer of materials across the membranes




# Going to work in the cell factory

- Cells transfer energy from food molecules to run their processes
  - Synthesis
    - Complex molecules (DNA, proteins)
    - Organized structures (organelles)
  - Organization
    - Generation of specialized compartments to store and organize materials for specific tasks
  - Creation of electrochemical gradients
    - Cells organize ions on either cell membrane sides to create electrochemical potentials
      - Mechanism that controls signaling and transfer of materials across the membranes
  - **Transport and movement**
    - Cells transport molecules from one side of the cell to another
    - Some cells swim or crawl using special motility proteins



# Where does cell energy really come from?

- Cells do not use the energy from food directly
- Adenosine triphosphate, the energy middleman




# The ATP/ADP cycle

Simple molecules  
glucose, amino acids, fatty acids

Catabolism – Exergonic reactions  
transfer energy from complex  
molecules to ATP


Complex molecules  
polycarbs, proteins, lipids

Anabolism – Endergonic reactions  
transfer energy from ATP  
to complex molecules




# The ATP/ADP cycle – energy coupling

- Cells make and break ATP all the time
- In endergonic reactions
  - Energy is provided by ATP hydrolysis to ADP and  $P_i$
- In exergonic reactions
  - Energy is captured to restore ATP by ADP and  $P_i$  condensation




# As easy as 1,2,3 – ATP, ADP and AMP



$$\Delta G = -30.5 \text{ kJ/mol} = 7.29 \text{ kcal/mol}$$




$$\Delta G = -30.5 \text{ kJ/mol} = 7.29 \text{ kcal/mol}$$





# Free energy difference

- Even if a reaction is thermodynamically spontaneous ( $\Delta G < 0$ ), it may not occur
  - It needs an “energetical push”
- **Activation energy  $E_a$  = energy barrier**





# Enzymes

- Even if a reaction is thermodynamically spontaneous ( $\Delta G < 0$ ), it may not occur
  - It needs an “energetical push”
- Activation energy  $E_a$  = energy barrier
- **Catalysts** = particular class of chemical substances that lower  $E_a$  and promote reactions
- **Enzymes** = biological catalysts (mostly proteins)




# Enzymes

- There are approximately 1300 different enzymes found in the human cell
- Each enzyme catalyzes a specific chemical reaction
- **ENZYMES DO NOT CHANGE THE  $\Delta G$  OF A REACTION BUT JUST SPEED UP THE REACTION RATE**
  - By breaking down the reaction into different steps, each of which has a low  $E_a$




# Enzymes



ENZYME CYCLE  
(reversible)


# Enzyme specificity

Cellulose and starch  
are both glucose polymers



Cellulase  
breaks  $\beta$ -1,4 bonds in cellulose  
People do not have this enzyme  
so we cannot digest grass!



Amylase  
breaks  $\alpha$ -1,4 bonds in starch  
People have this enzyme!



ENZYME SPECIFICITY


# Metabolic pathways

- Cellular chemical reactions all catalyzed by enzymes
- Cellular chemical reactions are often linked into **pathways**
  - Ordered sequence of chemical reactions
- Pathways = the “cell production line”
  - Starting from point A → land up with a particular product that the cell really needs
- Reactions are organized into multistep pathways
- Cellular pathways are governed by internal feedback mechanisms
  - **Positive feedback** = make more product(s) along that particular pathway
  - **Negative feedback** = make less and/or stop producing product(s) along that particular pathways
- Cellular pathways can also be mastered by external control signals




# Positive feedback loops

## Blood clot formation




- When the body gets injured the major threat to life is excessive loss of blood
- Blood pressure and blood flow at the site of injury are reduced
- At the site of the injury, blood clotting factors are released to initiate blood clotting
  - Clotting factors are responsible for the formation of a clot in the injured or wounded area
- Once the process begins it promotes the clotting process further → overall, the process of sealing the injured site is speeded up
- **This is one of the life-saving examples of positive feedback**



# Positive feedback loops


## Childbirth



- The onset of contractions in childbirth is also known as the *Ferguson reflex*
- The baby pushes against cervix, causing it to stretch
- Stretching of cervix causes nerve impulses to be sent to the brain
- Brains stimulates the posterior pituitary gland to to release oxytocin
- Oxytocin stimulates the uterine muscle to contract, initiating the birth process
- The fetus responds to uterine contractions by releasing prostaglandins, which triggers further uterine contractions


# Negative feedback loops

## Childbirth




- The onset of contractions in childbirth is also known as the *Ferguson reflex*
- The baby pushes against cervix, causing it to stretch
- Stretching of cervix causes nerve impulses to be sent to the brain
- Brains stimulates the posterior pituitary gland to to release oxytocin
- Oxytocin stimulates the uterine muscle to contract, initiating the birth process
- The fetus responds to uterine contractions by releasing prostaglandins, which triggers further uterine contractions

# Cellular pathways (examples)



## Cell Cycle Control: G1/S Checkpoint



## Cell Cycle Control: G2/M DNA Damage Checkpoint

