


Xist spatially amplifies SHARP/SPEN recruitment to balance chromosome-wide silencing and specificity to the X chromosome

Joanna W. Jachowicz, Mackenzie Strehle, Abhik K. Banerjee, Mario R. Blanco, Jasmine Thai and Mitchell Guttman




X inactivation




↓
Double dosing of X gene is incompatible with early development

↓
One of the X chromosome is **silenced** by heterochromatization



The X inactivation phenomena

- Choice of X




The X inactivation phenomena

- THE X CHOICE MECHANISM

→ Random

→ Conserved in the cell's progeny



- MOSAICISM PHENOMENA


→ Visual example of X choice



Mechanism of X silencing

Xist :

- Coded by X chromosome
- Non coding RNA 15000 nt
- Located all over the X chromosome



SHARP :

- Protein 400 kDA - 3664 AA
- 4 RNA reconizion motif
- Domain C-terminal



RNA recognition motif


Intransically disordered region

SPEN paralog and ortholog C-terminal

Mechanism of X silencing

1. Initiation


- X inactivation center



Mechanism of X silencing

1. Initiation


- Race to inactivation
- Concentration of X linked activation factor



Mechanism of X silencing



2. Establishment

- Xist transcription in X inactivation center
- Sharp recruitment by Xist
- Sharp recruit proteins complexes



Mechanism of X silencing

3. Maintenance




- DNA methylation
- Xist continuous expression

→ Silences the chromosome with **continued expression of Xist**

→ Xi **maintains its configuration** and **propagates** by cloning in all his divisions

Some genes escape the silencing




From : The Molecular and Nuclear Dynamics of X-Chromosome Inactivation

- 15 to 30% escape inactivation
- Xist coating responsible




Global inactivation



And if it's go wrong ?

If it didn't have X inactivation → Abortion

But some **diseases** can be in link with an **abnormal inactivation**



→ Code for 80% of immunity gene




Many autoimmune disorders have a female bias

Current studies on Xist

→ Influence of mosaicism

→ Reactivating the other x in case of x linked disease



From : Amitesh Panda et al, Cell, 2020,

PROBLEMATIQUE

- A greater understanding of those phenomenon at the molecular level is needed.
- All the complexes necessary to x silencing are recruited by Xist.
- But you have only 1 Xist mol for every megabase of DNA that need silencing

How does x silencing hapend in those condition ?



Xist spatially amplifies SHARP/SPEN recruitment to balance chromosome-wide silencing and specificity to the X chromosome


Joanna W. Jachowicz ^{1,3}, Mackenzie Strehle ^{1,3}, Abhik K. Banerjee^{1,2}, Mario R. Blanco¹, Jasmine Thai¹ and Mitchell Guttman ¹✉

To better understand how Xist works, the article address five main questions :

- 1. How do Xist recruit sharp ?**
- 2. How do SHARP assemblies form ?**
- 3. How does SHARP recruitment hapen?**
- 4. Does SHARP foci play a role in X silencing?**
- 5. Why is this strategy necessary?**


Methods

Experimental model



Femal F1 hybrids (Bl6xCast)
Mouse embryonic stem cells




Induction of Xist by dox



1. How do Xist recruit sharp ?

Methods

HaloTag construction



$$\text{SHARP} / \text{Xist} = 1 \text{ (constant)}$$



↓

Recruitment is stoichiometric

1. How do Xist recruit sharp ?


Result

RNA-fish on mouse mES



→ Stoecheometric excess

2. How do sharp assemblies form ?



→ Sharp form concentration dependant assemble in the nuclues

2. How do sharp assemblies form ?

Sharp deletion analysis :


FUS = Form multivalent homotypic interactions

Methods

Different models




Deletion



Δ IDR



Rescue



FUS- Δ IDR



Fluorescence technique

FL-SHARP



Δ IDR-SHARP




FUS- Δ IDR-SHARP




2. How do sharp assemblies form ?

Result

IDR deletion




Rescue (FUS)



→ Foci formation depend on homotypic multivalent interaction between IDRs

3. How does Sharp recruitment happen?

Methods



Visualisation by RNA Fish


Direct visualisation

3. How does Sharp recruitment happen?

Methods


→ Ezh2 use to mark Xi

→ Immunomarcage antibody against Ezh2



3. How does Sharp recruitment happen?

Result



3. How does Sharp recruitment happen?

Result


	Full-length <i>RRM</i> <i>IDR</i> <i>SPOC</i>	Δ RRM	Δ IDR	FUS- Δ IDR <i>FUS</i>
Sharp located on Xi ?	+++	-	+++	+++
Sharp can enrich Xi ?	+++	-	-	+++

→ RRM of SHARP are essential for localisation / link to Xi

→ IDR of SHARP are essential for enrichment on the Xi



4. Does SHARP foci play a role in X silencing?

Methods



4. Does SHARP foci play a role in X silencing?



Result



No inactivation of Xist without SHARP


4. Does SHARP foci play a role in X silencing?

Result



➔ No inactivation of Xist without RRM & IDR domain of sharp

Conclusion diagram of Sharp recruitment on Xist



Sharp
recruitment by
Xist

Excess
stoichiometric
recruitment

- RRM s allow Sharp linking on
Xist
- Interaction between IDRs
allow SHARP non
stoichiometric recruitment



Responsible of X
silencing

5. Why is this strategy necessary?

→ **Low number of Xist** mol is critical mechanical by which cells limit Xist spreading to autosomal region ?

Methods : **RNA - Fish quantification**


5. Why is this strategy necessary?



→ Xist quantity is correlate with Xist territory


→ Xist is in low quantity so as not to fix the autosomes

5. Why is this strategy necessary?



Perspectives ?

Discussion/conclusion




Do you have any question ?










Bibliography

- Wang W, Min L, Qiu X, Wu X, Liu C, Ma J, Zhang D, Zhu L. Biological Function of Long Non-coding RNA (LncRNA) Xist. *Front Cell Dev Biol.* 2021 Jun 10;9:645647. doi: 10.3389/fcell.2021.645647. PMID: 34178980; PMCID: PMC8222981.
- Furlan G, Galupa R. Mechanisms of Choice in X-Chromosome Inactivation. *Cells.* 2022 Feb 3;11(3):535. doi: 10.3390/cells11030535. Erratum in: *Cells.* 2023 Mar 21;12(6): PMID: 35159344; PMCID: PMC8833938.
- Dossin F, Heard E. The Molecular and Nuclear Dynamics of X-Chromosome Inactivation. *Cold Spring Harb Perspect Biol.* 2022 May 17;14(4):a040196. doi: 10.1101/cshperspect.a040196. PMID: 34312245; PMCID: PMC9121902.
- Maduro C, de Hoon B, Gribnau J. Fitting the Puzzle Pieces: the Bigger Picture of XCI. *Trends Biochem Sci.* 2016 Feb;41(2):138-147. doi: 10.1016/j.tibs.2015.12.003. Epub 2016 Jan 13. PMID: 26796628.
- Giaimo BD, Robert-Finestra T, Oswald F, Gribnau J, Borggrefe T. Chromatin Regulator SPEN/SHARP in X Inactivation and Disease. *Cancers (Basel).* 2021 Apr 1;13(7):1665. doi: 10.3390/cancers13071665. PMID: 33916248; PMCID: PMC8036811.

Annexes :




Annexes :



Annexes :

a



h

