



UNIVERSITÀ DEGLI STUDI DI TRIESTE

Dipartimento di Matematica e Geoscienze

Corso di Geologia Marina 2015-16



Università di Trieste
LAUREA MAGISTRALE IN GEOSCIENZE
Curriculum Geofisico
Curriculum Geologico Ambientale

Anno accademico 2015 – 2016

Geologia Marina


Parte II

Modulo 2.3 Metodi diretti: Sondaggi superficiali ed analisi dei sedimenti

Relatore
Dr. Renata G. Lucchi
rglucchi@ogs.trieste.it




BOTTOM SAMPLING SYSTEMS




Gravity corer

- classic gravity corer (Emery and Dietz, 1941; Hvorslev and Stetson, 1946)
- box corer
- kastenlot corer
- multi-corer



Piston corer

- classic piston corer (Kullenberg, 1947; 1955)
- long piston corer
 - Calypso piston corer (e.g. R/V Marion Dufresne, G.O. Sars)
 - Jumbo piston corer (e.g. R/V Araon)




Drilling systems



- ocean floor drilling systems (e.g. IODP-drilling vessels & semi-automated MeBo system)
- ice drilling systems (e.g. EPICA-European Project for Ice Coring in Antarctica & NorthGRIP- North Greenland Ice Core Project)



Classic gravity core system (Emery and Dietz, 1941)




It is the simplest coring device in which the weight of the coring equipment is used to force the barrel into the sea bottom. This system can work with or without a triggering system (sistema di sgancio).






Classic gravity core system: corer head



weight 600-800 kg
6000 kg



Classic gravity core system: core catcher and cutter



core cutter
(scarpa)





core cutter
with sediments




Additional gravity core systems: Kastenlot corer


The kastenlot (kastens) corer was originally designed by Kögler (1963) it was improved and modified by Zanger and McCave (1990). The barrel, of variable lengths, is square in section (15x15 cm) and it contains a base plate that can be raised to reveal a new cleaned core surface



Square section
of barrel



Core cutter and catcher



Head of
kastens
corer




The core catcher has a shutter-like closure consisting of two square doors held under tension and blocked in a retracted position by two levers located on the outside. During the corer pullout, the pressure of the surrounding sediments pushes down the two levers closing the doors.



- (a) removal of the barrel lid to reveal the core surface;
- (b) two PVC trunkings are pushed into the sediments
- (c) the inner base is moved upward to expose the trunkings that are cut at the base and removed from the main core using a cheese wire;
- (d) sampling with x-ray slabs, and a further set of trunkings (e). Each time the sediments are lifted upward and withdrawn with a cheese wire

Sampling of Kastens cores





Additional gravity core systems: BOX-corer

Designed for minimum disturbance of the sediment surface, ideal for coarse/stiff sea floor sediments, it allows the recovery of bottom waters.



lateral surface



coral sampling



sampling of
glacigenic
sediments



Box core sampling






Additional gravity core systems: Multi-corer

Especially designed for the sampling of sea bottom sediments-water interface, it permits to recover low disturbance sediment. Ideal for geochemical and biological sediment and water analysis.






Kullenberg piston corer system

Standard assemblage for piston coring. The core barrel penetration is maximised by the action of a piston located in the lower barrel (or into the lower plastic liner if present) that helps to overcome the friction between sediments and the coring tube by generating Vacuum behind the cutter. The sediment cores obtained are less compacted and distorted than gravity cores. This system is always used coupled with a trigger mechanism.


| Specifications | Kullenberg piston corer | Long piston corer |
|------------------------------|-------------------------|-------------------|
| headweight | 600 kg | 6000 kg |
| barrel length | 6 m | 13 m |
| barrel inner diameter | 65 mm | 140 mm |
| barrel thickness | 5 mm | 5 mm |
| plastic liner outer diameter | 63 mm | 113 mm |
| plastic liner thickness | 3 mm | 5 mm |
| maximum cable length | 5000 m | 10000 m |
| cable diameter | 12 mm | 30 mm |
| freefall | 4-5 m | 1.5 m |





Long piston corers:
Claypso, Jumbo etc.



Grab (benna)



Dredge (draga)





UNIVERSITÀ DEGLI STUDI DI TRIESTE


Dipartimento di Matematica e Geoscienze

Corso di Geologia Marina 2015-16



Fotos: Volker Diekamp, Marum

CORE ON DECK!






UNIVERSITÀ DEGLI STUDI DI TRIESTE

Dipartimento di Matematica e Geoscienze

Corso di Geologia Marina 2015-16




CUT INTO SECTIONS





SECTIONS' LABELING





The plastic liner is extracted from the barrel and cut into sections 1-1.5 m-long

The sections are numbered consecutively from bottom to top of the core or *vice versa* depending on the total length of the core.

Short cores are numbered consecutively from top to bottom.

Long cores are numbered consecutively on removal from the barrel from bottom to top of core.

Each section is labeled with a code indicating the name of the project and/or the name of the research vessel; the core number; the section number, and stratigraphic orientation (top-bottom)





CORE OPENING AND SEDIMENTS ANALYSIS


The plastic liner of each section is cut longitudinally. The plastic liner is cut by means of an electric saw/microvibro saw, while the sediments are cut using a “cheese wire”. The two splitted half-sections are labeled as **working section** and **archive section** and will undergo a different analytical process:

ARCHIVE SECTIONS: not destructive analyses
X-radiographs
multi-sensor core logger
XRF core-scan
photographs

WORKING SECTIONS: visual logging and sub-sampling



Half-sections'
labeling





CORE OPENING AND SEDIMENTS ANALYSIS


The plastic liner of each section is cut longitudinally. The plastic liner is cut by means of an electric saw/microvibro saw, while the sediments are cut using a “cheese wire”. The two splitted half-sections are labeled as **working section** and **archive section** and will undergo a different analytical process:

ARCHIVE SECTIONS: not destructive analyses
X-radiographs
multi-sensor core logger
XRF core-scan
photographs

WORKING SECTIONS: visual logging and sub-sampling



Half-sections' labeling





UNIVERSITÀ DEGLI STUDI DI TRIESTE

Dipartimento di Matematica e Geoscienze

Corso di Geologia Marina 2015-16



CORE DESCRIPTION






Visual core description 1° step



- **Definition of lithological units** based on:
 - Lithology including composition and texture
 - Color
 - Sedimentary structures (laminations, bioturbations, faults)
 - Boundaries (transitional, sharp not erosive, sharp erosive, irregular)
- **Definition of sediment disturbance**
 - Soupy sediments
 - Bended boundaries at the lateral ends
 - Flow-in (piston cores)
 - Core re-bouncing (repetition of stratigraphic sections) in gravity cores



| Project: | | Date | CORE | SECTION | |
|-----------------|---------|------------------|---------------------------------------|-----------|---------------------------|
| Observer(s) ... | samples | | LITHOLOGY (cm from top of section) | texture | LITHOLOGIC DESCRIPTION |
| depths | lengths | SEISMIC DEPTH | size | thickness | value |
| bottom | surface | 0 | 0 | 0 | 0 |
| 1 | 2 | 3 | 4 | 5 | 6 |
| 7 | 8 | 9 | 10 | 11 | 12 |
| 13 | 14 | 15 | 16 | 17 | 18 |
| 19 | 20 | 21 | 22 | 23 | 24 |
| 25 | 26 | 27 | 28 | 29 | 30 |
| 31 | 32 | 33 | 34 | 35 | 36 |
| 37 | 38 | 39 | 40 | 41 | 42 |
| 43 | 44 | 45 | 46 | 47 | 48 |
| 49 | 50 | 51 | 52 | 53 | 54 |
| 55 | 56 | 57 | 58 | 59 | 60 |
| 59 | 60 | 61 | 62 | 63 | 64 |
| 65 | 66 | 67 | 68 | 69 | 70 |
| 71 | 72 | 73 | 74 | 75 | 76 |
| 77 | 78 | 79 | 80 | 81 | 82 |
| 83 | 84 | 85 | 86 | 87 | 88 |
| 89 | 90 | 91 | 92 | 93 | 94 |
| 95 | 96 | 97 | 98 | 99 | 100 |

LEGEND

- day
- shallow boundary
- transitional boundary
- irregular boundary
- normal and reverse
- planar laminations
- cross laminations
- shells fragments
- pebbles rich
- trans. rich
- shaly sediment
- clay
- dark layer
- fault
- V.V. = very
- S. = soft
- D. = dark

length of section (cm)
total length of core (cm)
remarks:





Visual core description FORM






| Project: | | Date | CORE | SECTION |
|--|---------------------------------------|--|---------------------------------|---------|
| Observer(s) | LITHOLOGY (cm from top of section) | | | |
| samples | SEDIM. STRUCT. | texture ^(a) clay fine medium coarse grains-pebbles | LITHOLOGIC DESCRIPTION | |
| forams nannos diatoms s. slides | | | COLOUR | |
| | | | 10 | |
| | | | 20 | |
| | | | 30 | |
| | | | 40 | |
| | | | 50 | |
| | | | 60 | |
| | | | 70 | |
| | | | 80 | |
| | | | 90 | |
| | | | 100 | |
| LEGEND | | | | |
| - - - clay | — sharp boundary | ◇ pteropod rich | length of section (cm) | |
| - - - silt | — gradational boundary | ⊗ forams rich | total length of core (cm) | |
| - - - sand | ~~~ angular boundary | ● shell's fragments | remarks: | |
| pebbles | ▲▼ normal and reverse grading | ○ soupy sediment | | |
| turbidite | planar laminations | ◎ bioturbation | | |
| V V V tephra | cross laminations | | | |
| — dark layer | slump | | | |
| | fault | | | |



UNIVERSITÀ DEGLI STUDI DI TRIESTE


Dipartimento di Matematica e Geoscienze


Corso di Geologia Marina 2015-16





X-RAY

SEDIMENT
SURFACE







UNIVERSITÀ DEGLI STUDI DI TRIESTE

Dipartimento di Matematica e Geoscienze

Corso di Geologia Marina 2015-16



CORING DISTURBANCE






Visual core description 2° step sediment composition

Mud composition - Smear slides

- A smear slide is a thin layer of unconsolidated sediment embedded on a glass slide for petrographic microscopic examination;
- Smear slides are a useful tool to quickly assess the ***compositional content*** of clay-silty sediment samples;
- Smear slides are a powerful method for rapidly evaluating tiny quantities of sediment (mineralogy, components, form, size) as the basis for ***sediment classification***, and for ascertaining the presence of microfossils.





Foto: IODP Exp. 307






Quarz




IODP Exp .320


Plagioclase



Clay fraction




Composition/Quantification



Rothwell R.G., 1988. Minerals and mineraloids in marine sediments. An Optical Identification Guide. Elsevier Science Publishers, 279 pp



Diatoms




オホーツク海・中心部、水深 1107m, XP98, PC1, Sec.1, 20cm
200 倍。スケールは横 100 μm

Foraminifers



太平洋・熊野トラフ、水深 2190m, KT02-1, KK2PC
400 倍。スケールは横 50 μm

Calcareous nannoplankton



東地中海・キプロス沖、ODP Leg160, 967D, 1H, Sec.1, 0-5cm
1000 倍。スケールは横 20 μm

IODP Exp .320



Sand composition

- wet sievings at 63 microns are used to investigate the composition of sediment coarse fraction, the grains form and roundness, in order to define the sediment provenance. The sand fraction is also used for biostratigraphic purposes.



Beach sands



Glacigenic sediments

Tephra
(volcanic glass)



SEDIMENT SAMPLE DESCRIPTION

Project:

lat. N long. E water depth (m)
 core section cm cm from core top

sediment type lithology
 colour remarks

DESCRIPTION OF RESIDUE**textural characteristics**

| | | | | | |
|---------|----------------------------|------------|----------------------|------------|----------------------------------|
| sorting | very good good | sphericity | high low | angularity | very angular angular |
| | moderately | | | | sub-angular/rounded |
| | poor very poor | | | | rounded well rounded |

remarks

residue composition

(AA=very abundant; A=abundant; O=occurring; R=rare; RR=very rare)

terrigenous components: predominant not predominant

quartz calcite K-feldspar gypsum glauconite
 mica pyrite dark minerals volcanic glass micronodules
 rock fragments others

bioclastic components: predominant not predominant

benthic forams planktonic forams pteropods bryozoa
 echinoid spines sponge spicules ostracods corals
 bivalves gastropods corals algae
 radiolarians diatoms spores fish teeth
 plant debris coccoliths discoasters organic matter
 micrascidites of tunicates others

Foraminifera: list of the most characteristic taxa

| | | | |
|-----------------------|----------------------------|-----------------------|-----------------------------|
| <i>G. bulloides</i> | <i>G. tenellus</i> | <i>H. siphonifera</i> | <i>G. coriaconensis</i> |
| <i>G. trilobus</i> | <i>N. eggeri dutterrei</i> | <i>G. quinqueloba</i> | <i>G. inflata</i> |
| <i>N. pachyderma</i> | <i>G. ruber</i> | <i>G. scitula</i> | <i>O. universa</i> |
| <i>G. gomitiulus</i> | <i>G. sacculifer</i> | <i>G. glutinata</i> | <i>G. truncatulin. exc.</i> |
| <i>G. conglobatus</i> | <i>G. quadrilobatus</i> | <i>H. pelagica</i> | <i>G. digit./preadigit.</i> |
| <i>S. ionica</i> | others | | |

AGE
 ZONE
 FACIES
 REMARKS

SMEAR SLIDES DESCRIPTION

Project:

lat. N long. E water depth (m)
 core section cm cm from core top

sediment type lithology
 colour remarks

DESCRIPTION OF SMEAR SLIDES

Calcareous nannofossil: list of the most characteristic taxa and their abundance
 (AA=very abundant; A=abundant; O=occurring; R=rare; RR=very rare)

| | | | |
|---------------------|----------------------|------------------------|---------------------------|
| <i>E. huxleyi</i> | <i>G. oceanica</i> | <i>G. caribbeanica</i> | <i>small Gephyrocapsa</i> |
| <i>H. carteri</i> | <i>C. leptoporus</i> | <i>S. pulchra</i> | <i>S. histrica</i> |
| <i>C. pelagicus</i> | <i>S. recurvata</i> | <i>P. scutellum</i> | <i>O. antillarum</i> |
| <i>R. clavigera</i> | <i>R. stylifer</i> | <i>S. fossilis</i> | <i>B. bigelowi</i> |
| <i>U. tenuis</i> | <i>C. jonesii</i> | <i>T. saxeae</i> | <i>C. rugosus</i> |
| <i>C. cristatus</i> | <i>P. multipora</i> | <i>P. lacunosa</i> | <i>C. macintyreai</i> |
| <i>D. broweri</i> | <i>H. sellii</i> | <i>Discoaster sp.</i> | |

others

Sediment composition

(AA=very abundant; A=abundant; O=occurring; R=rare; RR=very rare)

terrigenous component: predominant not predominant

quartz calcite K-feldspar gypsum glauconite
 mica dark minerals pyrite volcanic glass micronodules
 dolomite aragonite Fe oxides zeolite plagioclase
 rock fragments clay minerals
 others

bioclastic component: predominant not predominant

benthic forams planktonic forams pteropods bryozoa
 echinoid spines sponge spicules ostracods corals
 bivalves gastropods corals algae
 radiolarians diatoms spores fish teeth
 plant debris coccoliths discoasters fish remain
 shell fragments silicoflagellates organic matter
 others


AGE
 ZONE
 FACIES
 REMARKS






UNIVERSITÀ DEGLI STUDI DI TRIESTE

Dipartimento di Matematica e Geoscienze



Corso di Geologia Marina 2015-16



SEDIMENT CORE ANALYSES



Plot of results





CORE REPOSITORY 4°C