



triphenylphosphine




triphenylphosphite






Ph₂PCH₂CH₂PPh₂, dppe

diphenylphosphinoethane



**2,2'-bis(diphenylphosphino)-
1,1'-binaphthyl, BINAP**



For Phosphines the σ -donor and the π -acceptor characters are inversely proportional

TEP = Tolman Electronic Parameter for phosphines


For $PX_1X_2X_3$ in $Ni(CO)_3L$

$$\nu = 2056.1 + \sum_{i=1}^3 \chi_i$$


TABLE I. Selected Values of ν (cm^{-1})

L	ν	$\Delta\nu$
P(<i>p</i> -Tol) ₃	2066.7	0.1
P(<i>o</i> -Tol) ₃	2066.6	
PMe ₃	2064.1	2.4
PEt ₃	2061.7	2.5
P(<i>i</i> -Pr) ₃	2059.2	3.1
P(<i>t</i> -Bu) ₃	2056.1	


Lewis basicity of phosphines





Need to introduce the steric parameter




Tolman cone angle in $\text{Ni}(\text{CO})_3\text{L}$






(a)



(b)



10

Table 21.4 Tolman cone angles (in degrees) for selected phosphines

PF_3	104
$\text{P}(\text{OMe})_3$	107
PMe_3	118
PCl_3	125
$\text{P}(\text{OPh})_3$	127
PEt_3	132
PPh_3	145
PCy_3	169
P^tBu_3	182
$\text{P}(o\text{-tol})_3$	193

TEP vs cone angle




FIGURE 4.4 Electronic and steric effects of common P-donor ligands plotted on a map according to Tolman (ν in cm^{-1} , θ in degrees).

Hydrides (1931)




$$\text{pK}_a = 8.3$$


^1H NMR: $-50 < \delta < 0$ (ppm)



IR: $2250 - 1500 \text{ cm}^{-1}$

Neutron Diffraction (structure)


H₂ coordination (1984): *side-on*
(typically in d^6 low-spin compounds)










H_2 -to-M donation
(a)



M-to- H_2 back-donation
(b)




Oxidative Addition




NMR: $-10 < \delta < 0$
 $^1J_{\text{HD}} = 20 - 34 \text{ Hz}$
IR: H-H $2900 - 2300 \text{ cm}^{-1}$


NMR: $-50 < \delta < 0$
 $^1J_{\text{HD}} = \sim 1 \text{ Hz}$
IR: M-H $2200 - 1500 \text{ cm}^{-1}$



NMR: $^1J_{\text{HD}} = 34 \text{ Hz}$







NMR: $^1J_{\text{HD}} < 2 \text{ Hz}$




End-on (η^1)
(most common)

IR: N–N 2150 – 1900 cm^{-1}




«hydrazine»



Side-on (η^2)
(more rare)


Nitrosyl



Linear: formally NO^+

Iso-electronic to CO

$2e^-$ donor, strong π -acceptor



Bent: formally NO^-
 $2e^-$ donor, σ -donor only

Switching from NO^+ to NO^- : formal loss of two electrons